



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.


We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





600035346R

G. 76. M. 22.

Fungi. Lindley, ord. 271.



E. BIBL. RADCL.

35.

W. Front

C. 3.

4.

9.
a. 2.

~~71. 13.~~

C

1639

e.

$\frac{62}{1}$

N





MYCOLOGIA EUROPAEA

SEU

COMPLETA OMNIUM FUNGORUM IN
VARIIS EUROPÆÆ REGIONIBUS DE-
TECTORUM ENUMERATIO,

METHODO NATURALI DISPOSITA;

DESCRIPTIONE SUCCINCTA, SYNONYMIA
SELECTA ET OBSERVATIONIBUS
CRITICIS ADDITIS.

ELABORATA

A

C. H. P E R S O O N.

SECTIONO PRIMA.

CUM TABULIS XII COLORATIS.

ERLANGAE

IMPENSIBUS IOANNI IACOBI PALMII.

MDCCCXXXII.

MYCOLOGIA.

CLASSIS PRIMA.

Fungi *exosporii* thecas aut sporulas (sporangia) in receptaculo *aperto* (in statu maturo), forma vario, plerumque dilatato proferentes.

ORDO PRIMUS.

TRICHOMYCI seu fungi byssoidei, filamentosi, aut fibrosi, rigidi, rarissime subtremelloso; nunc simplices, nunc vero plus minusve intertexti, floccos pannumque referentes, passim septati. *Sporulae*, thecis non inclusae, vel sunt dispersae, vel in capitulum, spiculam, racemulum aut verticillatim aggregatae, passim seriatim dispositae, et in nonnullis, hujus ordinis, fungis omnino nullae aut saltem nondum observatae.

I. ERINEUM. *Fila simplicia (brevia), rigida, in cespitulum aggregata, attenuata, aut turbinata, vel in crustam oblitterata.*

* PHYLLERIUM, *filis tenuioribus, crassitudine aequalibus.*

1. *rubeum*, effusum mollissimum griseo-virens.

Habitat in parte superiore foliorum *Rubi corylifolii*. Cespituli appianati, forma et magnitudine varii, $\frac{1}{2}$ unciam et ultra lati. Fila tenuissima, recta.

An *Phyllum Rubi*, effusum planiusculum album, floccis subintricatis rectis cylindricis apice acuminatis. *Fries Obs. Mycolog.* 2. p. 371. junior hujus speciei aetas?

Obs. Ambiguum est vegetabile, nam et *Bedeguar* in ramis *Ruborum* passim occurrens huic quoad colorem est simile, attamen filis crassioribus et paullo rigidioribus differt.

2. *juglandinum*, cespitulis immersis appianatis majusculis dilute pallidis.

Hab. in fol. *Juglandis regiae*. Sub nomine *E. Juglandis* a nonnullis descriptum. Fila tenuia, brevissima, mollia, caespitulos efficiunt saepius lineari-quadrangulares, majusculos, nonnunquam orbiculares immersos.

*) *Erinea* vegetabilia sunt ambigua, aspora incertaeque originis, in foliis, plerumque arborum, adhuc virentibus proveniunt: *Bedeguar* rosarum aliarumque plantarum non dissimilia, nec

3. *tiliaceum*, cespitulis pulvinatis subrotundis confertis opace pallidis. *Synops: fung. p. 700.*

β. Tiliae albae, cespitulis immersis superis.

γ. roseum, subcrustaceum rubro-roseum.

Phyllerium tiliaceum. Fries Obs. Mycol. I. p. 219.

Crescit var. *α.* in *Tilia europea*, et plerumque in superficie fol. inferiore, aestate; *β.* in Hungaria lecta et mecum communicata a D. *Funck*, in pagina fol. superiore occurrit, ubi cupulas sistit; *γ.* vere collecta a D. *Chaillet* (prope Neuchâtel) etiam suprafoliacea est, sed adhuc junior plantula videtur, textura enim est subgrumosa.

4. *coryleum*, cespitulis immersis opaco-griseis subvariegatis inferis.

Rarius in fol. *coryli Avellanae* crescit.

Cespituli cupulaeformes, molles, 2-3 lin. lati, subrubro-variegati.

5. *quercinum*, cespitulis immersis laxis rufescente-pallidis nitidis.

Phyllerium quercinum. Kunze Mycol. Hefte I. p. 87.

A 2

tamen, ut illa, insectorum opera sunt. Nunc superficialia, subeffusa, nunc immersa, cupulas seu depressiones in foliis (plerumque in pagina inferiore) efformant. Colore sunt pallida, lutescentia, badia, passim rubella vel albidia in aetate praesertim juniore.

Hab. in fol. *Quercus pubescentis*. Fila compressa, intricata, mollia.

6. *Vitis*, cespitulis majusculis immersis effusisque subpulvinatis, primo albido-roseis, dein spadiceis.

Erineum *vitis*. Duval et Phyllerium *viteum* Fries.

Frequens in fol. *Vitis viniferae* et *laciniaetae*. Cespituli congesti, subconfluentes.

7. *pyrinum*, effusum sublaxum flavo ferrugineum. *Syn. fung.* p. 700.

Hab. in fol. *Pyri* parte superiore.

8. *sorbeum*, effusum laxum superficiale confluens spadiceum.

Crescit in utraque superficie fol. *Sorbi Aucupariae*. Lanam refert et habitu a consimilibus parum recedit. — Primum album, exsiccatum, flavescens. *Fries Obs.* I. p. 218.

9. *mespilinum*, irregulare effusum ex rufescente olivaceum, filis compressis membranaceis. *Decand. Flor. Gall. Suppl.* p. 13. †

Hab. in *Mespilo germanica*.

10. *ilicinum*, superficiale, primo orbiculare, dein effusum opacum umbrinum. *Decand. Flor. Gall. Suppl.* p. 14. *Funck Pl. cryptog.* n. 505.

Hab. in Europa meridionali ad fol. *quercus Ilicis*. Ab initio albidum, tunc rufum,

demum opace spadiceum, nonnunquam totam superficiem occupat.

11. *Alni incani*, superficiale subpulvinatum rufo-badium, filis tortis aequalibus.

Erineum alneum. *Nees d'Esenbeck Comment. p. 14. t. I. f. 10.*

Hab. in Helvetia. Cespituli forma et filorum structura ab *E. alneo* distinctum.

12. *Platanoidis*, filis densissime intricatis tenuissimis albido-cinereis. Phyllerium acerinum. *Fries Obs. Mycol. I. p. 218. †*

In fol. *Aceris plantanoidis*, vere; in utraque pagina crescere dicitur.

13. *Gei*, floccis subliberis tenuissimis retiusculis filiformibus albidis. *Fries l. c. p. 200. †*

In fol. *Gei rivalis*, immersum; habitu a reliquis recedit. Fr. Forte junior adhuc plantula quae in monte Kullaberg in Suecia lecta.

** GRUMARIA, filis rigidioribus, incrassatis turbinatis aut cupulatis, passim difformibus.

14. *Pseudoplatani*, subrotundum (profunde) immersum umbrinum, floccis (filis) dense intertextis longissimis crassis. *Schmidt Mycolog. Hefte I. p. 84. sub Phyllerio.*

Hab. in Lusatia superiore ad folia *A. pseudoplatani*. Cespit. 2 - 3 lin. lati, plerumque orbiculares ita immersi ut fol. bullata reddant.

Color ab initio albedo pallens, dein aureo nitens, tandem umbrinus. *Synops. fung. p. 700.*

15. *acerinum*, superficiale, passim immersum caespitulis variis, primo pallidis (roseis albidisque) dein spadiceis. *Syn. fung. p. 700.*

Hab. copiose inferne in fol. *Aceris Platanoidis*, ubi colore plerumque badium, sed etiam griseo-fuscum nec non irregulare effusum; in *Acere campestri* plerumque pallidum fere grumosum; in *A. Pseudoplatano* lacte spadiceum, magis molle et in *A. Opulifolio* primo roseum dein lutescens; hinc ab auctoribus minus recte plures constitutae sunt species.

16. *fagineum*, caespitulis immersis confertis subrotundis obscure badiis. *Synops. fung. p. 700. Nees System der Pilze t. 5. f. 63.*

β. *Fagi purpurei*, caespitulis amoene purpureis.

Non infrequens in fol. *Fagi sylvaticae*, sed magis versus marginem superiorem. Ab initio album, tenerum et tunc *Erineum lacteum*, *Fries Obs. Mycol. 2. p. 371*, videtur.

17. *betulinum* effusum planiusculum primo subgrumosum albidum, purpureum et roseum, dein rubiginosum obscurum.

Cresc. in utraque pagina fol. *Betulae albae*, sed magis in superiore, praesertim si co-

lore est rubro. Huc pertinent ut varietates *E. betulinum roseum* et *purpureum* auctorum.

18. *Pyracanthae*, superficiale oblongum obscure rubrum subgrumosum. — *Decand.*

Flor. Gall. Suppl. p. 13.

Hab. in Gallia meridionali in Mespilo *Pyracantha*. An color constans remanet? Fila subdifformia, capitata.

19. *Oxyacanthae*, sublineare immersum dilute flavum nitidum, foliorum marginibus replicatis.

In Crataego *Oxyacantha*. L. rarius crescit, nonnunquam ad latera nervi principalis occurrit, plerumque vero ad margines foliorum nunc reflexos oritur.

20. *populinum*, cespitulis orbicularibus immersis grumosis opacis spadiceis. *Syn. fung. p. 700.*

Frequens in fol. *Populi Tremulae*.

21. *alneum*, grumosum effusum superficiale fulvo-ferrugineum, subnitidum. *Syn. p. 701.*

In Alno *glutinosa*, sed minus frequens.

22. *nervisequum*, superficiale lineare parallelum grumoso-farinaceum albido-pallens.

Schmidt et Kunze Fung. Germ. exstec. fasc. 7. no. 173.

Hab. in Lusatia, in fol. fagineis adhuc junioribus. Forte nondum plane evoluta spe-

cies est, quae, secundum auctores, primo pallide rosea est, et dein pallens. Fibrillae crassae, elongato-clavaeformes.

*** *TAPHRIA*, filis brevissimis subobsoletis, in crustam aggregatis.

23. *aureum*, cespitulis dilatato-depressis magnis aureis. *Syn. fung. p. 700.*

β. cinereum, cespitulis minoribus cinerescentibus.

Provenit passim in *Populo nigra*, *β.* in *P. fastigiata* lecta et mihi missa, cujusque color alienus aetati veteriori attribuendus videtur cum *α.* etiam colorem suum flavum in griseum fere commutet.

24. *griseum*, superficiale tenuissimum orbiculare plerumque effusum griseo-cinereum.

Erineum platanoideum, latissime effusum superficiale infrafoliaceum tenue cinereum, cupulis irregularibus variis. *Fries Obs. p. 229?* (vix ob fila cupulata).

Hab. in fol. fere exsiccatis *Aceris platanoideis* et *Quercuum*. Sub lente tenuiter villosum.

25. *leprosum*, late crustaceum compactum tenue villosum flavescens. †

Autumno satis frequens ad folia viridia quercina, latas maculas exhibens. Vix hujus generis, sed ubi referendum. ?

26. *Cassiae*, angustum nervisequum flavescens.

Inveni ad latera venarum fol. *Cassiae marylandicae* (cultae). Cespituli lineares subinterrupti.

27. *hyperici* effusum tenue lutescens.

Passim in *Hyperico Nummularia*, cujus folia fere tota obtegit.

† *BYSSI DISCRETAE* filis plerumque rigidis (nigris) non intertextis gaudentes sporulis ubique adpersis.

II. *FUMAGO*. Crustaeformis (nigra) ex materie subcompacta, fibrillis raris intertexta formata, foliis afflata. (Color niger.)

* *Conformatione aequali.*

1. *vagans*, late effusa subtuberculosa.

Aestate et autumnii initio, durante praesertim aëre calido, folia *Tiliae*, *Aceris*, *Ulmii*, *Populi*, *Alni* et *Salicis*, materie nigra obducta inveniuntur, si fumo exposita fuissent.

2. *Mali*, orbicularis tenuis subpulverulenta.

Rarius in fol. *Pyri Mali* occurrit. An huius generis?

** *POLYCHAETON*, crustaceum magis villosum, setulis rigidis sparsis distinctum.

3. *Quercinum*, late effusum submembranaceum aterrimum, setis densis rigidis.

Valde rare occurrit in fol. quercinis aridis, quorum totam fere occupat superficiem

pronam. Setulae in pelliculam crustaeformem et fol. leviter adhaerentem coalitae.

4. *Citri*, late effusa crassiuscula nigro-grisea.

Provenit in Europa meridionali ad folia *Citri Medicae*, quae saepe tota induit.

5. *Fagi*, sparsa tenuis suborbicularis.

In Vogesia, supra folia *Fagi sylvaticae* crescentem invenit D. Mougéot. Ab initio maculas sistit sparsas. Color nigro-fuliginus. Setulae magis sunt distinctae.

6. *Ilicis*, orbicularis atra, crusta s. subiculo villosa.

In utraque superficie foliorum *Ilicis Aquifolii*. Lin. 1 - 2 lata.

7. *Typhae*, acervulis irregularibus confertis, setulis copiosioribus.

Occurrit passim in fol. *Typhae angustifoliae*, et reperta prope Dax a D. Thore.

III. CONOPLEA. *Fila* aggregata, cespitulum subglobosum sporulas includentem efformantia.

* BLEPHARIA, filis rigidis passim subramosis divergentibus, exsertis. Sporulis paucis aut nullis.

1. *hispidula*, parvula subrotunda nigra, filis longiusculis. *Syn. fung.* p. 235.

Crescit in foliis, passim et spiculis exsiccatis graminum.

2. *Tiliae*, erumpens conferta subglobosa atro-brunnea.

Exosporium *Tiliae*. Link in Berl. Magaz. d. Naturkunde, 1809. p. 9. (qui fila s. fibrillas pro sporulis s. sporidiis considerat).

Hab. in ramis mortuis *Tiliae*, praesertim gelu tactis, quos saepe totos investit. L. Primo obtutu nigra, sed sub lente observata, badia est. Lin. 1-2 lata.

3. *clavuligera*, erumpens nigra, filis aliis aequalibus filiformibus, aliis clavatis pellucidioribus. — Link in Berl. Magaz. 1815. p. 33.

Proven. in ramis exsiccatis, in Germania.

4. *hypodermia*, elliptica gregaria sub epidermide prorumpens denum confluent, filis elongatis distantibus. Link, l. c.

In ramis dejectis. Parva, magnitudine granorum pulveris pyrii. Link.

5. *quercina*, epiphylla fasciculata nigra, filis mollibus.

Satis frequens occurrit in fol. aridis quercinis. Dematium refert, sed pulvere nullo adspersa; fasciculus obconicus, e filis pro parvitate satis longis, compositus.

6. *Eryngii*, fasciculis confertis brevissimis atris, filis rigidis, aliis obovatis septatis.

Prope Parisios non infrequens ad caules aridos *Eryngii campestris* se propagat, vere.

Crustam elongatam primo intuitu sistens. Sub lente indagata, observantur filis intermixta corpuscula turbinata; passim subglobosa substipitata, interstitiis distincta. Hinc genus distinctum ex hac specie confici posset, *Elosia* denominandum.

**** CONOPLEAE VERAЕ**, *cespitulis mollibus, filis intertextis subfloccosis, pulvere sat copioso refertis.*

7. *sphaerica*, gregaria subconfluens nigrescente-olivacea globulosa, recente in statu subgranulosa. *Syn. fung. l. c.*

Provenit ad ramos exsiccatos fagineos.

C. *Tiliae* quadantenus similis, at duplo major et pulvere sat copioso referta.

8. *fusca*, subrotunda confluensque fusca.

Circa Lipsiam a D. *Kunze* lecta, mecumque sub nomine *Helmisporium subramosum* communicata.

9. *cinerea*, subrotunda confluensque cinerea.

Rarius prope Parisios ad ramos aridos invenit. 3-4 lin. lata.

10. *gilva*, sparsa subcompacta ex gilvo subrufescens.

Ad caules herbarum majorum passim inventa a D. *Cauvin*. Acervuli longe inter se distant, pulvere minus conferti.

11. *atra*, acervulis mollissimis subrotundis con-

fertisque atris pulvere copiose refertis. *Syn. fung. p. 235.*

Provenit satis frequens locis praesertim suffocatis, ad ligna aliaque corpora,

Obs. Confiteri debeo, quod specimina, Parisiis collecta, obtulerunt *sphaeriae speciem*, *Sph. Byssisedae* similem, villo obvolutam, an forte *parasiticam*? cum simul pulvere digitos inquinante scatet. Hinc nisi *Sphaeriae* sint, ad genus *Chaetomium* Kunze referri debet.

12. *cylindrica*, sparsa subcylindrica ferrugineo-fusco. *Syn. l. c.*

In culmis arundinaceis in sterquiliniis locisque cryptis olim circa Gottingam collecta, A priore forte non specie distincta.

IV. DEMATIUM. *Fila* subrigida, ut plurimum fasciculata (simplicia aut subramosa) erecta, *sporulis* globosis aut ovalibus, passim seriatis adspersa, demum in crustam subconfluentia.

1. *vulgare*, *α. herbarum*, acervulis compactis subtuberculatis olivaceo-nigricantibus. *Syn. fung. p. 699.*

*Cladosporium herbarum. Link Berl. Magaz. 1813, p. 37. *)*

*) Character genericus *Cladosporii*, secundum autorem, sequens est: Thallus e floccis cespici-

Crustae in plantarum subexsic-
cente corpore, ibique crustam rudem effi-
pitata, acervulis subrotundis subcon-
tincta. In fol. Irideis, Brassicae etc.

denotatur, tenerius effusum suboliva-
ceis, exsiccatis praesertim agaricis.
In foliis typharum, aestate.

7. atrum sparsum, ac effusum, floccis
septatis. *Link, l. c. p. 38.*

Colore atro ramos dejectos investit floccis
obscurioribus non septatis obscurioribus, minus
lucidis alienum refert habitum. *L.*

Abietinum, tenue late effusum, primo fla-
vo-cinnamomeum, demum canescens. *Syn.*
flaig. p. 699.

Cladosporium abietinum, fuscum expal-
lens, demum effusum, floccis septatis. *Link,*
l. c.

Hab. in cortice *Pini Abietis*, adhuc vi-
gentis.

4. *virescens*, cespitosum laxum, subfarinaceum,

tos erectis simplicibus aut subramosis, api-
cibus in sporidia secedentibus, — sporidiis
(seriatis) apici primum innatis, dein dilaben-
tibus. *Link, l. c. Confer quoque Nees Sy-*
stem d. Schw. 2. p. 18.

primo laete virens, dein subolivaceum. *Syn-
fung. p. 698.*

Sporotrichum virescens. Link, l. c. p. 13.

*β. ? Chloridium dispersum, effusum ni-
gro-olivaceum. Nees Syst. 2. p. 18. t. 5.
f. 64.*

Hab. ad ligna exsiccata et passim caules
plant. majorum, hyeme lene, *β.* in cortice
quercuum, semiputrido. — Flocci erectiusculi,
non ita ramosi ut in reliquis *Sporotrichis*, et
sporidia saepe seriata. *Link*, hinc *Dematii* aut
Cladosporii Link, species.

5. *asseraculorum*, cespitulis tenuissimis, floccis
laete viridibus, sporidiis majusculis. *Lk.*,
l. c. t. I. f. 16. Chloridium viride *).

6. *griseum*, effusum, filis subulatis rigidis fus-
co-nigris, sporulis copiosis inspersis cylin-
dricis albo-griseis.

*Chloridium griseum. Ehrenb. Sylv. My-
col. p. 23.*

Nascitur hyeme in trunco alneo. Spori-
dia nonnunquam auctori visa sunt ramos ar-
ticulatos fingere.

*) *Chloridium*, thallus e floccis simplicibus et
subramosis erectis non septatis. Sporidia ubi-
que inspersa. *Lk.* Hinc verae *Dematii* (*mi-
hi*) species sunt, cum quibus, ceterum, *Clad-
osporia* quoad habitum conveniunt.

7. *fuscum*, acervulis variis tenuioribus submol-
libus dilute fuscis.

Non raro occurrit in caulibus herbarum
exsiccatis. Acervuli sparsi, parvuli, nec ces-
pitem seu crustam efformant.

8. *grumosum*, acervulis densis sparsis minutis
grumosis fusco-nigricantibus.

Cum priore, cujus forsitan est plantula
junior, sed acervuli compacti sunt et colore
obscuriores.

9. *epiphyllum*, acervulis distantibus in or-
bem dispositis nigrescentibus.

β. *Chionanthi*, subcrustaceum, fil. brevis-
simis.

Satis frequens in variarum arborum foliis
siccis, praesertim populneis.

10. *graminum*, sparsum fasciculatum minutum
griseo-nigrescens, filis distinctis laeviuscu-
lis.

Autumno sat vulgare in foliis et culmis
aridis graminum.

Species dubiae et alius forsan generis.

11. *articulatum*, nigrescens fasciculatum, filis
divergentibus subramosis articulatis. *Syn.*
fung. p. 694.

Helmisporium carispermum. Link Berl.
Mag. 1815. p. 39.

Hab. in caulibus plantarum et in grami-
nis. *Conopleae* habitu.

Obs.

Obs. Sporidia globosa, non septata, foramine conspicuo dehiscencia, an inde genus proprium *Coëlosporium* dicendum? Link & c.

12. *verticillatum*, nigrum, filis subramosis verticillatis. Syn. fung. L. c. Hoffm. Flor. germ. 2. t. 13. †

Spondylocladium fumesum. Martius Flor. Cryptog. Erlang. p. 355.

Hab. in caulibus plantarum; et in lignis putrescentibus. Aspermum.

Obs. Duæ posteriores species mihi denuo sese non obtulerunt, ut eas curatius examinare possem. Fila seu fibrae in hac, secundum Martium, sunt annulato-moniliformes, erectae, rigidae, simplices vel subramosae: ramulis verticillatis. Sporidia distincta nulla, an articuli solvuntur?

V. HELMINTHOSPORIUM. Fila (setulae) rigida, plerumque divisa, erecta. Sporulae oblongae, septatae filis adhaerenti-pendulae. (*Helmisporium* Link, Nees.)

1. *nappum*, effusum nigrum, fibris furcatis simplicibusque subnodosis, sporis subcylindricis sparsis fibris paullo brevioribus. Nees Commentatio, p. 17. t. 1. f. 13. b. Ejusd. Syntem, 2. p. 67. t. 3. f. 65.

Hab. in ligno emortuo, maculas longitudinaliter effusas sistens.

2. *simplex*, dense caespitosum effusum; setulis erectis simplicibus in medio subconglutinatis, sporis utrinque acutis obsolete septatis. *Nees, l. c. p. 15. t. 1. f. 11.*

Crescit aestate, in ligno salicum. Olivaceo-nigrum.

3. *tenuissimum*, laxe caespitosum, fibris erectis simplicibus olivaceo-nigris, sporis ad basin inconcervatis ovato-clavatis. *Nees, l. c., p. 16. t. 1. f. 12.*

In caulibus plantarum prope Lipsiam detectum a Dr. Kunze.

4. *subulatum*, setulis distantibus subulatis subsimplicibus nigris, sporulis clavatis parvis basi adhaerentibus. *Nees, l. c. t. 1. f. 13.*

Hab. in ramis quercinus putrescentibus.

Vere.

5. *velutinum*, floccis sparsis subramosis nigris, sporidiis basin versus adhaerentibus pyriformibus elongatis. *Link, Berl. Mag. 1805. t. 1. f. 9.*

Hab. ad ramos exsiccatos. Cum *Periconia* non confundi debet.

VI. MELCOSPORIUM. *Setulae* erectae, rigidae, simpliciusculae. *Sporulae* spirales, remote geniculatae, fugaces, inspersae.

1. *vegetum*, fibris nigris distantibus, sporis vi-

Ordo primus.

19

ridi-lutescentibus. *Nees System*, 2. p. 19.

t. 5. f. 66.

Crescit in assulis quercinis.

2. *pulvinatum*, irregulare olivaceum, fibris decumbentibus ramosis implexis, sporis luteo-viridibus.

Helicotrichum pulvinatum. *Nees Comm.*

p. 20. t. 1. f. 15.

Hab. ad truncos caesos quercuum. Vere. Fibræ tenuissimæ, opacæ, annulis obscurioribus distinctæ, ramulis patentibus. N.

VII. *CIRCINOTRICHUM*. Fibræ decumbentes, tenuæ, circinato-implexæ, opacæ. Sporulae subfusiformes inspersæ. *Nees*.

1. *maculiforme*, olivaceo-nigrum tenue, fibris sericeo-floccosis. *Nees, System*, 2. p. 19. t. 5. f. 66.

In foliis quercinis putrescentibus. Vere.

VIII. *GONOTRICHUM*. Fibræ implexæ, ramosæ, nodoso-articulatæ: ramulis subverticillatis. Sporulae globosæ, inspersæ.

1. *caesium*, caespitulis pulvinatis fusco-caesiis. *Nees, Comment.* p. 18. t. 1. f. 14.

Unica vice in ramulo quercino, fol. putrescentibus oblecto, locis humidis inventum. Sporulae minutæ, in ramulorum apicibus solitariae, in fibris copiose inspersæ, sub aqua fugientes. N.

Obs. Duo antecedentia genera, nec non ambo sequentia a me ipso nondum reperta sunt.

IX. CAMPSOTRICHUM. *Fibrillae* breves, liberae, intricatae, divaricato-ramosae, nigrae. *Sporidia* ramis extremis opposita pellucida.

1. *bicolor*, fibris apice ramosioribus olivaceo-nigris: ramulis extremis pellucidis, sporidiis magnis subovatis rufo-fuscis. *Ehrenb. in Annalib. Botanic. Berol. Fasc. 2. p. 55.*

Usneae destructae ramos ambit tomenti instar, sed filor. rigidorum ac ramosorum laxa compage. Granula sunt magna, ramorum apicibus multo crassiora pellucida. *Ehrenb. l. c.*

X. ALTERNARIA. *Fila* erecta, simplicia sparsa *nodosa*: nodis ovalibus, remotis; interstitiis filiformibus.

1. *tenuis*, filis tenuissimis strictis nigro-olivaceis. *Nees Syst. 2. p. 19. t. 5. f. 68.*

Hab. in ramulis aridis, hinc nigro-maculatis; autumnino.

2. *rudis*, fibris rudibus variis rigidis complexis brevibus nigris. *Ehrenberg Sylv. Mycolog. p. 22.*

Truncos pineos colore nigro obducit. Vere.

XI. TORULA. Effusa subincrustans. *Fila* simplicia, brevia, moniliformia seu articulata: articulis approximatis, passim deciduis.

Ordo primus.

21

* *Articulis durantibus, plerumque ovatis.* (Monilia. Link, Nees.)

1. *antennata*, effusa atra, filorum articulis ovatis. *Syn. fung. p. 694. Hoffm. Flor. Germ. 2. t. 13. f. 4.*

Monilia antennata. Link, Nees Syst. 2. p. 19. t. 67. a.

Frequens autumno et hyeme leni in assulis exsiccatis quercinis. Primo obtutu villum subcongestum refert; junior subolivacea.

** *Articulis globosis, saepius secedentibus.* (Torula Link, Nees.)

2. *herbarum*, incrustans tenera atra, articulis globosis subcontiguis.

Monilia herbarum. Syn. fung. p. 693.

Torula monilis. Obs. Mycolog. 1. p. 25.

Passim in caulibus exsiccatis plantarum majorum invenitur.

3. *tenera*, acervulis subrotundis confluentibus, articulis globosis pellucidis. *Link Berl. Magaz. 1815. p. 40.*

T. tenera, maculaeformis confluenta, atrovelutina, articulis subglobosis inaequalibus. *Nees Syst. 2. p. 20. t. 5. f. 69.*

Provenit in ramis aridis.

4. *fuliginosa*, villosa compacta atra, filis articulatis crassis; articulis globosis subconfluentibus.

α . *pinophila*. *Antennaria pinophila*, nigra hirsuta confluens in ramis Pini Piceae parasitica. *Nees Syst.* 2. p. 72. t. 39. f. 298. *)

β . *ericophila*. *Antennaria ericophila*. *Link in Schrad. Journal für die Botanik*, 1809. vol. 3. p. 17. f. 24. **)

Crescit α frequens in ramis foliisque Pini Piceae, β in *Erica arborea* quam destruit et suffocat.

*** *HORMISCIMUM*. Fibrae aggregatae vel solitariae, simplices, strictae, rigidiusculae, suppelliculae articulatae: articulis globosis continuis. *Kunze Mycol. Hefte* 1. p. 12.

5. *expansa*. *Hormiscium expansum*. *Kunze l. c.* p. 13.

Cespites magnos efficit in plantarum caulibus. Articuli non secedunt.

6. *alta*, fibris variis rigidis aggregatis obscuris atris, articulis magnis. *Ehrenberg. Sylvae Mycolog.* p. 22. sub *Hormiscio*.

Crescit in cortice Alni glutinosi. Aprili.

Obs. *H. expansum* fibras habet subpellu-

*) *Antennaria*. Perithecia irregularia astoma, thecis liberis, pluriannulatis referta, hypothallo fibroso monilioide circumvallata. *Nees l. c.*

**) *Antennaria*. Sporangia tomento imposita e filis moniliformibus econtexto, intus thecis oblongis repleta. *Link l. c.*

cidas, duplo tenniores ac breviores, et articulos triplo fere minoris. *Ehrenb.*

†† *Mucedines*, quarum limites inter ceteros fungos byssoideos non equidem facile circumscribi possunt, forma magis sunt determinatæ, stipitatæ, substantia teneræ, plerumque fugaces, in mucidis et supra corpora fermentescentia plerumque provenientes; colore albidæ, glaucæ, aut griseæ, rarius nigrae; earumque fila (flocci) minus sunt intertexta, plus minusve septata, decumbentia, adscendentia, ut plurimum erecta, stipites et ramulos imitantia, in quibusdam in sporulas (sporidia) secedentia, eas vero in aliis, in capitulum, spiculam, racemulum aut verticillatim aggregatas, vel seriatim dispositas, passim dispersas ferentes.

XII. ACROSPORIUM. *Fila simplicia*, distincta, (passim cespitulos efformantia) moniliiformia: articulis (sporulis?) ovatis secedentibus.

* *Filis discretis erectis.*

1. *monilioides*, candidum cespitose effusum, contactu pulverem movens. *Nees, System der Pilze, 2. p. 14. t. 4. f. 49.*

Monilia hyalina, filis aggregatis simplicibus erectis, sursum crassioribus hyalinis, articularis subgloboso-ovatis. *Fries Obs. Myc. I. p. 210. t. 3. f. 4.*

Hab. in culmis et foliis gramineis; aestate. Color etiam griseus.

*** *ALYCIUM*, floccis aggregatis erectis simplicibus pellucidis, articulis ovalibus in sporidia solvendis. Kunze Mykol. Hefte, 1. p. 11. t. 1. f. 6.

fulvum, cespitosum fulvum.

Alysidium fulvum. Kunze, l. c. p. 12.

In Salicibus cavis detectum in Lusatia superiore.

*** *OTIUM*, floccis implexis discedentibus in articulos ovatos, floccis vage inspersos. Nees, l. c. p. 12.

3. *fructigenum*, caespitibus subrotundis effusisque griseis pallidisque.

Monilia fructigena. Syn. fung. p. 693.

Torula fructigena. Pers. Obs. Mycol. 1. t. 1. f. 7.

Oidium fructigenum. Mycol. Hefte, 1. p. 80. t. 2. f. 22.

Hab. aestate in fructibus putrescentibus, natura paullo firmioribus, praesertim prunis, pyris.

Obs. *Epochnium* *) *monilioides*, floccis griseis, sporidiis nigrescentibus. Link, tam quoad characterem genericum, quam iconem

*) *Epochnium*. Thallus e floccis caespitosis septatis, ramosis, Sporidia oblonga, appendiculo filiformi, inspersa, subseptata. Link, Berl. Mag. 1809. p. 18. t. 1. f. 28. — Albo cinerascens, orbiculare in pyris vetustis exortum. Nees, Syst. 2. p. 11.

longe alius fungillus, eodem tamen in loco proveniens, videtur.

4. *laxum*, floccis erectis divergentibus pallide cinereis, articulis rarius confluentibus pellucidis magnis. Ehrenberg *Sylv. Mycolog.* p. 22. †

Cresc. in Pruno armeniaco putrido.

5. *aureum*, floccis aureis expallescens, sporis concoloribus. Nees, *System* 2. p. 12. t. 3. f. 44. Link *l. c. t. 1. f. 23.*

Hab. in lignis putridis.

Obs. *Trichoderma aureum*. *Syn. fung.* p. 232. quod laudati auctores, ut synonymum allegarunt, vera *Trichodermatis* species est, pulvere grosso copioso, nec seriatim disposito scatens, ac tegmine villosa involutum. Et *Oidium rubens*. Link *Berl. Mag.* 1815, p. 37. (in caseis vetustis) etiam *Trichoderma* videtur.

XIII. ACROTAMNIUM. *Fibrae* decumbentes, ramosae, continuae, opacae, laxae implexae, ramulis extremis subpellucidis articulatis. Nees.

1. *violaceum*, violaceum, ramulis ascendentibus clavatis saturatioris coloris. Nees *System*, 2. p. 26. t. 5. f. 71. Ehrenb. *Sylvae Mycolog.* p. 10.

Cresc. in interiore corticis quercei latere.

Hoc genus ab auctore pone *Racadium* collocatum, a me, uti sequens nondum observatum.

XIV. GEOTRICHUM. Flocci decumbentes, caespitosi, septati, ramosi. Sporulae ovales, utrinque truncatae, inspersae.

1. *candidum*, effusum album. — Link, Berl. Mag. 1809. p. 17. t. 2. f. 26. †

Plantula fugax, tenuis instar tomenti terram in sylvaticis sterilibus et ericetis obtegit.

XV. MYCOGONE. Effusa byssoidea forma indeterminata. Fila globulifera; globuli solitarii pellucidi concolores.

1. *incarnata*, compactior, colore carneo-rufescente opaco.

β. *rosea*, caespitibus densis effusis, sporidiis roseis. Link in Berl. Mag. l. c. p. 18.

Agaricos putrescentes denso villo obvolvit eosque penetrat, odorem foetidissimum spargit, Leprae animali (morbo) quoad colorem similis.

2. *alba*, laxior tota alba. †

Frequentior in fungis putridis occurrit. An forte junior speciei antecedentis?

3. *cervina* dense effusa colore ceryino.

3. *cervina*, caespitibus densis, sporidiis cervinis. Ditmar Fung. Germ. p. 107. t. 53.

Nascitur supra *Pezizam* macropum.

Obs. D. Ditmar affirmat, in globulis se reperiisse sporulas parvas *Uredinis* similes,

hinc potius genus hocce ad familiam *Mucorum* translocari debet, cum quibus convenit globulis solitariis et stipitatis.

XVI. ACREMONIUM. *Fila* repentia subimplexa septata ramosa: ramulis singulis globulis terminatis.

1. *alternatum*, album, ramulis alternatis. *Link*, l. c. p. 15. *Ditmar*, l. c. p. 3. t. 2.

Obducit folia delapsa sicca, verno tempore.

2. *verticillatum* album, ramulis verticillatis. *Link*, l. c. t. 1. f. 20.

Truncos abietinos late investit villo sat dense implexo. Flocci tener et longi, ramis plerumque quaternis. L.

3. *fuscum*, expansum fuscum, ramulis crebris alternatis. *Mycolog. Hefte*, I. p. 79. t. 2. f. 23.

Ad ligna pinea abietina, vere. Filis arachnoideis teneritate simile.

Obs. Genus mihi plane ignotum, Himantias quodammodo aemulans. Quos sub hoc nomine accepi fungillos byssum confusam referebant, minime figuris similem quae formam forsitan nimis distinctam exhibent. Tamen illud genus omittere nollem, cum post ejus conditorem ab aliis quoque adoptatum sit.

XVII. HAFLARIA. *Flocci simplices aut subramosi, distantes. Sporae globosae, hinc inde coacervatae. Nees.*

1. *grisea*, floccis sparsis simplicibus bifidis dichotomisque griseis. *Link, Berl. Mag. 1809. p. II. t. I. f. 12. †*

In fol. siccis *Spargani*, *Arundinis* et aliarum plantarum non raro occurrit. Sporidia facillime delabuntur et flocci nudi supersunt. *Link. Conf. Botrytem cineream.*

XVIII. ACLADIUM. *Flocci simplices erecti simplices aut subramosi caespitosi. Sporae (majusculae) ovatae vel oblongae, circa apices densius congestae. Nees.*

* *Sporulis subdispersis.*

1. *conspersum*, caespitulis densis ex olivaceo et luteo albis. *Link, l. c. p. II. t. I. f. 13. Nees, l. c. p. 14.*

Cresc. in truncis emortuis. Albidum, flavo-virens, effusum; sporis deciduis. *Nees.* An hujus generis? *Sporotrichum* quoddam potius refert.

** *Sporulis in apicibus collectis.*

2. *capitatum* (albidum), caespitulis laeviusculis, floccis albis. *Link, p. 12. f. 13.* Albidum, lutescens, sporis capitatis. *Nees, l. c. p. 14.*

In truncis putridis. Ad *Aspergillum* (*Moniliam*) transit. *Link.*

Ordo primus.

29

3. *roseum*, cespitosum roseum. Ehrenberg,
p. 11.

Monilia rosea. Syn. fung. p. 691. ? Batsch
Cl. fung. t. 12. f. 58.

In chartis, fructib. putrescentibus.

Obs. An species duae posteriores sequen-
ti generi potius adnumerandae sunt?

XIX. MONILIA. Flocci discreti, aut ces-
pitosi, erecti (stipitiformes) subincrassati. *Spo-
rulae* in capitulum aggregatae, in nonnullis
seriatae, subrotundae. (*Aspergillus*. Link.)

1. *glauca*, caespitosa sparsaque cinereo - glau-
ca. Syn. fung. p. 691. Bull. Cham. p. 106.
t. 504. f. 10.

Vulgaris in rebus corruptis praesertim plan-
tarum. Plures hujus dantur varietates aut sub-
species, quarum praecipuae sequentes sunt:

α. fungorum, cespitosa olivaceo - grisea,
capitulis majusculis.

Frequens in fungis siccis, domi locis sub-
humidis conservatis. Stipes sublongus.

β. grisea, maculaeformis orbicularis et
confluens grisea vel cinerascens - glauca.

Crescit in lignis, locis humidis et suffo-
catis.

γ. clavata, congesta alba, capitulis clava-
tis. Supra carnem coctam exsiccatam, rarius.

2. *albicans*, crustacea minutissima albicans, stipitibus penicillatis.

Ad coria (calceos), locis humidis.

3. *candida*, sparsa aut subcespitosa parva candida subpersistens. *Syn. fung.* p. 692.

Omnium vulgaris, omnique tempore provenit in plantis exsiccatis, fungis, etc.

4. *pulla*, subcaespitosa nigra. *Syn. fung.* l. c.

Rarius in fol. et in charta. †

5. *flava*, floccis simplicibus sociatis albis, capitulis flavescentibus. *Nees System*, 2. p.

16. t. 4. f. 60. sub Aspergillo.

In plantis exsiccatis.

6. *ferruginea*, subcespitosa ferruginea nitida.

Mucor ferrugineus. *Sowerby Engl. fung.* t. 378. f. 10?

In lignis, putrescentibus caesis, hyeme.

Forma nec non magnitudine *M. glaucae*.

7. *spongiosa*, late effusa crassa, filis dense contextis flavo-ferrugineis, capitulis primo albis, demum concoloribus.

In fungis putrescentibus rarius in sylvis.

Obs. Plantula singularis, habitu quodammodo Spongiae est, etiam diu ut illa humiditatem conservat. E basi floccoso-byssoides assurgunt stipites hyalini, qui capitula gerunt primo albidula, in nonnullis guttulam solummodo referentia, quae dein fulvo-grisea evadunt.

8. *virens*, caespitibus densis, floccis intricatis virentibus erectiusculis, capitulis concoloribus. *Link, l. c. (Aspergillus).*

Reperta in sebo supra res conditas infuso. Capit. magna, sporidiis globosis seriatis. *L.*

* *POLYACTIS*, floccis erectis virgatis, ramis inferne nudis, ad apicem ramulis brevibus obtusis confertis sporosis (*Nees*). Sporidiis in ramis congestis capitulum sistentibus. *Lk.*

9. *vulgaris*, caespitibus effusis, floccis griseis, sporidiis glaucis. *Link, l. c. p. 16. t. 2. f.*

22. *Nees, l. c. p. 15. (Polyactis).*

Partes plantarum putridas in hortis et caldariis tegit. Cespites format laxos, et late saepe extensos. *L.*

Polyactis ab *Aspergillo* non differt nisi apicibus divisis. *Link in Berl. Mag. 1815. p. 36.*

10. *turbinata*, caespitibus laxis turbinatis, floccis caesiis, sporidiis candidis. *Schmidt in Mycol. Hefte, 1. p. 83. (Polyactis).*

Hab. in lignis putrescentibus pineis.

11. *quadrifida*, effusa extans, floccis teneribus ramosis griseis, apice quadrifidis fertilibus. *Link, l. c. p. 36. sub Aspergillo.*

Provenit in pomis et peponibus putridis.

XXX! *ACTINOCLADIUM*. Fibrillae erectae, rigidae, annulatae, apice umbellatae. Sporidia pellucida, vaga, globosa. *E.*

1. *rhodospermum*, effusum, fibris raris rigidis, radiis stipitem longitudine aequantibus obtusis, sporidiis globosis majoribus dilute violaceo roseis. Ehrenberg in: *Link's Jahrb. der Gewächskunde*, 2tes Heft, p. 52. t. 3. f. 3.

Crevit, tempestate humida, in trunco arido circa Berolinum. Fila pellucida apparent, teretia, annulata, apice in partes tres obtusas fissae. E.

XXI. MENISPORA. Fibrillae erectae, subcorymbosae. Sporulae lineari-lunatae s. curvatae.

1. *glauca*. *Camptosporium glaucum*. Link, Ehrenb. *Sylvae Mycolog. Berol.* p. 11. (absque descriptione).

Nascitur in latere interiore corticis dejecti quercini et betulini omni tempore. E. Sporulis forma peculiari a ceteris Mucedinibus sane differt. Color glauco albidus.

XXII. BOTRYTIS. Stipites erecti, patuloramosi. Sporulae discretas, pedicillatae, in corymbulum aut ad verticillum aggregatae, rarius subsolitariae.

* *B. verne*, ramulis corymbosis aut racemosis.

1. *cinerea*, congesta subsparsaque late effusa fuligineo cinerea. Syn. fung. p. 690. Albert. et Sweinitz *Fung. Nisk.* p. 363.

Sat vulgaris ad caules variarum plantarum exsiccatos; passim parasitica supra *Sclerotium durum*. Sporulae albicantes. Stipites nonnunquam valde septati, praesertim in speciminibus, quae locis humentibus proveniunt.

Obs. Tres sequentes tantum hujus esse varietates videntur.

2. *polyspora*, caespitulis densis effusis ex oliveo griseis, ramulis creberrimis. *Link, Berl. Mag. 1809. p. 14. Ditm. Fung. germ. p. 71. t. 35.*

Cresc. in ramis humi projectis. Sporulae copiosissimae. *Lk.*

3. *cana*, caespitibus rubrotundis laxis incanis, floccis sterilibus decumbentibus, fertilibus erectis, superne ramosis, sporidiis ovalibus magnis. *Mycolog. Hefte. 1. p. 83.*

Hab. in foliis flaccescentibus *Scrophulariae nodosae*, locis umbrosis humidis.

4. *cinereovirens*, caespitibus densis effusis, medio cinereis, versus ambitum virentibus, floccis aggregatis laxis, sporidiis globosis magnis. *Schmidt, l. c. p. 82.*

In ramis decorticatis praesertim *Salicis capreae*, vere detecta, uti prior, prope *Lipsiam*.

5. *macrospora*, floccis sterilibus decumbentibus albis roseisque, fertilibus erectis albis

superne ramoso-corymbosis. *Link, l. c. p. 15. Ditm. fung. t. 101.*

Cladobotryum varium. Nees, System 2. p. 15. t. f. 55.

Prov. in foliis deciduis, praesertim *Quercinis*. Flocci in apice tantum ramosi: ramis brevibus. Sporidia maxima subcylindrica. *Lk.*

6. *agaricina*, alba, ramis divaricatis ramosissimis. *Link, l. c. Ditmar p. 103. t. 51.*

Fungos obducit lana alba. Sporidia maxima, ovalia, creberrima. An satis diversa a sequente?

7. *dendroides*, ex albo fuscescens, stipitibus strictis ramosissimis, sporulis subovato-oblongis. *Decand. Synops. pl. Galliae. p. 14. Bull. Hist. d. champ. p. 105. t. 504. f. 9.*

Hab. in fungis aliisque corporibus semiputridis. Plantula pulchra et congenerum altissima.

8. *elegans*, effusa, primo alba, demum flavescent, ramis plurimis erectis. *Link, l. c.*

In stercore vaccino cespites laxos efficit.

9. *allochroa*, primo candida, demum flava, floccis superne ramosis. *Link, Nees Syst., p. 15. t. 4. f. 53.*

In caulib. plantar. siccis. Stipites albi, demum rufescentes. Sporae exalbidae.

10. *flavicans*, effusa tenuis, ramis virgatis flavicans, albo-liniata, sporidiis minutis globosis. *Link, Berl. Mag. 1815. p. 36.*

Cresc. in ramis arborum dejectis.

11. *aurantiaca*, rotunda effusaque aurantiaca, ramis crebris. *Link, l. c. p. 14. f. 18.*

In caulibus plantarum, praecipue umbelliferarum.

12. *rosea*, effusa laxa extans rosea, ramis virgatis, sporulis globosis minutis in apicibus ramulorum collectis. *Link, l. c. p. 36.*

In ramis exsiccatis arborum.

13. *variosperma*, effusa densa candida, ramis virgatis, sporidiis ovalibus magnitudine diversis. *Link, l. c. †*

Cresc. in ligno putrido. Sporotrichum refert, sed ramulos profert sporidia sustentantes, quae alia majora, alia minora sunt. *Lk.*

14. *parasitica*, caespitosa ramosa candida, racemulis laxis. *Pers. Obs. Mycol. 1. p. 96. t. 5. f. 6.*

Mucor Botrytis. *Sowerby fung. t. 359.*

Aestate provenit in acervulis *Uredinis Thlaspeos Bursae Pastoris.*

15. *ramulosa*, candida rotunda effusaque, ramulis brevissimis. *Link, l. c. p. 14.*

Hab. in caulibus herbarum siccis. Flocci longi, sed rami pauci et parvi. Sporidia majuscula. *Lk.*

16. *diffusa*, candida ramosa, ramis longe lateque diffusis: ramulis undique racemosis. *Alb. et Schwein. Conspect. fungor. Lusat. p. 362.*

Nascitur in caulibus Solani tuberosi semiputridis. Caespites biunciales et majores formans. Sporulae in racemulo singulo subquaternae et fere globosae. *l. c.*

17. *densa*, caespitibus densis rotundis albis, ramis virgatis, sporulis majusculis ovatis albis. *Ditmar fung. german. p. 105. t. 52.*

Hab. ad muscos aut ad ligna semiputrida in Germania. Habitus diversus et nonnullis Sporotrichis similis.

18. *epiphylla*, crustaeformis alba, stipite simplice brevissimo. †

Inveni hanc speciem ab aliis parum diversam in foliis adhuc virentibus plantae cujusdam, quae hunc crispa et bullata erant. Sub lente examinata farinae dispersae enim similis est.

** VIRGARIA. *Flocci erecti, virgati, ramis furcatis. Sporae diffluentes globosae, sursum magis confertae (nec ramulos terminantes?) Nees.*

19. *murina*, caespitibus densis effusis murinis, floccis approximatis rigidis, ramis remotis patentibus. *Ditmar in Sturm Fung. German., Fasc. 3. p. 73. t. 36.*

Hab. autumnno, ad ramos dejectos et se-

miputridos. Quoad iconem sporulae glomeratae sunt magis in axillis ramulorum, qui non pellucidi sunt.

20. *olivacea*, caespitibus effusis tenuissimis olivaceis, floccis remotiusculis brevissimis. *Lk.*

Berl. Mag. 1809. p. 14.

Cresc. in ramis dejectis.

21. *nigra*, effusa nigra, floccis (ramulis) approximatis. *Link, l. c. t. 1. f. 19.*

Virgaria nigra. Nees, System. 2. tp. 14. f. 52.

Cespites densos velutinos efformat in truncis dejectis. *Lk.*

22. *didyma*, effusa fuligineo-nigra, floccis approximatis subramosis; ramis incrassatis, sporulis magnis ovatis didymis. *Mycolog.*

Hefte, Fasc. 1. p. 81.

Proven. ad ligna exsiccata. Caespes unc. 1 magnus. Sporulae copiosae subovatae, septo distinctae.

*** *STACHYLIS. Floccis* adscendentibus, basi implexis; ramulis superne verticillatis abbreviatis obtusis. *Sporis* globosis ad verticillos congestis. *Nees (Stachylidium).*

23. *bicolor*, effusa, floccis sterilibus decumbentibus complicatis griseis, fertilibus erectis superne simplicibus albis. *Link, l. c. p. 15. Nees, l. c. p. 15. f. 56.*

Late investit caules *Rhei undulati* exsiccatos.

24. *terrestris*, effusa alba, floccis sterilibus decumbentibus paucis, fertilibus erectis ramosis. *Link, l. c. f. 21.* sub *Stachylidio*.

Ad terram, praesertim in caldariis observatur.

25. *tenera*, erecta rufescente-grisea, ramulis patulis verticillatis monosporis.

Verticillium tenerum. *Nees, System. 2. p. 15. t. 4. f. 55.*

Ad stipites *Althaeae roseae* inventa.

26. *capitata*, nivea minima, ramulis brevibus ad apices collectis, vesiculis (sporulis?) pellucidis. *Ehrenb. Sylv. Mycolog. p. 25.* sub *Verticillio*.

Hab. in excrementis larvorum. Vere.

27. *acinorum*, fasciculata, stipitibus rectis subglaucis, ramulis ut plurimum ternis verticillatis terminato, sporulis albido-griseis.

Frequens in uvulis putrescentibus. Singulus cespes ex pluribus individuis basi unitis compositus. Stipites stricti, nitidi. Ab initio albidus est hicce fungillus.

Obs. Confer. *Polyactis* (Monilia) *turbinata* et *Aspergillus* (Monilia) *quadrifidus.* *Lk.*

XXIII. SPICULARIA. Stipes simplex subramosusque apice bi-quadrifidus. *Sporulae* in spiculam ovatam (pedicellatam) arcte aggregatae, terminales.

1. *simplex*, grisea sparsa, caule simplice, spiculis pluribus radiatis.

Botrytis simplex. *Syn. fung.* p. 690. *Micheli N. Gen. pl.* p. 212. no. 1. t. 91. f. 1.

Ad ligna putrescentia et graminum culmos.

2. *umbellata*, caespitosa ex albo nigricans, floccis decumbentibus exserentibus stipites plures ad apicem umbellatim divisos.

Botrytis umbellata. *Decand. Syn. pl. gallic.* p. 14. *Mucor umbellatus*. *Bull. champ.* p. 105. t. 504. f. 8.

Crescit in fructibus conditis. A priore omnino diversa videtur. Primo albida, dein ex griseo nigrescens. *D.*

3. *ramosa*, cinerea ramosa, spiculis (4) radiatis. *Syn. fung. l. c.* *Micheli, l. c.*, no. 2. t. 91. f. 3.

Provenit in rebus corruptis.

4. *gemina*, ramosa subdichotoma grisea, spiculis ut plurimum geminis.

Botrytis spicata. *Syn. fung.* p. 691. *Micheli* p. 212. no. 4. t. 91. f. 4.

Provenit in umbrosis. *M.*

5. *alba*, cespitosa alba subsimplex, spiculis radiatis.

Botrytis non ramosa alba. *Micheli, l. c.*

no. 2. t. 91. f. 3. (ubi stipes apice ternatim divisus exhibetur).

Hab. in pomis putrescentibus, ubi individua dense approximata crustam efficiunt. Stipes simplex, hyalinus. Spiculae opacae, rarius bene explicatae. Observavi quoque basin byssoideam colore olivaceam, quae a nostro fungillo diversa videtur. Spiculae quaternae ex Michelio.

6.? *racemosa*, cespitosa ramosissima, sporulis subovato-elongatis spicato racemosis ex albo-cinereis. Bull. l. c. p. 105. t. 504. f. 3. sub *Mucore*. +

Botrytis racemosa. Decand. Flor. Gall. 2. p. 70.

Nascitur in oleribus putrescentibus. D.

XXIV. *Dactylium*. Sporulae simplices oblongae annulatae ex apice stipitis divergentes.

1. *candidum*. Nees System, 2. p. 16. t. 4. f. 53.

Supra corticem quercinum. Candidum gregarium, nudo oculo punctiforme. N.

XXV. *Penicillium*. Sporulae serratae digitatim divergentes in stipitibus erectis.

1. *glaucum*, crustaceum glaucum, stipite simplice. — Link Berl. Mag.

Monilia digitata. *Syn. fung.* p. 693. *Micheli N. Gen. pl. t. 91. f. 3.*

Mucor penicillatus. *Bull. champ.* p. 107. t. 504. f. 11.

Vulgaris in variis rebus putrescentibus.

β. fasciculatum, aeruginosum, stipitibus brevibus dense congestis subconnatis villosis albicantibus.

Non infrequens in bulbis putrescentibus *Allii Cepae*. Mergem in forma minuta exhibens.

δ. expansum, floccis albis, sterilibus magis minusve contextis, capitulis sporidiisque glaucis. *Link, l. c., Nees, t. 4. f. 59.*

Cresc. in fructibus, fungis, et in rebus conditis, ubi membranam quamdam sistit. *Lk.*

2. *racemosum*, albicans, stipite racemoso. *Syn. fung. p. 692.* sub *Monilia*. *Michel. p. 213. t. 91. f. 4.* sub *Aspergillo*.

Terrestris, in viridario quondam observatum a *Michelio*.

3. *candidum*, caespitibus rotundis effusisque, floccis candidis, capitulis concoloribus. *Lk., l. c. p. 17.*

Prov. in fung. aliisque plantis putrescentibus.

4. *roseum*, effusum tenue album, floccis decumbentibus ramosis, erectis simplicibus,

sporidiis roseis albisve. *Link, Berl. Mag.*
1815. p. 37.

In caule Solani tuberosi.

Obs. Characterem genericum aliter dederunt Link et Nees, secundum quos flocci (stipitis) dilatati apex finditur in fasciculum ramulorum capitulum sporarum globosarum colligentem (*N.*). Sporidia in apicibus penicillatis collecta (*Lk.*). Dicit etiam cl. Link, cave ne sporidia *seriata* cum hisce penicillis confundas. Penicilli tamen in duabus prioribus speciebus nihil aliud videntur quam sporulae concatenatae. Species Linkianae hinc parum a sequente genere discrepant.

XXVI. COREMIUM. Stipes crassus, subcapitatus undique tectus floccis penicillatis, quibus sporulae inspersae.

1. *glaucum*, stipite brevi lutescente, capitulo concolore, sporidiis glaucis. *Link B. Mag.*
1809. p. 19. t. I. f. 31.

Prov. in fructibus conditis corruptis frequens. *Lk.*

2. *leucopus*, gregarium grumosum, capitulis glaucis, stipitibus albis subglabris.

Inveni hunc fungillum, sero autumnio, in leguminibus putrescentibus Viciae *Fabae*. Stipites plerumque uniti, passim solitarii, floccosi vel glabri.

3. *candidum*, album, basi floccosum. — *Nees System. I. p. 87. et 2. p. 24. t. 7. f. 8. (excl. synonymo).*

Ad semina Rubi *Idaei*.

4. *citrinum*, gregarium citrinum, stipite tomentoso. *Conf. Link l. c.*

Monilia Penicillus. Syn. fung. p. 692. Obs. Mycol. 2. p. 34. t. 4. f. 2.

Rarius in muscerdis occurrit.

Obs. Species hujus generis habitu peculiari a *Penicilliis* differt, etiam magis perdurant tamen iis nimis affines. Plura individua videntur stipitibus unita, nam in *C. leucope* etiam reperi nonnulla stipite tenerrimo, unicum *penicillum* vel filum ferente instructa. Ad *Isarias* transitum faciunt.

XXVII. CERATIUM. Primo subgelatinosum, dein submembranaceum brachiatum, pilis *sporulas* solitarias (elastice?) defficiendis obtectum.

* *Forma regulari.*

1. *hydroides*, gregarium candidum simpliciter divisum.

Ceratium Hydroides. Alb. et Schweinitz Conspect. fung. p. 358. Nees Syst. 2. p. 25. t. 7. f. 82. Flor. Dan. t. 688.

Isaria mucida. Syn. fung. p. 688.

Provenit aestate post pluvias non raro ad

ligna semiputrescentia, in Salicibus cavis etc. Vix 2 lineas longum, niveum. Primo amorphum, mollis est plantula, uti *Trichiae* juveniles.

2. *aureum*, effusum, ramulis filiformibus creberrimis. *Link in Berl. Mag. 1815. p. 39.*

Ad truncos putridos in Silesia detectum.

** *Forma anomala,*

3. *pyxidatum*, sparsum pyxidatum, trunco simplice apice radiatim cornuto. *Conspect. fung. Lusat. superior. p. 359. t. 12. f. 9.*

Passim ad ligna putrida. Cornua 3 — 10 crassa, obtusa, in pyxidem composita. *l. c.*

4. *porioides*, congestum poriaeforme, in orbem confluent, cornubus simplicibus distantibus erectis. *Conspect. l. c. t. 2. f. 6. †*

In lignis abiegnis. Forma constanter porioides, poris majusculis faviformibus angulatis, 5-8-gonis, in singulo angulo primum tuberculatis, dein cornutis. Sporulae rariores. *l. c.*

XXVIII. ISARIA. Elongata, subclavaeformis, simplex ramosave, sicca, floccoso-furfuracea. (Color plerumque albus.)

* *In Insectorum cadaveribus provenientes.*

1. *truncata*, ramosa, ramis truncatis crenulatis. *Syn. fung. p. 687. Ramaria farinosa. Holmsk.*

Crescit ad larvas et chrysalides insectorum. Pro more alba.

β.? *Damaecornu*, cespitosa alba, clavulis subsimplicibus obtusis, apice nonnullos ramulos exserentibus.

Hab. quoque (in Gallia) ad chrysalides emortuas et terra sepultas. An forte distincta species. Unc. 1 longa, subincrassata, lin. 1. fere crassa, nivea. Stipes, materie farinacea detera, lutescit.

2. *crassa*, simplicissima crassiuscula subconica, stipite distincto glabro. *Syn. fung. p. 687.*

Ut plurimum solitarie provenit ad *chrysalides*, terrae infossas. Tota alba.

3. *velutipes*, simplex clavata nivea, stipite distincto floccoso. *Link, Berl. Mag. 1809. p. 20. f. 32. Ditmar fung. p. 119. t. 54.*

Affinis antecedenti, cum qua loco natali convenit.

4. *arachnophila*, simplex subfasciculata, ramulis clavatis albidis farinaceis. *Ditm. fung. p. 111. t. 55.*

Nascitur supra *araneas* mortuas. Substantia floccosa laxa.

5. *eleutheratorum*, stipite (trichophoro. N.) filiformi torto subcompresso ramuloso: ramulis patentibus. *Nees, l. c. p. 23. t. 7. f. 84.*

Provenit supra *Carabos* aut *Cerambyces* emortuos, hyeme aut autumno. N.

** In diversis plantarum partibus crescentes.

6. *epiphylla*, clavulis elongatis simplicibus basi in cespitulum connatis albis. *Syn. fung.* p. 688.

Rarius in foliis exsiccatis. Subincurva.

7. *bulbosa*, stipite basi bulboso saepe connato, simplice aut subramoso: ramulis spiniformibus. *Nees, l. c. f. 83.*

Cresc. in lignis putridis.

8. *umbrina*, fasciculata umbrina subramosa, ramis strictis aridis. *Syn. fung. p. 689. Icones pictae fung. fasc. 4. p. 51. t. 21. f. 4.*

Ad ramos dejectos. Non farinacea.

9. *glauca*, fasciculata, ramulis strictis glaucis. *Ditm. Fung. Germ. p. 39. t. 19.*

Nascitur supra ramos fagineos dicorticatos. Antecedenti quoad formam similis. Sporulae minutae, globosae, parcissimae. *D.*

10. *clavata*, gregaria subfasciculata clavata umbrino-straminea. *Ditm. fung. p. 113. t. 56.*

In ligno Fagi reperta, aestatis fine. Contextu floccoso, magnitudine lineari.

11. *citrina*, ramosissima pilosa, trunculo citrino, ramulis tenuissimis albis, sporulis globosis concoloribus. *Ditmar, l. c. Fasc. 3. p. 75. t. 37. Syn. fung. p. 689. Ic. et descr. fung. p. 9. t. 5. f. 6.*

Rarius in vetustis truncis. Autumno. Plumulae similis.

12. *agaricina*, conferta alba subramosa rigida: ramulis rectangulis. *Syn. fung. p. 688.*
Batsch Cl. fung. f. 163.

Non raro autumno in Agaricis siccis, semiputridis. Byssos similis.

13. *carnea*, effusa dilabens, primo albida, dein carnea, stipite flaccido albido. *Syn. fung. l. c.*

Ad folia sicca et supra muscos, initio autumnis; Verum cito in pulverem dilabitur. Affinis *Sporotrichis*.

14. *monilioides*, simplex erecta subgregaria, stipite firmulo subpellucido, capitulo oblongo subturbinato aut tereti, filis racemosis farinosis brevibus. *Alb. et Sw. Consp. fungor. p. 362. t. 12. f. 8.*

Hab. in Lusatia, ad ligna corticesque putrescentes Pinorum. Pusilla, vix linearis. Stipes colore variat albo, lutescente aut rufescente. *loc. cit.*

15. *byssosidea*, caespitosa alba byssosidea tenera, ramulos parvos ubique exserens. †

Passim apparet ad *Pezizas*, ex. gr. *Pez. virginiam*, sed minutissima sub lente determinanda species.

16. *sacharina*, effusa candida, filis parvis intertextis, pulvere denso dispergibili oblecta. *Syn. fung. p. 689. Alb. et Sweinitz Consp. fung. p. 362.*

Aleurisma sacharinum. Link *Observ. in Berl. Mag.* 1809. p. 38.

Ad fungos exsiccatos praesertim *Boletos* nascentem olim hanc speciem reperi; cum vero illa deinde sese mihi non obtulerit, incertus haereo cuinam generi adnumerari debeat.

+++ *Byssi fibrilliformes*, quarum fila elongata, simplicia aut sparsim ramosa, repentia, divergentia, cornea, rarius subtremellosa sunt, et sporulis manifestis destituta.

XXIX. *CERATONEMA.* *Fibrillae* liberae, teretes, farctae aut passim cavae, simpliciusculae subcorneae. (Color pallidus, fuscus aut nigricans).

1. *Crabronis*, fasciculatum simplex elongatum medio nodosum, dilute ambrinum, apice pubescens cinereum.

Isaria sphaecophila Ditm. apud Sturm *Fung. Germ.* p. 115. t. 57.

Aestate supra *Vespanti Crabronem* emortuum repertum.

Fila subuliformia, plurima, 4 unc. longa, flexuosa, ultra medium versus basin nodulo praedita, hanc speciem singularem facile distinguunt, quae cum multis *Isariis* quoad locum natalem convenit, sed melius huc trahenda videtur.

2. *capillare*, croceum tenuissimum subpubescens, ramosissimum capillare subcompressum.

Roth in Usteri Annal. Bot. vol. I. p. 8. t. I. f. 3. sub Rhizomorpha.

Hab. ad radices truncorum putridorum sub foliis delapsis.

Lanam croceam primo intuitu refert hic fungus, senescens colorem induit fuscum, repens est et corneus, oculo armato pubescens; ramus primarius durus, flexuosus, griseus, filiformis. *Roth.*

3. *pallidum*, filis elongatis subconfertis pallidis: ramis primordialibus subcrassioribus.

Hab. in Gallia meridionali, inter folia sicca. *Dufour.*

Rami fibrillosi subintricati, amoene pallidi.

4. *tenerum*, filis tenerrimis flaccidis aequalibus, primo aquoso-albidis, dein pallentibus.

Crescit quoque intra acervum foliorum putridorum, sero autumno. Fila paululum ramosa.

5. *bicolor*, fibrillis nigris ramosissimis, ramulis ultimis pallidis.

Hab. passim inter muscos et ramulos dejectos, cum quibus cohaeret.

6. *Hippotrichodes*, fibrillis rigidis crassiusculis fusco-nigris, brevi ramosis.

Rhizomorpha setiformis. *Syn. fung. p. 705.*

Obs. Foliis et ramulis exsiccatis pineis plerumque adhaeret. Non videntur stipites *steriles* Agarici cujusdam; interea similes fibrillae paullo tamen validiores et multo longiores plerumque simplices, quae non raro in foliis quercinis, etc. observantur, verosimile ad Agaricum *Rotula*, cujus pileus nondum evolutus aut delapsus fuit, pertinent.

Species subterraneae.

7. *variegatum*, spadiceum, ramulis junioribus vel lateralibus fasciculatis pallidis.

Rhizomorpha variegata. *Sowerby Engl. fung. t. 431. f. 2.*

Prov. in Anglia sub terra ad profunditatem trium pedum.

8. *Martini*, cespitosum ramosissimum, criniforme brunneum. *Sowerby, l. c. f. 3.* sub *Rhizomorpha*.

Repertum a cl. *Martin* in fodinis metallicis, prope *Derbyshire*, in latos cespites crescens.

XXX. CAPILLARIA. *Fila* laevia, tenuissima (capilliformia), solida, matrici (ligno aut foliis) arcte adhaerentia. (Color fusco-nigricans).

1. *nigra*, filis ramoso-divaricatis distinctis nigris.

Hab. in fol. vaccinii. Fila brevissima.

2. *Arundinis*, effusa maculiformis, filis ramosissimis densis fusco-nigricantibus.

Rhizomorpha confervoides. *Chaillet.*

Hab. in interiore culmorum *Arundinis Phragmitidis*. Oculo inermi ut maculae nigrae, 4-5 lin. latae, apparet.

3. *Phytolaccae*, filis ramosissimis elongato-divergentibus flexuosis badiis.

In interiore et ad dissepimenta caulis *Phytolaccae vulgaris*.

4. *badia*, filis ramosis vagis dilute badiis.

Passim in fol. siccis Fagi sylvaticae. Fila parva, sat longa, minus ramosa.

5. *epidermidis*, effusa, filis intricato ramosissimis sordide fuscis.

Rarius supra epidermidem ramorum invenitur.

- 6? *grammica*, filis (?) nigris brevissimis horsum forsum flexis simplicibus ramosisque.

Copiose in foliis aridis praesertim quercinis. Lineolis similis. An vegetabile quid? An tunc ob naturam et formam a ceteris separanda? *Venularia* denominanda.

XXXI. ACTINONEMA. *Fibrillae* ex centro radiantes, noduloso-exasperatae, matrici arte adnatae.

1. *Crataegi*, suborbiculare maculiforme fusco-nigrum, filis ubique distinctis.

Nascitur supra folia adhuc viridia *Crataegi* torminalis, prope Neuchatel.

Ab initio maculam solummodo efficit fuscam (saltem eam ad nostrum fungillum pertinere puto), dein fibrillae apparent distinctae. E longinquo *Xylomati* similis, sed fibrillae non in crustam continuam unitae sunt, etiam non adsunt thecae, nec sporulae.

2. *caulincola*, fuscescens, fibrillis densis medio in maculam confluentibus.

Rarissime in caulibus Herbarum (umbelliferarum). 2-3 lin. latum, quasi pingue.

Obs. Non videtur subiculum esse *Sphaeria* cujusdam, uti hoc locum habet in *Sph. Himantia*, saltem nullas sphaerulas vidi.

XXXII. FIBRILLARIA. Rhizoidea, fibrillosa, vage ramosa, libera, farcta, subfarinacea (alba).

1. *implexa*, ramosissima implexa subdichotoma, ramulis attenuatis tenuioribus.

β ? *Rhizomorpha fibrillosa*. *Achar.*

Rarius observatur intra truncos cariosos et in cryptis inter quisquilias.

2. *Arbuscula*, ramosissima alba, ramis crassis subcomplicatis: primario dilatato compresso. †

Turino missa a D. *Bulbis*. Ultra 2 unc. longa et lata. An forma indicata ei constans? 3? *vestita*, subpilosa albida, cortice exteriore laxo membranaceo.

Rhizomorpha vestita Rebentisch.

Hab. in Germania. Singularis ob corticem laxum, circumdantem fibrillam glabram, pallidam et solidam. Externe pilis brevibus et sparsis obsita est.

4. *subterranea*, subramosa crassa teres subfarinacea, ramulis crassitudine subaequalibus divaricatis.

Lipsiae reperta sub terra, trunco vetusto adhaerens, fere octopedalis. *Schwaegrichen.*

5. *felina*, subintertexta ramosa farinoso-villosa alba.

Clavaria felina. Decand. Flor. Gall. Suppl. p. 30.

Passim in cellis ad excrementa Felis, quae demum fere tota fibrillis complicatis et dense aggregatis farcta sunt. An potius *Isariae* species?

6. *Xylotricha*, ramosissima implexa tenuis subglabra albida pallens.

Obs. *Rhizomorphae corticata* et *xylotroma Achar. Syn. Lich.* quoque pertinere videntur ad hocce genus, quod, fateor, mere artificiale est, et *Himantiis* nonnullisque *Isa-*

vis affine, sed ab his forma et crescendi modo, *cet.* differt.

Hab. ad palos putredine emollitos, quos penetrat, plantarum radículas capilliformes simulans. Ramulos multo copiosiores et flexiles habet quam ceterae species. Communicata a cl. *Labillardiere*.

XXXIII. RHIZOMORPHA. Rigida, glabra, fibrosa, ramosa, repens, medulla villosa farcta.

Obs. Species plerumque magnae, radicioformes, colore nigrescente aut fusco-badio distinctae. Subterraneae sunt, aut in locis cryptis et humidis, etiam inter lignum et corticem arborum vetustarum crescunt.

1. *subcorticalis*, (α . *cancellata*) magna compressa, fibris primariis parallelis subcanaliculatis, fibrillis transversalibus inter se cohaerentibus.

Rhizomorpha fragilis Roth, Catal. I. p. 232. Rh. hybrida Sowerby fung. t. 392. — Micheli gen. plant. t. 66. f. 3.

Hab. sub cortice variarum arborum, praesertim quercuum. Nitida, et hinc inde materie albicante obducta est. Fibræ nonnunquam in corpus membranaceum satis latum concrescunt.

β . *Tendo*, longissima sublibera compressa fibris primariis aequalibus parallelis, latere fibrillas transversales hinc inde exserentibus.

Provenit eodem in loco. An distincta?
Funiculiformis, minus nitida et colore magis
ad fuscum vergens.

2. *scandens*, crassiuscula fragilis teres sparsim
ramosa laevis, ramulis breviusculis sublibe-
ris aggregatis.

Rhizomorpha scandens. *Rebentisch Flor.*
Neomarch. p. 391. exclus. synonym.

Rhiz. subcorticalis, var. *teres*. *Decand.*
Flor. Gallica, vol. 2. p. 281. Roth. in Usteri
Annal. Bot. vol. 1. p. 8. Obs. III.

Hab. ad cortices arborum. Magna sub-
complicata, horsum versum flexa, passim sub-
compressa, opaca, forma non regulari dis-
tincta.

Obs. Hac in specie praesertim observan-
tur tubercula sphaeruliformia, ab aliquibus, an
jure?, pro fructibus habita.

3. *obstruens*, fibrillis longissimis liberis tereti-
bus simplicibus fuscis, aut badiis.

Rhiz. imperialis. *Sowerby fung. t. 429.*

In Anglia in tubis ligneis aquam ducen-
tibus inventa eosque tandem obstruens; ad lon-
gitudinem forte 100 pedum (Sow.). Cortex fa-
cile rumpitur. Floccus medullaris candidus.

4. *putcalis*, subrespitosa opaca nigra subteres
flaccida: recens apice albicans, axillis sub-
compressis. *Syn. fung. p. 705.*

Hab. passim in puteis locisque aliis suffocatis. Fibrae subtenerae, flaccidae, ramulis fibrilloso-capillaribus.

5. *molinaris*, flexilis laevis subnitida teretiuscula ramosa: ramulis praesertim superioribus oppositis verticillatisque.

Hab. in Vogesia, ad palos semper humidos molas chartaceas sustentantes, *Mougeot*.

Obs. Habitu quodammodo *Rh. scandentis*, sed fibrillis subintertextis tenuioribus et magis flexilibus differt. Ramuli nunc oppositi, nunc tenues, subverticillati, passim fasciculati.

6. *subterranea*, teres nigra glabra ramosissima, ramis ramulisque confertis attenuatis liberis, *Ach. l. c. p. 588*.

Hab. in Fodinis metallicis, ad ligna subputrida. In Anglia nondum detecta,

Rami glaberrimi, atri, subdichotomi, ramis quandoquidem radicanibus, intus flavescentibus, *Humb. Friberg, p. 34*.

7. *dichotoma*, teres fusco-glaucescens, ramis dichotomis patentibus elongatis cylindricis nudis liberis, *Achar. l. c. p. 589. Sowerb. fung. 298*.

Cresc. in Fodinis plumbeis et cupreis in Anglia. Sex pedes longa, hinc inde tuberculifera. An revera diversa a sequente specie? qui

secundum cl. de Humboldt etiam dichotoma est; vid. *Usteri Bot. Annal. vol. I. fasc. 3. p. 55.*

8. *verticillata*, cinereo-fusca pendula, ramis omnibus verticillatis. *Humb. Flor. Friberg. Spec. p. 35. sub Lichene.*

Cresc. imprimis in Fodinis metallicis in Saxonia. 4 ad 6 pedes longa, ramuli juniores virides ex Humb., color ceterum in plan-
ta aqua irrigata niger est.

9. *palmata*, erecta ramosissima cinereo-fusca, ramis compressis palmatis, apice dilatatis. *Humb. l. c. p. 36. Scop. Dissert. I. p. 94. t. 7. f. 1.*

Crescit cum priore. Triuncialis, cinereo-fusca, apice nigricans, rami rigidi. *H.*

10. *pinnata*, fusco-nigricans subnitida ramosa: ramis (brevibus) omnibus oppositis horizontalibus. *Humb. Flor. Friberg. p. 33. t. 2. f. 3. Lichen pinnatus.*

Hab. ad columnas ligneas in Fodinis metallicis Saxoniae. Rami setacei, compressi, juniores laete virides viscosi, intus tomentosi. *Humb.*

11. *criniformis*, ramosissima intertexta cespitosa, fibrillis tenuibus teretibus.

Inventa in Hercyniae Fodina *Dorothea* ad 200 orgyrum profunditatem. Crinum e-

quinum quoad colorem et dimensionem refert. Fibrillae hinc inde subasperae.

12. *Harrimanni*, subterranea, fibrillis simplicibus aut sparsim ramosis flexuosis fragilibus spadiceis: cortice exteriori articulatim secedente. — *Sowerby Fung. t. 431.*

Hab. in Fodinis Anglicis. Rami longi, undulati.

13. *spinosa*, tereti-subcompressa spinosa obscure fusca, ramis sparsis passim anastomosantibus. *Achar. Lichenogr. p. 588. Sowerb. l. c. 299.*

Hab. in Anglia, in Fodinis?

Diffusa, ramis ad angulum fere rectum prodeuntibus. *A.*

β. acutangula, compressa nitida, fibrillis angustis ramosissimis subflaccidis: ramulis breviusculis rectangulis.

Crescit sub cortice in Germania, et a b. *Ehrhartho* olim accepi sub nomine: *Lich. dendrophilus*. Diversa quoque videtur a *Rh. subcorticali*.

14. *phosphorea*, fusca flexilis, fasciculato-ramosa: ramulis patentibus incrassatis. †

Clavaria phosphorea. Sowerby fung. t. 100.

Hab. in Anglia in cellis vinariis. Tam quoad formam quam ob locum natalem a *Rh. subcorticali* omnino distincta est.

15. *fusca*, flaccida mollis subintertexta rufo-badia, ramis plerumque dilatato-palmatis. — *Mougeot et Nestler Stirp. cryptog. vol. 6. no. 558.*

Hab. in Vogesia, ad parietes cuparum cornis vacuarum, sed aqua adhuc semiplenarum, ibique late ac profunde expansa. Rami inferi in massam informem confluunt. *Moug.* Species sat magna et nitida.

16. *terrestris*, cespitosa parvula opaca rugulosa, fibrillis intertextis subangustis brevibus.

Hab. ad terram, locis suffocatis. *Thuillier.* Cespitulum, duas fere uncias latum efficit, terrae arcte adhaerens eaque saepe inquinata, opaca nigra, in statu sicco rugulosa.

17. *filicina*, parvula compressa nitida dilute spadicea, ramis rectangulis ramulis brevissimis digitatis. — *Rebentisch Ind. Florae Berolinensis.*

Hab. prope Berolinum. Species pulchella, 2 unc. longa.

18. *simplicissima*, erecta atro-nitida simplex aut subramulosa, apice albicans.

Nascitur ad caules aridos Chaerophylli sylvestris prope Neocomium. *Chaillet.*

Gregarie crescit instar Clavariae. Individua juniora crassiora sunt apiceque magis albida; adulta 2 — 3 uncias longa, attenuantur

et hinc inde ramulos ut plurimum breves exserunt. Intus medulla alba farcta.

19. *fontigena*, fibrosa plana sanguinea, fibris basi connatis apice divergentibus. *Rebenth.*

Flor. Neomarch. Prodr. p. 391. t. 4. f. 19.

Hab. ad fontium siphones putridos. Valde depressa ligno adhaeret, et flocco medullari destituta videtur. Ulterius examinandus fungus.

20. *Stokesii*, parvula reticulata spadicea, fibrillis primariis crassis angulato-ramosis; fibrilleolis copiosis reticulatis apice furcatis. —

Sowerby, fung. t. 430.

Hab. in Anglia ad lapides. Species singularis, 2 unc. fere lata et alta.

21. *flabelliformis*, parvula, cauliculis basi teretibus sursum dilatatis et divis in ramulos aequales subpalmatos albidos.

Hab. in quisquiliis.

Species pulchella, quadantenus referens, magnitudine excepta, *Sphaeriam ramosam*. *Sowerb. fung. t. 395. f. 3.*, sed intus cava est et sericea, nec apices pulverulenti sunt et minus divisi; ramuli quoque fastigiati sunt.

Obs. Nonnullae ut *Rhizomorphae* adhuc descriptae sunt a cl. *Achario* in *Actis Suecicis*, an. 1814. (vid. quoque *Fries Observ. Mycol. I. p. 212 — 215.*), quas vero videndi et

examinandi facultas mihi defuit, et harum aliquae forsitan ad antecedens genus pertinent. Haec vegetabilia quoad naturam et crescendi modum magis ad familiam Fungorum (sensu latiore), quam ad Lichenes pertinent.

XXXIV. PHLEBOMORPHA. *Fibrillae repentes, tremelloso-gelatinosae, reticulato-venosae.*

1. *rufa*, crassiuscula subpersistens anastomosans rufa. Tab. VI. Fig. 1. et 2.

Provenit hyeme leni ad trabes putridos, passim quoque ad *Boletos suberosos* exsiccatos. Non videtur status junior *Physari* cujusdam, quod etiam magis fluxile est massam amorpham sistit, nec ita sero invenitur. Fibrillae ad marginem praesertim nodulose confluunt.

2. *lurida*, epiphylla elongata grisea fluxilis vage venosa.

Autumno pluvioso ad folia putrida passim observatur, super quae ad 2-3 unciam serpit, vermis adinstar. Ab initio fere aquosa, hinc propemodum evanescit et vestigium, ut maculas lineaeformes, colore obscuriore relinquens.

+ *MESENTERICA*, ramosissima, ramificationes membranula gelatinosa secum junctae ut mesenteriforme expansum constituent. *Tode*,

3. *lutea*, citrina vel aurea.

Mesenterica tremelloides var. *æ*. *Tode*
Fung. p. 7. t. 2. f. 12.

Hab. ad ramos exsiccatos aut assulas lignorum.

4. *caerulea*, glauco-caerulea. *Tode*, l. c. p. 8.
var. β .

Autumno in assulis semiputridis. *T.*

Obs. Plantulae hae (mihi ignotae) delicaetae sunt et intra 12 horarum spatium, aura calidiore ita corrumpuntur, ut praeter bracteolas quasdam minutissimas conglomeratas, nullum earum vestigium relinquatur. *Tode*.

5. *grisea*, tota grisea. *Albert. et Schwein. Consp. fung.* p. 374.

Arbusculum perfectissime effiguratum offerunt venae gelatinosae repentes, margine tumidulo, colore a sordide rufo cinerascente. Inventa fuit in hymenio *Boleti fusci* semiputridi. Septemb. l. c.

6. *sanguinolenta*, latissima expansa lactea, ad contactum intense sanguinolenta. *Consp. fung.* l. c.

Latubat inter corticem lignumque trunci ejusdam crassi putridi ad spithamas complures expansa. Intacta colorem habebat ex albo dilutissime glaucescentem, sed ad merum aëris liberi contactum aut digiti pressione, illico colore cruento inquinatur. l. c.

++++ BYSSOIDEI VERI, ex filis tenuissimis, intertextis flocci-lanae-pauniformes, passim coriacea, plerumque difformes, maxima ex parte thecis et sporulis destituti; colore varii, et extensionem saepe satis magnam acquirentes.

XXXV. HYPHA. Flocciformis, mollissima ad tactum diffuens, fugax.

Obs. Fungi sunt subterranei, aut in locis suffocatis et cryptis vegetant, colore plerumque albi.

1. *bombycina*, subrotunda indeterminata, humida floccos gossypinos referens.

Hyphasma floccosum. *Rebentisch, Flor. p. 396. Dill. Hist. Musc. p. 5. t. 1. f. 9.*

Dematium bombycinum. *Syn. fung. 696.*

Hab. in fodinis et cellis, lignis et lapidibus adhaerens. Colore nivea.

2. *flabellata*, dilatata fasciculato-membranacea subpeltata basi in petiolum condensata.†

Provenit iisdem in locis, sed rarissime. Magis compacta est.

3. *elongata*, plus minusve in ramos teretes varie elongatos et subdichotomos divisa.

Byssus elongata. *Decand. Flor. Gall. 2. p. 67.?*

In subterraneis, 3 — 7 lin. crassa, et fere spithamam longa.

4. *muralis*, latissima flocculosa candida, exsiccata submembranacea. †

Hab. in cavis et cellis humidis, ad muros aut ligna, nonnunquam latitudine bipedali sese extendens.

Obs. An forte idem vegetabile ac sequens? sed nulla forma indutam eam hactenus vidi. An subiculum *Poriae* cujusdam? conf. *Sowbery Engl. fung. t. 387. f. 6.*

5. *argentea*, fibroso - membranacea, filis e centro radiantibus in ramos tenuissimos divisis.

Byssus parietina, β . *argentea*. *Decand. l. c. p. 66. Vaill. Bot. Paris. t. 8. f. 1.*

Mesenterica argentea. *Syn. fung. p. 706.*

Hab. in domibus, ad parietes et laqueatam contignationem, locis humidis et suffocatis. Elegans, latitudine palmari.

β .? *cellaris*. filamentosa alba, filis repentibus tenerrimis implexis. *Humb. Flor. Friberg. Spec. p. 66. (exclusis synonymis?)*

Hab. in omnibus puteis atque cuniculis Fodinarum, saxa atque trabes obducens. Elegantissimos interdum orbiculos fingit latitudinis semipedales. *Humb.*

6. *speciosa*, repens nivea ramosissima, ramis flexuosis compressis apice dilatatis orbiculatim dispersis. *Humb. Flor. Friberg. Spec. p. 69.*

p. 69. t. 4. f. 18. (sub Byssos, uti sequentes).

Hab. in omnibus cuniculis Fribergensibus lignum atque saxa obducens, circulos formans diametri 2 - 5 - pedalis, mollissimos. *H.*

7. *plumosa*, nivea ramosa, ramulis oppositis pinnata. *Humb. l. c. t. 2. f. 1.*

Crescit cum priore. Bipollicaris, dependens, nivea, pappo plumoso simillima, sericeo-splendens. *H.*

Obs. Ill. de Humboldt in Libro citato nonnullas adhuc descripsit Byssos subterraneas, forma anomala plerumque gaudentes, quas ulterius observandas enumerare quoque velim.

8. *digitata*, dependens, filis longissimis in funem niveum contextis, apiceque 5 - 7 - dactyla. *Humb. l. c. p. 67. t. 3. f. 10.*

Hab. in Fodinis metallicis Saxoniae, e columnis ligneis pendula. Funis mollissimus 3 - 4 - uncialis, ex filamentis parallelis contextus, sed versus apicem in 5 - 7 ramos digitatim divisus. *H.*

9. *clavata*, erecta nivea versus basin attenuata clavaeformis. *Humb. p. 67. t. 4. f. 17.*

Attactu destruitur, uncialis. Fila apice intertexta, sericeo-splendentia. *H.*

10. *Penicillum*, erecta nivea, filamentis fasci-

culatim congestis. *Humb. p. 68. Scopol. Diss. I. p. 91. t. 2. f. 3 et 4.*

11. *minima*, filamentosa nivea, filamentis erectis simplicissimis minimis. *Humb. p. 66. Scop. l. c. p. 90. t. 2. f. 1.*

Cresc. ad columnas ligneas, B. *Penicillo* affinis, sed filam. rarius fasciculata. *H.*

12. *radiciformis*, nivea tenerrima, filamentis ramosis pendulis, globis floccosis hinc inde adhaerentibus. *Humb. p. 68.*

Ad columnas ligneas. Corpora globosa celluloso - membranacea, fere fungosa, semper adsunt et cum ramulis arcte contexta, hinc non parasitica. *H.*

13. *globosa*, expansa radians dein in globum pendulum glomerata. — *Humb. p. 70. Scop. Diss. p. 93. t. 6.*

In omnibus Fodinarum cuniculis passim occurrit. Expansio filamentosa, parte inferiore sustinet saepe ingentes glomeres concolores niveos. *Scop.*

14. *membranacea*, explanata nivea, membrana ex filamentis constans dense intertexta. *Humb. p. 69. Scop. l. c. p. 92. t. 4.*

In membranam sinuatam panni ad instar expansa. *Scop.*

15. *laciniata*, e simplici petiolo pendens substantia (e filis rectis contexta) et in lacinias

multas planas imbricatas dissecta. *Scop. l. c. p. 92. t. 5. Cum priore.*

XXXVI. RACODIUM. Expansum molle, filis intertextis persistentibus seu rigidioribus, sporulis manifestis non adpersum.

* *Colore nigro.*

1. *cellare*, late effusum, ab initio laxo contextum flavescens, dein ex olivaceo nigrum. *Syn. fung. p. 701. Nees v. Esenbeck System d. Schwämme. 2. p. t. 5. f. 70. Dill. Musc. t. 1. f. 12.*

Hab. frequens in cellis, dolia vinaria vetusta longe lateque obducens. Primum lanuginosum, et colore flavum aut rubellum. Maturum magis compactum nigrum, et granulis e filis submoliniformibus conglobatis passim adpersum.

2. *vulgare*, expansum molle laxo contextum nigrum, filis aequalibus. *Fries Observ. Mycol. 1. p. 215. Racodium nigrum. Schumacher. Flor. Saelland. p. 447.*

Hab. in ligno et ramis *Betulae*, *Fagi*, *Pini*, inque muscis et foliis, locis umbrosis. In Suecia et Dania.

3. *Resinae*, expansum molliusculum dense contextum nigrum, filis inaequalibus divaricatis. *Fries l. c. p. 216.*

Crescit in resina *Pini Abietis*.

β. *piceum*, filis per fasciculos elongatos dispositis.

Etiam supra resinam proveniente[m] ex Helvetia habeo; sed forte fila ob nimium imbrem talem acquisiverunt dispositionem.

4. *rupestre*, complanatum nigrum filis rudioribus contextum. *Syn. fung. l. c. Dill. Musc. p. 9. t. 1. f. 18.*

Crescit ad rupes, in regionibus montosis; crusta *Leprariae* cujusdam albae ei saepe immixta invenitur.

5. *turfaceum*, subcompactum dilatatum nigricans, filis brevissimis.

Hab. in Helvetia, in turfaceis et sphagnosis subtus magis condensatum ibique colore subspadiceum.

* *cornutum*, late effusum atro-fuligineum corpusculis crassis elongatis compactis intermixtis.

Hab. in Gallia (Rouen), et Helvetia ad caudices *Betulae albae* quos destruit, latitudine saepe magna.

Obs. Corpuscula Slerotioidea? nunc recta, saepius tamen subincurva, parum compressa et striata, substantia sicca et colore paullo sunt dilutiora. Forte sui generis vegetabile.

** *Colore gilvo, ferrugineo.*

7. *fodinum*, subcompactum molle, superne opacum gilvo-spadiceum, inferne pallens.

Hab. ad januas in Salifodinis in Helvetia etiam in aliis cryptis, nec non in cellis (Pari-siis, Beaur.) muris adhaerens. Spithamiam plus minusve longum, unc. 1. latum, ad tactum molle.

β. *dealbatum*, fere albicans cum ligno totum connatum.

8. *Aluta*, late expansum densius contextum cinnamomeum nitidum. *Syn. fung. p. 702.*

Prov. rarius in interiore arborum excava-tarum. Pulchra, sed parum cognita species.

β. *quercinum*, minus molle, superficie inferio-re cum ligno arcte connatum.

Hab. ad caudices caesos quercinos prope Neuchâtel. *Chaillet.* Colore exacte cum an-tecedente convenit, sed paullo durius est.

9. *subcorticale*, late effusum subtenue inter-ruptum ferrugineum, superficie subunita de-mum canescente.

Hyphasma flavescens, pallide flavescens tomentosum, aetate pulverulentum ferrugineum. *Rebent. Fl. Neom. p. 397?*

Hab. in Gallia inter corticem et lignum. Hinc ad *Xylostromata*, etiam ob superficiem accedit, sed substantia interna lanuginosa. Ae-tate profectiore etiam in pulverem parum so-lutum est, tunc *Sporotrichis Linkianis* simile.

10. *strigosum*, crassiusculum densius compac-

tum, obscure ferrugineum, filis nonnullis rigidioribus subtrigosum. †

Hab. in Pedemontio. *Balbis*. Ligno molliori arctius adhaerens. Filis dense intertextis et colore opaco non ad *Ozonium auricomum* pertinere potest.

11. *badium*, longitudinaliter effusum, filis densis parallelis contextum.

Hab. ad ramos exsiccatos, prope Parisios.

Obs. Colore amoene ferrugineo - badio et dispositione fibrillarum, quae ad marginem saepe in superficiem unitam condensantur, sese distinguit.

12. *fuscum*, sublate effusum tenuissimum dense contextum opacum fusco - badium.

Parisiis in cellis ad muros detectum a cl. Beauvois; 2 unc. latum, 4 unc. longum, calce saepe inquinatum, filis non distinctis.

13. *rubiginosum*, mollissimum rufo - spadiceum, fibrillis nonnullis distinctis crassiusculis.

Ad muros in cellis habitare videtur ob particulas calcareas adspersas.

*** *Colore pallescente.*

14. *striatum*, sordide pallens in membranam tenuem subcompactum, striatum liberum.

Racodium *Aluta* Schleicher (excluso synonymo).

Hab. in Helvetia.

15. *ochroleucum*, subparvum effusum laxius contextum ochroleucum nitidum.

Hab. in Helvetia, prope Neufchatel, ad ligna exsiccata cellulosa, iisque inmersum. Colore primo subochraceo, grato distinguitur. Vix $\frac{1}{2}$ unc. magnum, applanatum.

16. *linteaceum*, dilatatum submembranaceum liberum, amoene pallidum.

Hab. in Gallia. Locus natalis proprius mihi ignotus. Textura Lintei fere, et ejus color chirothecis coriaceis similis est, sed paullo dilutior forte exsiccatione.

17. *lanatum*, laxum mollissimum albidopallidum lanam referens.

Hanoviae repertum ad ligna a D. Nees.

**** Colore albo.

18. *papyraceum*, subtenue late effusum, ligno (quod saepius intrat) arcte adhaerens.

Syn. fung. p. 703. Sowerby Engl. fung. t. 387. f. 10?

Crescit supra et intra ligna cariosa, praesertim Salicum. Subtomentosum, sed ad substantiam submembranaceam tendit.

Obs. Non raro ad Ligna exsiccata per intervalla et intra cavitates reperiuntur flocci sericei, amoene albi, sed non intertexti, quos ne cum Racodio nostro confundas, cave. Hi enim nihil aliud videntur, quam *fibrae* ipsius

ligni in *Asbestum* vegetabilem, ut ita dicam, solutae.

18? *Entomogena*, globoso-flocculosum niveum dein confluens.

Sporotrichum densum. Link, *Jahrbücher d. Gewächsh.* p. 272?

Hab. ad Scarabeos diutius emortuos, inter folia sicca, praesertim in *Melolonthae* majalis *elythris*. Nullas sporulas vidi, quod etiam valet de sequentibus.

19? *Mycobanche*, latius effusum tenuissimum subcompactum albidum.

Non raro autumnno post multas pluvias ad *Pezizas* majores, praesertim *P. Macropum* externe et interne et in disco cupulae *P. hemisphaericae*, quas extinguit.

20? *Rhizophila*, subterraneum candidum subflosculosum.

Observatur ad extimas fibrillulas *Arbustularum* languescentium aut emortuarum, quibus valde noxium.

Obs. Tres hae species, quoad locum natalem inter se et ab aliis valde diversae, omnino melius et repetito examine observari merentur.

XXXVII. *SPOROTRICHUM*. Expansum persistens molle, filis densius et cum pulvere seu

sporulis (sporidiis *Link*) intertextum aut iis adspersis *).

* *Colore albedo.*

1. *laxum*, tenuissimum laxum demum effusum, floccis adpressis vagis, sporulis oblongis. — *Link*, *Jahrb.* p. 166. *Nees ab Esenb.* *System.* I. p. 49. t. 3. f. 45.

Cresc. in truncis caesis et putridis, super quos tenuiter, sed nonnullas uncias longum, effusum est.

2. *candidum*, minus tenue laxum demum effusum, floccis adpressiusculis vagis, sporidiis globosis. *Link*, l. c. p. 166. *Nees*, l. c. f. 45. B.

Cum priore, a quo praesertim differt sporulis rarioribus globosis.

3. *obducens*, caespitibus tenuibus densis limi-

*) Hujus generis monographiam dedit cl. *Link* in *Annalibus Botanicis* (*Jahrbücher der Gewächskunde*) fasc. 1. p. 163 — 185. Fatendum tamen est, quod hoc et antecedens genus solummodo caractere negativo separanda sint, cum habitu et crescendi more inter se conveniunt. Etiam nihil impedit, quin *Racodii* species, melius observatae, favente loco et caelo, sporulis plus minusve copiosis adspersae invenire possint. Nonne *Aleurisma* ut genus particulare separandum?

tatis, floccis strictis adpressis, sporidiis globosis. *Link, l. c. p. 168.*

Hab. in sylvis circa Berolinum ad cortices arborum. *Sporulae* paucae et villo arcte adhaerent. An forte initium alius speciei. *L.*

4. *polysporum*, crassiusculum densum limitatum, floccis implicatis, sporulis globosis minutis copiosis. *Link, l. c. p. 169.*

Hab. in ligno putrido. Lineas quasdam tantummodo latum; aetate juniore globulos sparsos efformat, qui demum confluent. *L.*

5. *sporulosum*, crassiusculum limitatum, fibrillis discretis, sporulis globosis minutis copiosissimis. *Link, p. 169.*

Aleurisma sporulosum. Link, in Berlin. Mag. 1809. p. 19. t. 1. f. 25. (et Ejusd. A. in-
persum.

β. *Aleurisma erubescens. Nees, fung. 2. p. 13.*
t. 3. f. 48.

Caespitulos 2-4 lin. latos format in ramis dejectis. Fila tantum adsunt parva, sed pulvere copioso oblecta. *L.*

6. *fructigenum*, caespitulis crassis confluentibus, floccis densis implicatis, sporulis globosis magnis. *Link, l. c. p. 169. Ejusd.*

Aleurisma macrosporum. Obs. 2. p. 38.

In fructibus *Cerasorum*, acervulos lin. 1-2 latos efficit, qui demum confluent. Fila val-

de tenacia, quibus sporulae firmiter adhaerent.
Lk.

7. *densum*, caespitibus crassis limitatis, floccis densis implicatis, sporidiis globosis minutis.
Link, p. 172. Nees, p. 49. t. 3. f. 45.

Hab. in Melolonthis emortuis, earum pedes et antennis praesertim obducens. L. Etiam ad ramos. Prof. Nees.

8. *fungorum*, tenue effusum, floccis lanatis crispis, sporulis minutis globosis. Link, l. c. p. 170.

Hab. in Agaricis putrescentibus, quos lanae ad instar nonnunquam totos involvit. Flocci non semper crispi sunt.

Obs. Cavendum, ne *Moniliae* aut *Botrytidis* aliusque fungilli species nondum evoluta, pro hac plantula sumatur. — *Sporotrichum agaricinum*, Link (l. c. p. 170.), de quo auctor quoque dubitat, an verus sit fungus, potius materiam farinaceam, nullis filis intertextam, in superficie pilei Agaricorum exsiccatorum secretam, quam vegetabile quid sistere videtur.

9. *latebrarium*, rupestre, albido-griseum crassiusculum stuposum, sporulis copiosis globosis. — Link, p. 171. Achar, Syn. Lich. p. 551. (Pulveraria).

Hab. locis praesertim umbrosis, in rupium fissuris, latitudine nonnunquam insigni.

ni. Fila sub microscopio observata geniculata seu septata sunt, dense intricata et ob sporulas copiosas farinosa apparent. *Link.*

10. *chioneum*, cespitosum laxè cohaerens subgrumosum niveum.

Cresc. ad rupes aut ad terram in Gallia. *Delise.*

Cespitulum pulvinatum latitudine sesquiunciale efficiens.

11. *canescens*, late effusum tenuissimum canescens seu albido-pallens breve villosum.

Prov. ad ligna exsiccata latitudine fere palmari, e longinquo Thelephorae simile, sed sub lente observata villum tenuissimum basi in membranam subconnatam offert. An initium fungi alicujus? Sporulas quidem non vidi, tamen *Racodium* non refert.

**** Colore griseo.**

12. *murinum*, terrestre tenue griseum, floccis divaricatis, sporulis minutis globosis. *Link.*
p. 173.

Hab. ad terram udam, quam lanugine tenera, saepe latitudine ampla obducit. Fila exsiccando evanescunt. *L.*

13. *griseum*, subtenue griseum, floccis implicatis, sporulis crebris globosis. *Link, l. c.*

Provenit ad caules plantarum, praesertim prope radices. Sporulae murinae. Tomentum rufescente-griseum.

- 14? *epiphyllum*, tenue laxius contextum griseum, filis ramosis.

Rarius inveni hanc speciem, forsitan alius aut proprii generis, ad paginam inferiorem foliorum *Rumicis Acetosae*, quam nonnunquam totam obtegit.

15. *fuliginosum*, cespitulis parvis: aliis discretis, aliis confluentibus fuligineis, sporulis minutis subrotundis creberrimis.

Rarius provenit ad cortices plantarum. *Chaillat.*

*** *Colore villi albo, sporularum fusco.*

15. *fusco-album*, tenue laxum effusum, floccis lanatis, sporidiis crebris magnis globosis. *Link, p. 177.*

Ad latitudinem unc. 1. et ultra supra cortices arborum putrescentium nascitur. Lanam compactam, in ambitu albidam, et medio sordide fuscam sinit. Sporidia materie quadam referta auctor quoque observavit.

- 16 *oosporum* (phaeosporum), effusum crassiusculum albicans, floccis intricatis validis mollibus hyalinis, sporulis crebris magnis ovatis fuscis. *Ehrenb. Sylv. Mycol. p. 11.*

Prov. in ramis putridis quercinis Julio. A S. *laxo*, quod auctor etiam enumerat, sporularum colore tantummodo distinctum videtur.

**** Colore ferrugineo, fusco aut badio.

16. *Bryophilum*, floccis laxiusculis primo rubiginosis, sporulis globosis creberrimis.

Hab. rarius ad *Hypna* majora, quorum ramos tractu longitudinali obducit, locis umbrosis et subhumidis sub arboribus. Color demum obscurior et opacus fit.

17. *fuscum*, late effusum crassum fusco-badium, demum subcompactum, sporulis parvis.

β. *Sporotrichum badium*. Link?

Corticium fuscum, *Obs. Mycol.* 1. p. 38.

Thelephora vinosa. *Syn. fung.* p. 578.

Hab. passim ad ramos exsiccatos.

18. *Hypnophilum*, opacum obscure pallidum subcompactum, sporulis grumosis albidis.

Autumno passim ad *Hypna* majora invenitur, cespitulos complanatos, non equidem magnos sistit.

19. *lapidum*, laxius contextum spadiceum, filis nonnullis crassioribus. †

Semel inveni (circa Parisios) ad lapides. A priore differt colore nitidiore et textura minus compacta. Sporulae subrotundae parvae et copiosiores. *Ozoniis* quadantenus affine.

***** Colore virescente, olivaceo.

20. *chlorinum*, virescens densum mollissimum, sporulis globosis conglutinatis flavo-viridibus. — Link, p. 181.

Hab. supra fol. dejecta; textura laxum. Sporulae sparsim glomeratae, facile deciduae, colore flavo-viride distinctae.

21. *aeruginosum*, tenue contextum aeruginosum, sporulis parvis.

Racodium aeruginosum. Syn.fung. p.700.

In truncorum caesorum fissuris solummodo bis inveni in Germania. Aetate colorem mutat in pallide olivaceum, etiam volumen amittit.

22. *olivaceum*, villo laxius contexto aeruginoso, sporulis medio congestis lutescente-olivaceis.

His in regionibus unicum quidem inveni specimen parvum lapidem ad terram obducens sed integrum, et uti pellicula a lapide facile detrahi potest.

23. *psittacinum*, densum crassiusculum, flavo-virens, floccis tenuibus septatis, sporulis raris. *Link, Obs. 2. p. 35.*

Ad ramos dejectos in Lusitania.

***** Colore toto rosco, rubro.

24. *anthochroa*, subcompactum amoene roseo-rubrum, ambitu albicante.

Thelephora anthochroa. Syn.fung. p.576.

Rarissime ad ligna cariosa inveni. Cespitem vix $\frac{1}{2}$ unc. latum efficit, colore ut in sic dicto flore cobaltino distinctum.

25? *roseum*, expansum roseum dein grumosum griseum. Hypasma *Rebent. Flor. p. 397. t. 4. f. 20.*

Ad vetustas molarum januas. Primo ut caespitulus provenit, demum maculas 2 unc. latas repraesentat. *Reb.*

***** *Villo colore albo, sporulis rubris.*

26. *mycophilum*, caespitibus tenuibus subeffusis margine albis, floccis sublanatis, sporulis globosis rubris copiosissimis. *Link, l. c. p. 179.*

Prov. ad fungos majores exsiccatos tractu magis elongato, passim quoque acervulos efformat dispersos. *L.*

β. *rubicundum*, totum albo-incarnatum. —

In Agaricis destructis. *Nees ab Esenb.*

27. *scotophilum*, caespitib. crassiusculis limitatis, floccis implicatis laxiusculis, sporulis globosis rubris. *Link, p. 180.*

Hab. in merdis exsiccatis.

28. *stercorarium*, crassiusculum limitatum, floccis raris, sporulis copiosissimis globosis. *Lk. l. c.*

Obs. Forte una cum *Sporotricho merdario Lk.* melius ad genus *Trichoderma* referri debet.

29. *muscorum*, ex pallido rubescens, sat late effu-

effusum, villis intricatis, sporulis globosis copiosis.

Passim supra cespites muscosos effusum invenitur, nonnunquam ultra unc. 1. latum. Affine quoque videtur Trichodermatibus, imo et Isariae roseae.

30. *vesicarium*, cespitibus tenuibus ambitu albis, versus medium sporulis conglomeratis rubris. *Link, l. c.*

Supra vesicas, quibus orificia lagenarum, praeparata anatomica continentium, clauduntur, repertum. Lanam teneram offert. Sporulae conglutinatae, primo albae, dein amoenae rubrae. *L.*

31. *ollare*, caespitulis tenuissimis effusis, floccis lanatis, sporulis versus medium coacervatis globosis roseis.

Sporotrichum roseum. Link, p. 177.

Nascitur super terram in ollis, ubi plantae hyemales conservantur. Ab initio totum candidum, sporulis evolutis, roseum.

32. *rhodochroum*, caespitulis crassiusculis sublimitatis parvis floccisque raris, sporulis minutis roseis. *Link Anal. l. c.*

Aleurisma roseum. Ejusd. Berl. Magaz. 1815. p. 58.

Etiam terrestre est, praesertim in turfosis provenit. Acervuli colliculosi demum inter se confluunt.

***** *Villo albo, sporulis nigris.*

33. *Collae*, crassum effusum, floccis implicatis stuposis, sporidiis in acervos globosos congestis. *Link, l. c. p. 182. †*

Collarium nigrispermum. Ejusd. in Berl. Mag. p. 17. t. I. f. 30. Nees Syst. d. Schw. p. 47. t. 3. f. 42.

Nascitur super collam siccam, ut lana alba sat crassa, in qua acervuli observantur nigrae ex sporulis seu granulis compositi, sibi arcte agglutinatae. *Lk.*

34. *lyococcos*, effusum laxum, floccis implicatis tenuibus longis albis, sporulis in acervos globosos collectis, in aqua diffluentibus nigris. *Ehrenb. Sylv. Mycol. Berol. p. 22.*

Prov. in Pruno armeniaco.

***** *Colore toto nigro.*

35. *parietinum*, effusum tenue, floccis laevibus vagis, sporulis versus medium collectis globosis. *Link, l. c. p. 181.*

Crêsc. ad parietes calce recentes dealbatos et humidos.

36. *calcigena*, laxum tenue nigrum, sporulis minutis undique dispersis concoloribus. *Link, l. c.*

Latas plagas occupat in cubiculis recentior calce illitis, humorem conservat et attra-

hit, quam ob rem parietes peiores reddit.
Link.

37. *chartaceum*, nigrum, primo orbiculare,
dein confluens, acervulis pulveraceis.

Non infrequens ad chartas scriptorias aeri
humido et pluviis expositas. Caespitulus pri-
mo tenuis circularis, dein pluribus confluenti-
bus, latitudinem acquirunt palmarem.

XXXVIII. *ATHELIA*. Depresso-applan-
ta. *Flocci* intertexti tenues, medio in pellicu-
lum laevem sporuliferam condensati (saepius
fissilem).

Obs. Habitu quodummodo *Thelephorarum*
resupinatarum.

1. *strigosa*, obscure lutescens opacum crassi-
usculum irregulare dense strigoso-villosum.

Corticium strigosum. *Obs. Mycol.* 2.
p. 18.

Passim provenit ad salicum cortices, qua-
rum fissuras explet villo strigoso, 1-2 unc.
lato, membranula sporulifera brevi et colore
paululum dilutiore distinctum.

β. *nuscigena*, irregularis, sordito lutescens
opacum molle, floccis subfuniculatis den-
sis.

Thelephora byssoides. *Syn. fung.* p. 577.

Praesertim in pinetis provenit supra et

inter muscos quos una cum ramulis siccis coadunat in ferruginem.

2. *epidermea*, late effusa tenuissima submembranacea albido-pallens tota glabra.

Non raro observatur supra epidermidem ramorum exsiccatorum quercuum, ubi nonnunquam ad longitudinem spithamaeam est effusa. Vix initium *Thelephorae* cujusdam.

3. *Typhae*, longitudinaliter effusa tenuis demum tota glabra, subfarinacea alutacia.

Hab. ad culmos *Typhae latifoliae* exsiccatos. Ab initio orbicularis alba et byssina.

4. *Salicum*, latissima tenerrima ubique glabra albicans aut subcinerascens.

Ad ligna emortua salicum prope Neuchâtel.

Tres unc. fere lata, margine sinuato.

5. *ochracea*, subnitens compacta, medio dilute ochraceo, margine byssino albicante.

β. pallida, colore dilute pallido.

Hab. ad ligna quibus arcte adhaeret, 2-3 unc. lata.

6. *Taxi*, irregulariter effusa tenerrima, subpalescens grisea.

Hab. ad cortices *Taxi*. Chaillet.

7. *epiphylla*, tenuissima glabra cinerascens.

Pers. Traité p. 67.

Rarius in pagina superiore fol. quercus.
Cavvini.

Irregulariter et magis in latitudinem effusa, unc. 1. lata.

8. *bombacina*, epiphylla tenerrima candida, margine glabro.

Sporotrichum, *bombacinum*, effusum tenuissimum densiusculum albam, floccis tenerimis septatis, sporidiis obscuris. *Link, Berl. Mag.*

Provenit in fol. dejectis, filicinis et quercinis.

9. *lanuginosa*, late effusa lanuginosa albida, pellicula sporulifera brevi.

Hab. ad muscos, fol. putrida et sub cortice ramorum dejectorum. Margo latas byssinas, ab initio *Himantiae* subsimilis, etiam subfibrillosa.

10. *velutina*, longitudinaliter effusa, disco carneo-rufa, margine byssino: fibrillis rectis aut subdivergentibus.

Thelephora velutina. Decand. Flor. Gall. 6. pag.

Hab. ad ligna exsiccata quercina. Ultra 2 unc. longa, $\frac{1}{2}$ lata.

11. *sericea*, disco reticulato-texto pallido fuscascente, fibrillis marginalibus sericeis rectis subdivergentibus sulphureis.

Thelephora sulphurea. *Syn. fung. p. ?*

Hab. ad ramos exaridos, muscosos. Species satis pulchra, a sequente omnino diversa.

12. *citrina*, terrestris, fulvo-sulphurea, primo subfibrillosa. *Traité sur les champ. comestib. p. 67.*

Crescit pulchra haec species ad pedes arborum supra terram et muscos; gregarie crescit, sed vix ultra $\frac{1}{2}$ unc. lata, et lin. crassa.

XXXIX. OZONIUM. *Fibrillae* elongatae (liberae), ramosae, discretas (non intricatae) subtomentosae repentes: primariis crassioribus, teretibus, glabriusculis.

Obs. Himantiis forte nimis affine genus.

1. *croceum*, dispersum flavo-croceum, fibrillis subdifformibus vage ramosis.

Himantia sulphurea. *Syn. fung. p. 703.*

Sporotrichum croceum. *Kunze Mycologische Hefte, 1. p. 81.* (quoad specimen missum).

Hab. in quisquiliis, passim intra terram. Fibrillulae vix $\frac{1}{2}$ unc. longae, in medio saepe difformiter incrassatae.

Obs. An forte status junior *Atheliae citrinae*? sed secundum D. Kunze, sporulis minoribus ovalibus oblecta est haecce plantula.

2. *lateritium*, fibrillis vage ramosis subdifformibus incrassatis glabris sublateritiis. †

Himantia lateritia. *Syn. fung.* p. 704.

Clavaria filiformis. *Sowerb. Engl. fung.*

t. 387. f. 4.

Hab. supra folia exsiccata praesertim *castaneae*, sero autumnino, supra quae aliquando totum effusum et iis fere agglutinatum colore tunc subcroceo, plerumque vero liberum ascendens rarius ramosum, colore ex rufescente, lateritio et spadiceo variat, apice passim albicante. Plantula dubii loci, etiam ad *Isarios* ob formam quodammodo accedit, sed omnino glabra, et substantia sicca est.

3. *stuposum*, subterraneum, magnum subimplicatum opacum spadiceum subglabrum.

Dematium stuposum. *Syn. fung.* l. c.

Byssus intertexta. *Decand. Flor. Gall.*

Hab. in Fodinis et aliis locis subterraneis.

Fibrillae latitudine sensim diminuuntur.

4. *fulvum*, cespitosum densum fulvo-ferrugineum fibrillis rigidis strictis.

Ozonium auricomum. *Link in Berl. Mag. pag. 21.* (quoad communicata specimina).

Dematium strigosum. *Syn. fung.* p. 695.

Byssus fulva. *Huds. Humb. Friberg.* p. 62.

Byssus barbata, *Eugl. Botany*. t. 701.

Hab. ad truncos exsiccatos et humiditati expositos; praesertim sub cortice aut intra huius fissuras, demum saepe canescit.

β. *Gentianae*, filis magis discretis elongatis apice divergentibus, colore dilutiore flavescente.

Hab. ad caules aridos *Gentianae luteae*, in Vogesia. An forte species distincta?

γ. *radians*, dilatata compressa ferruginea, fibrillis centrifugis radiantibus.

Byssus parietina, var. α. *Decand. Flor. Gall.*

Hab. in domibus humidis et obscuris ad parietes.

Species elegantissima cum priorē quoad colorem convenit, sed dispositione fibrillarum, quarum primariae incrassatae et glabrae et minores subimplicatae villosae sunt, ab ea distinguī debet. Ad *Himantias* transitum facit.

XL. HIMANTIA. Repens depressa, (tenuior). *Fibrillae* ramosissimae divergenti-ramosae discretae; omnibus conformibus, minoribus nonnunquam in membranā connatis,

* *versicolores*.

1. *rubiginosum*, mollissimum rufo-spadiceum partim intertextum villosum, fibrillis paucis distinctis crassiusculis teretibus.

Turino misit cl. *Balbis*, ad muros et in cellis habitare videtur, cum particulis calcareis inquinata est et subtus unita ceterum unc.

1. lata et *Racodius* affinis. †
2. *domestica*, maxima subviolaceo-fuscescens mollis in membranam subcohaerens. *Syn. fung. p. 703.* †

Crescit in domibus intra ligna fabrefacta, quae destruit, longitudine nonnunquam insigni, demum fere membranacea. Affinitatem quandam habet cum *Xylostromatibus*.

3. *cellaris*, magna ramosissima villosa nigrescente-umbrina.

Hab. in cellis ad muros, longitudine aliquando bipedali, ramis divergentibus in membranam coalitis in individuis majoribus; ramulis quasi anastomosantibus.

4. *umbrina*, compactior minus divergens ramis cancellatis, umbrina. *Syn fung. p. 704.*

Rarissime provenit ad cortices arborum in Germania. Antecedenti valde affinis. Longitudine palmari.

5. *flabellata*, ramosissima villosa-hirta umbrina, ramis flabellatis apice in membranam concretis.

Fibrillaria ramosissima. *Sowerb. Engl. fung. t. 387. f. 2.*

Species ex icone valde distincta, quae,

uti brevis descriptio dicit, duos pedes habet in diametro, et saepe concentrice crescit. Rami crassi sunt, sed breves.

6. *rosea*, radiato-ramosa rosea, fibrillis crassiusculis, extimis byssaceis.

Himantia Rosae. *Fries Obs. Mycol. I. p. 211.*

Hab. in Suecia in cortice Pini et Juniperi.

7. *helvola*, repens subadnata glabriuscula subcinnamomea: ramulis aequalibus divergentibus: nonnullis fasciculatis.

Hab. ad Ligna, in Germania. *Funck.*

Unc. 1 magna, tenuis. Fibrillae aequales, basi convergunt.

8. *dendritica*, depresso plana adnata glabriuscula pallida, ramulis sparsis divergentibus.

Hab. in Germania. *Esenbeck.*

Species elegantula exacte dendritica, substantia submembranacea a ceteris parum recedit.

** *Albae, epiphyllae.*

9. *plumosa*, epiphylla repens depressa uniformis candida, apice dilatata plumosa.

Himantia candida var. α . *Syn. fung. p. 704.*

Byssocladium candidum. *Link, Berl. Mag. 1815. p. 36.*

Frequens in foliis delapsis praesentim quercinis. Ramoso-dichotoma et undique villosa.

10. *radians*, gregaria subparva, fibrillis demum adscendentibus glabriusculis subsimplicibus, dein pallentibus, basi villosa radiante (fol.) affixa. *Champ. Comest. p. 65.*

Himantia farinacea. *Consp. fung. p. 373.*
Exclus. Synonymo.

Chordostylum hispidulum. *Tod. Fung. Mecklenb. I. p. 39. t. 7. f. 54?*

Equidem non rara in foliis siccis delapsis.

11. *Tela*, ramosissima repens tenerrima farinosa, fibrillis tenuissimis pallidis glabriusculis.

Passim provenit quoque in foliis, quae ad dimidium occupat. Telae arachnoideae e longinquo haud absimilis. Ramuli extimi villosa farinacea. Mihi quoque missa sub nomine *Acremonii verticillati*.

12. *nitens*, albido-nitens subcrustacea, fibrillis obsoletis.

Hab. passim sero autumnno, in sylvis frondosis, inter acervos foliorum exsiccatorum, quae inter se quasi agglutinat et in altum elevat. Fibrillae quidem in pelliculam quasi cohaerent, sed sub lente et nonnullae quoque in margine satis distinctae apparent.

- 13? *pulchella*, tenerrima candida appressa, fibrillis distinctis radiato-divergentibus.

Elegantula haecce planta passim crescit supra folia adhuc viridia Rosae gallicae in campis circa Parisios. Textura valde delicata, vix villosa est. Fibrillulae angustae, ut venae supra fol. pag. superiore sese expandunt. An forte subiculum *Erysiphes cujusdam*?

*** *Albae epiphyllae.*

14. *finbriata*, magna effusa ramosa, filis longis subargenteis, fasciculatim basi congestis, apicibus diffusis applanatis plumosis. *Schumacher, Saell. p. 444.* H. plumosa.

Inter corticem et lignum nec non in truncis cavis putrescentibus haec elegans species nascitur. S.

15. *subcorticalis*, primo villosa divergente radians, demum in membranam latam glabram coalescens.

Hab. praesertim inter corticem et lignum (sed illi adnata) ramorum putridorum, locis humidis jacentium. Saepe guttulas aquae sat magnas exsudat, an forte tunc a b. Tode ut *Myrothecium roridum* descripta?

16. *Ornithogala*, sublata incrustans farinosa, fibrillulas nonnullas liberas exserens.

Hab. ad sepes in quisquiliis, ad ramulos foliaque sicca ibi accumulatos, quae ejus fibrillis inter se cohaerent, colore albo, excremento avium e longinquo simili mox discernitur.

XLI. XYLOSTROMA. Diffusum, coriaceo-suberosum, compactum, laeve, asporum.

1. *radians*, tæne submembranaceum ligno arcte adhaerens, pallide spadiceum, a fibrillis obsoletis radiante-striatum, dein compactum nigrescens.

Hab. ad ligna dura Juglandis regiae.
Chaillet.

Substantia sicca, membranacea, glaberri-
ma, quae ex fibrillis conflata est, harum enim
nonnullae, colore spadiceae, radiatim promi-
nent. Demum in areolas aut frustula dilabi-
tur, nec non nigricat. Haecce plantula parum
dubia et ulterius indaganda.

2. *cinereum*, crassiusculum laeve dilute cine-
reum.

Hab. in Gallia. Valde compactum, utrin-
que planum, 1-2 lin. crassum, nonnullas unc:
latum.

3. *candidum*, orbiculare confertum tenuissi-
mum candidum glabrum lignum ubique pe-
netrans.

Repertum in ligno carioso quercino. Conf.
Racod. *papyraceum* et obs. ibi adjectam.

4. *loreum*, coriaceo-membranaceum tenue ru-
fo-spadiceum.

Hab. in Gallia. Lora exacte simulat. *Spi-
thamum* longum, 1-2 unc. latum.

5. *ramentaceum*, tenuissimum parvum laeve ferrugineum.

Ad ligna exsiccata. Unc. et quod excurrit magnum. An vera *Xylostromatis* species?

6. *Corium*, latissimum coriiforme laeve pallide alutaceum flexile.

Xylostroma giganteum. *Tode fung. i. p. 36. t. 6. p. 51.* *Sowerby fung. t. 358.* (optima). *Syn. fung. p. 702.* sub *Racodio*.

Hab. in interiore truncorum *quercuum*. Corio simile, ab Anglis hinc *Oak-leather* nominatum. Nunc ligno arete adnatum, nunc liberum, ad longitudinem nonnunquam insignem extensum. Substantia et colore paululo differt ut et crassitudine.

7. *Suber*, latissimum crassum suberosum durum dilute alutaceum, inferne spadiceum.

Inventum in cupa, ubi vinum fuit fabricatum, et cujus fundum fere totum explevit. Communicatum a D. *Thiaillier*.

Tres quatuorve lineas crassum; superficies in specimine inaequalis, hinc inde lacunosa. Color in pagina inferiore, quae scabra est, fere nigrescit, et verosimile ab uvulis rubris residuis ita mutatus.

ORDO SECUNDUS.

SARCOMYCI (*Hymenomyci*, *Pilomy-*
ci) seu fungi stricte sic dicti, textura sunt
stipati, substantia carnosi, coriacei vel tre-
mellosi, exceptis nonnullis speciebus prae-
sertim *Thelephorae*, *Boleti* et *Hydni*, quae
nonnunquam praesertim versus ambitum
tollae villosae sunt. Magnitudine plerum-
que ceteros superant, nec non formam ha-
bent determinatam, quae est aut teres vel
elongata, plerumque plusminusve orbicu-
laris et *pileus* (cum ejus varietatibus) au-
dit, hic, saltem in majoribus et terrestri-
bus, suffultus est pediculo (*Stipes*). *Se-*
minula (*Sporulae*, *Sporidia*) proferunt
parca in copia, nonnunquam solummodo
prunae ad instar, in omni aut in una al-
terave superficie, quae antea inclusa fue-
runt in *thecis* membranaceis, oculi inermi
vix percipiendis, quae vero dense adpro-
pinquatae sunt, ut membranam homoge-
neam efficere videantur, et *hymenium* de-
nominatam; hoc enim vel laeve est vel
tuberculosum, aut porosum et aculeatum,
saepius vero plicatum et lamellosum.

† *Tremelloidei*, substantia, uti nomen in-
dicat, sunt molles, tremellosi, aut interdum

carnoso - coriaceo - tremellosi (non fluxiles), nunc laeves, nunc rugoso - plicati, forma satis varii, plerumque vero minus regulari; sporulas quoque (nonnullis forte exceptis) secernunt quae vero thecis (in omnibus?) sunt destitutae.

XLII. MYCODERMA. Orbiculare, coriiforme, primo molle, subpellucidum dein induratum, substantia ubique aequali. (Aspermium? natura Mucedinum).

1. *ollare*, fusco - badium magnum, utrinque plicis densis rugosum, demum fragile.

Repertum in olla quadam, ubi folia *Rumicis Acetosae* cocta asservata fuerunt.

2. *mesentericum*, crassiusculum album subviscosum, utrinque undulato - plicatum, mesenteriforme.

Supra orificium ruptura ortum in lagena vino adhuc plena reperta a cl. *Chaillet*. Aqua irroratum, carnosum et ad tactum viscosum. Conf. *Traité sur les champ. comestib.* p.

3. *Lagenae*, rubens orbiculare subpeltatum laeve, inferne obsolete rugosum.

Eodem modo uti antecessens natum et mecum ab eodem inventore benevole communicatum, cum multis aliis fungis raris.

4. *pergameneum*, orbiculare tenuissimum album superficie exasperatum.

- Dete-

Detectum a Dom. Desmazieres, Lilloae, supra succum cerasorum, inclusum in lagena, ubi pellem sistit, 2-3 unc. latum.

Obs. Hae enumeratae formae originem ceperunt in locis a ceteris fungis majoribus ita alienum ut adhuc dubitandum, an revera ad horum vegetabilium familiam pertineant; cum vero facie et substantia fungosa omnino gaudent, hinc eas silentio non praeteriri existimavi.

XLIII. *AURICULARIA*, sessilis, coriaceo-tremellosa in superficie sporulifera (disco) vage plicato-venosa, plicis paucis divaricatis, subtus tomentosa.

1. *mesenterica*, subimbricata, pileo demum reflexo fasciato fuscescente griseo, disco purpurascente plicato.

Thelephora mesenterica. *Syn. fung. p. 571. Bolt. fung. Append. t. 172. Michel. gen. t. 66. f. 4.*

Auricularia tremelloides. *Bull. champ. t. 290. (nala).*

Hab. ad truncos arborum, praesertim *Juglandis*. Fungus junior resupinatus, late effusus. Pil. superne griseo- aut cinereo-brunescentis, zonis concentricis distinctus et hirsutus. Disens in maturitate a sporulis albo-farinosus.

2. *Sambuci*, resupinata concava, utrinque ve-

noso - plicata externe tenui tomentoso fuscescente, interne nigrescente.

Tremella Auricula. Syn. fung. p. 6. 14.

Peziza Auricula Lin. Bull. t. 427. f. 11.

Hab. ad truncos *Sambuci nigrae*; saepe gregaria. Sat magna. Plicae in pileo qui non zonatus est magis distinctae sunt quam in disco, qui nigrescit.

XLIV. TREMELLA. Gelatinosa, forma varia, undique sporulifera. Pileus distinctus nullus. Sporulae demum emersae, thecis distinctis non inclusae *).

* ENCEPHALIUM, capitulo subrotundo plicato-gyroso, continente nucleum compactum.

1. *encephala*, sessilis subcompacta carneo-fuscescens. — Willd. Bot. Mag. vol. 2. fasc. 1. p. 17. t. 4. f. 14. Albert. et Swein. Conspect. fung. p. 301. (Descr. optima.)

Encephalium aurantiacum. Stroma ambitu plicato gelatinoso, nucleo duro compacto,

*) Hocce genus comprehendens species forma valde diversas, in posterum dividendum. Illud sub nomine: *Encephalium* a cl. Link a ceteris separatum, quamvis quoad habitum externum cum nonnullis ex divisione secunda conveniat, tamen ab iis differt nucleo quodam, typo forte pilei; sed *Tubercularia vulgaris* a ceteris etiam nucleo distinguitur.

sporidiis in ambitu sparsis demum emersis.
(Char. gener.) *Link in Berl. Mag. 1815. p. 33.*

Nematelia encephala. Fries Obs. Mycol.
2. p. 370.

Hab. in ramis *Pini sylvestris.*

Obs. B. Willdenow hanc speciem oblongam, sessilem, nec non pallide carneam, exsiccando vero fusciscentem descripsit: *cl. Schumacher (Flor. Saelland. p. 433.)* ei attribuit stipitem brevissimum, basi dilatatum et colorem subumbrinum, qui vero secundum *cel. Link* est aurantiacus. Species hinc videtur variabilis, aut ex pluribus composita.

2. rubiformis, gyrosa lutea. Fries Obs. Mycol. 2. p. 370. (Naematelia).

Hab. cum priore rarior. Absque descriptione. †

** *GYRARIA. Effusa (major), plicata gyrosa, vel undulata, submembranacea.*

3. fragiformis, subcompacta rotunda rubra, sublobata, plicis confertis. Syn. fung. p. 622. Icon. pict. fung. fasc. 2. p. 23. t. X. f. 1.

Crescit quoque ad cortices *Pini*. Intus colore pallidiore (an nucleo?) distincta, hinc omnino examinari meretur, an *Encephalii* species sit.

4. mesenterica, plicato-undulata flavo-aurantiaca. Syn. fung. p. 622. Jacq. Miscell. au-

striaca. 1. p. t. 13. Vaill. Botan. t. 14. f. 4.

Satis frequens prorumpit initio hyemis, post pluvias, e ramis exsiccatis, lignis et sepi-
mentis $\frac{1}{2}$ unc. ad 1 unc. magna, nunc subro-
tunda, nunc oblonga, passim subcompressa pro
loco natali; colore plus minusve flavo-auran-
tiaco. Substantia subtenax.

β ? *lutescens*, gyrosa mollissima fluxilis (?)
pallide flava. Syn. fung. l. c. Ic. et Descr.
fung. p. 33. t. 8. f. 9. Decand. Flor. Gall.
Suppl. p. 28.

Autumno in Germania rarius eam inveni
ad ramum fagineum, nec denuo, nec alibi.
Forte nonnullis ab auctoribus antecedens sub
hoc nomine descripta vel enumerata est spe-
cies.

5. *Helvelloides*, gelatinoso-tremula roseo-au-
rantiaca, erecta, stipite basi sulcato frondes-
cente, margine sinuato. Decand. Syn. pl.
Gall. p. 19.

Hab. ad terram subhumidam in fagetis.
An hujus divisionis? Sex unc. alta. Capitul.
erectum, aut nutans. D. †

6. *candida*, subparva rotundata gyrosa alba.
Syn. fung. p. 624.

Passim ad ramos siccos tum in sepi-
bus, quam in sylvis, autumno et hyeme. Parva,
vix 4 lin. lata, erumpens.

7. *violacea*, subcompressa parva compacta gyrosa violacea. *Syn. fung. p. 623.*

Hyeme ad truncos Pyri communis; ex quibus erumpit, ad quatuor lineas longa, vix unam crassa.

8. *tingtoria*, magna cespitosa undulato laevis vinoso - violacea, exsiccata (fragili) fusco-purpurea.

Tremella mesenteriformis, var. 4. *violacea. Bull. Hist. d. Champ. p. 230. t. 499. f. VI.*

Hab. ad truncos, rarius. 2-3 unc. magna, foliacea. Tincturae inservire potest ex Bulliard.

Obs. An *Tremella undulata Hoffm. veg. crypt. 1. t. 7. f. 1.* hujus loci?

9. *foliacea*, magna cespitosa utrinque laevis undulato - plicata basi crispa cinnamomeo-carnea. *Syn. fung. p. 626. Obs. Myc. 2. p. 98.*

Hab. in umbrosis ad truncos. Subpezizoida.

10. *succina*, cespitosa tenuis pellucida plicata fuscescente succina.

Tremella mesenteriformis. Bull. Champ. t. 499.

Rarius ad ligna exsiccata observatur autumno seto. Substantia tenuis, tremula, magnitudine unciali. A *Tr. mesenterica* quam maxime diversa.

Cespitem non tam densum eam formantem inveni, qualem depinxit beat. Bulliard. Antecedenti valde affinis.

11. *salicum*, gregaria orbicularis peltata submembranacea pellucida disco late plicata fusco-succina.

Hab. in Germania ad ramos exsiccatos salicum. D. Ehrenberg.

Exsiccata valde contrahitur, difformis fit et fere nigrescit; aqua irrigata formam genuinam acquirit satis gratam auriculae subsimilem; $\frac{3}{4}$ unc. fere lata, plicato rugosa, subnitens.

12. *impressa*, explanata impressa subconchiformis, recens laevis opaca nigrescente-fusca.

Tremella rufescens. Ehrenberg (ined.)

Unc. lata est haecce species, crassiuscula sicca rugosa, sed non valde diminuitur, et rufo fusca fit; humectata tota laevis, aut tantummodo versus marginem parum plicata, subtus uti videtur sterilis est.

13. *spiculosa*, effusa applanta nigra papillis conicis spiculosa. *Syn. fung.* p. 624.

Tremella glandulosa. Bull. Champ. t. 42. f. 1.

Provenit sero autumnno ad ramos putridos, supra quos longitudinaliter est effusa. Ab initio suborbicularis plana, et tunc magis plicata videtur. Inferne tomenti tenui oblecta hinc

generi *Auriculariae* valde affinis, tamen *Tremellarium* faciem habet. Ab hac *Trem. papillata* Kunze *Mycol. Hefte*, p. 86. non diversa videtur.

*** *TREMISCUS*, erecta dimidiata, subauriculata.

14. *sinbriata*, umbrino - fuliginea suberecta, latero incisa, margine undulata. *Syn. l. c.*
Obs. Mycol. II. p. 97. — Hoffm. veg. cryptog. I. t. VII. f. 1. 2.

In ramis putrescentibus, locis humidis, passim. Sursum dilatata.

15. *rufa*, magna erecta carneo - rufa dimidicata - infundibuliformis, marginibus laevibus subcinnatis. *Jacq. Misc. Austr. I. p. 143. t. 14.*

Hab. in Austria et Helvetia.

Obs. Species singularis forte sui generis, nam solummodo interne fructificare videtur. Ultra 2 unc. longa, versus basin canaliculata. Juniores ligulatae, adultiores subauriculatae vel spathulatae.

**** *AGYRIA*. *Pezizae* — aut tuberculiformis, recens laevis, aut subrugoso plicata. (Species plerumque parvae.)

16. *reçisa*, mollis obconica truncata plana aut subconvexa subrepanda fusco - succina, stipite obliquo. — *Ditmar in Sturm Flor. German., cryptog. Fung. fasc. I. p. 27. t. 13.*

Peziza gelatinosa. Bull. champ. t. 460.
f. 2. Syn. fung. p. 633.

Provenit ad ramulos aridos, praesertim salicinos. *Pezizam* exacte refert. Maxima hujus subdivisionis. Exsiccata valde diminuitur et spadicin evadit.

17. *Urticae*, conferta minuta plana aut excavata flavo-rubra. Syn. fung. p. 628.

Vulgaris ad caules exaridos *Urticae dioicae*, ubi est immersa, saepe in seriem longitudinalem crescit et applanata est. — A reliquis speciebus floccis rectis parallelis differt, quippe qui in aliis magis contorti sunt. Link.

18. *abietina*, gregaria parva subrotunda convexa aurantiaca. Syn. fung. p. 627.

Satis frequens ad ligna abietina fabrefacta, hyeme.

Tuberculiformis, laevis, immutabilis, rarius subplicata, lin. lata.

Obs. Flocci intertexti septati albi, sporidiis creberrimis aurantiis globosis interspersis. Link, Obs. 2. p. 33. A cl. Nees ab Esenbeck cum sequente sub nomine *Dacryomyces stillatus* (Fung. 1. p. 89. t. 90.) conjuncta.

19. *lacrymalis*, subrotunda aut inaequalis subplicata lutescens nitida. Syn. fung. p. 628.

Tremella deliquescent. Bull. champ. t. 453. f. 3.?

Autumno passim ad ligna quercina crescit. Ab initio globosa laevis, demum oblonga, subconfluens quasi lobata. Recens flava, exsiccata fuscescit, et valde tenuis depressa fit.

20. *hyalina*, congesta hyalina alba subrotunda oblongaque laevis aut subplicata.

Autumno passim ad ramos cariosos quercinos, his in regionibus, inveni; 2-5 lin. magna est.

Obs. An a *Tr. candida* sufficienter distincta?

21. *finetaria*, subexigua gregaria gelatinosa orbiculata depressiuscula tuberculosa fusco-carne subdiaphana sessilis. *Schum. l. c.*

In fimo vaccino. Jul. in Dania. †

22. *Syringae*, minor gregaria subconfluens carnosa orbiculata subtuberculosa aurantiorubra, stipite brevissimo superne dilatata. *Schum. Flor. Saelland. p. 440.*

In rimis truncorum *Syringae* vulgaris. Decemb. †

23. *alutacea*, gregaria subconfluens carnosogelatinosa suborbiculata minutula subgyrosetuberculosa, alutacea, stipite brevissimo cylindrico. *Schum. l. c. p. 439.*

In ramulo subputrido, in Dania reperta. †

Obs. Tres hae species, mihi plana ignotae, ob discum tuberculosam et stipitis pre-

sentiam forte alius sunt generis, et denuo examinari merentur.

***** *Coryne* (Nees). *Elongata, teres, erecta, clavaeformis.*

24. *clavata*, solitaria erecta simplex incrassata carneo-rubra. *Syn. fung. p. 630. Icon. pictae fung. fasc. 2. p. 24. t. X. f. 1.*

In ramis putridis. Unc. 1 alta, 2-3 lin. crassa.

25. *cornuta*, gregaria subcompressa ramoso-cornuta obscure carnea. *Syn. fung. l. c.*

Vere ad truncos, aequae ac rare ac antecedens.

26. *dubia*, cespitosa viscosa fastigiata mollis carneo-subalbicans. *Syn. fung. l. c. Schaeff. fung. t. 323. f. 1. 3-6.*

Coryne Acrospermum. Nees fung. t. 15. f. 143.

Hab. autumnno ad caudices putridos. Affinis *Pezizae sarcoidi*. Forma variabilis.

27. *aculeiformis*, gregaria fasciculata flavescens simplex ramosaque basi connata latiore. *Syn. fung. p. 596.*

Clavaria cornea. Batsch El. fung. f. 161. Clavaria aculeiformis. Bull. champ. t. 463. f. 4.

Hab. in lignis putrescentibus, vero autumnno. Minuta, 2-3 lin. alta.

28. *ferruginea*, gregaria submagna carnosio-gelatinosa fusco-ferruginea simplex passim bifida flexuosa, basi angustata. *Schum. Flor. Saelland. p. 441.* †

In ramulis dejectis semiputridis, in Dania Hyeme.

29. *palmata*, solitaria carnosio-subcoriacea aurantio-lutea (fulva) compressa dilatata apice ramoso: ramulis subteretibus divaricatis. *Schum. Saelland. 2. p. 442.*

Hab. in ligno quercino, autumno. Magnitudine media, basi angusta. Ramuli obtusi. *S.*

30. *ligulata*, gregaria subminuti rugulosa subcompressa obtusa diaphana pallida. *Schumach. L. c.*

Inventa, in Dania, in stipitibus Pteridis aquilinae. Clavata $1\frac{1}{2}$ lin. longa.

31. *glossoides*, flava parvula simplex subtremellosa, clavula ligulata, caul. tereti. *Syn. fung. p. 596.*

Apparet rarius in ligno exsiccato subputrido quercuum. Altitudine trilineari.

32. *subclavata*, subsolitaria media clavata subgelatinosa, apice subtuberculosa sordide lutea, compressiuscula, basi teres. *Schumach. L. c. p. 442.*

33. *applanata*, submagna applanata elevata tuberculosa subgyrosa lutea. *Schum. L. c.*

In ligno subputrido quercino. Novembr.
 $\frac{1}{2}$ unc. longa.

***** *Acrospermum*, apice capitulo sporulifero distinctum ibique colore plerumque dilutiore.

33. *galeata*, gregaria e fuligineo subpurpurascens, capitulo subalbicante. *Comment. de fung. clavaef.* p. 91. *Flor. Dan. t. 1100. f. 2.*

Helotium galeatum. Syn. fung. p. 628.

Clavaria galeata. Holmsk. Fung. Dan. p. 25. ic.

Autumno passim ad ramulos dejectos, locis praesertim humentibus. Unc. 1. fere magna.

34. *cylindrica*, solitaria submagna carnosogelatinosa elongato-clavata purpurea, apice obtuso dilutiore. *Schum. l. c. p. 441.*

Acrospermum cylindricum. Flor. Dan. t. 1076. f. 4.

In ligno putrido et ramis dejectis. Pini *Abietis* et *Betulae albae*, in Dania.

Berea, subflexuosa, basi angustata purpureo-nigricans et fuliginosa.

β. cornuta, obscure purpurea, basi stipitiformi nigra, superne bi-trifido, ramulis subdivaricatis simplicibus subincisisve tuberculatis compressis. *Schum.*

Hab. cum antecedente, a qua ut species forsā distingui meretur.

***** *Leioderma*, tremelloso - coriacea, laevis, exsiccatione immutabilis, ligno arte adnata, applanata.

35. *elliptica*, aquose pallida elliptica tenuis.

Semel ad caules aridos *Urticae dioicae* mihi obvia, sed tunc sat magna in copia. Lin. 1 - 2 longa.

36. *linearis*, minuta elongata linearis pallens.

Aequae rara in ramulis dejectis quercinis, in quibus ut striae apparent. Substantia mollior.

37. *Arundinis*, majuscula subcoriacea nigrens demum subrimosa.

Ad culmos exsiccatos *Arundinis Phragmitidis*. *Chaillet*. Unc. 1 effusa, subinterrupta opaca, tenuis.

†† *THELEPHOREI* fungi substantia sunt firmiter coriacea, nonnullis exceptis quae aut toti aut ad marginem byssoidei vel textura tomentosa sunt, et forma magis regulari, sed pileo bene distincto nondum gaudent: plurimi enim effusi aut effusoresupinati sunt, pauci imbricato - horizontales, nonnulli integri, infundibuliformes, nonnulli clavaeformes ramosi. In plurimis perspicuum adest hymenium, quos, in superficie est tuberculosum vel papillosum, aut papilloso - pulveraceum, in aliquibus interea omnino laeve, in perfectioribus ex thecis compositum, aut sporulis tantum adpersum.

XLV. THELEPHORA. *Pileus* forma varius. *Hymenium* papillis, passim simul setulis distinctum, vel omnino laeve.

* POLYFILUS, stipitatus, cespitosus. *Pilei* subcarnoso - coriacei, numerosi horizontales, aut iufundibuliformes. *Hymenium* laeve. (*Species majusculae, terrestres, pallidae.*)

1. *frondosa*, maxima stipitata forma varia, stipite tuberoso difformi, pileis imbricatis undulato - crispis glabris pallescentibus.

In Gallia ad pedes *Pini maritimae*, mensibus Octobri, Novembri crescens detecta a D. *Cauvin* *Descr.* Boletum (Polyporium) *frondosum* quodammodo refert, sed minor, spithamam et ultra est haec species. *Stipes* simplex, aut ramosus, compresso - irregularis, 2 uncias longus, unc. 1 crassus, terram ad 2 uncias fere intrat, basi subtuberosa paucasque fibrillas emittens. *Pileoli* lobiformes numerosi, dimidiati, tenues, substriati, margine incisi, suberistati; nonnulli integri, subinfundibuliformes, recenti in statu subalbidi. *Hymenium* laeve, aut subplicatum, colore intensius. *Odor* gratus *Agarici campestris* aut *Morchellae exculenta*.

2. *intybacea*, stipitibus elongatis basi unitis, pileis imbricatis fibroso - squamosis, subtriquetris pallidis, subtus fusco - cinereis. *Syn. fung.* p. 567.

Olim in fagetis circa Gottingam hunc fungum inveni, cespitem efformantem obconicum, 3 unc. latum. Cum vero denuo sese mihi non obtulit, plura de illo in medium proferre hactenus non possum.

3. *pannosa*, cespitosa pallide flavescens, pileo infundibuliformi margine lacero.

Helvella pannosa. Sowerby fung. t. 155.

Cresc. in Anglia. Pil., in specimine ab auctore recepto, margine plus minusve lacernis subglaber, et inferne setulis destitutus; an etiam aliquando integerrimus? Stipes sublongus. Sequenti affinis.

4. *pallida*, cespitosa pallida, pileo integro infundibuliformi squamoso strigoso, hymenio setulis distincto. *Syn. fung. p. 565. Ic. et Descr. fung. Fasc. I. p. 5. t. 1. f. 5.*

Hab. in Germaniae umbrosis, sed rarius. Stipes basi villosus. Pileus ex pallido rufescens, cum vicinis saepius connascens, inferne longitudinaliter striatus et setulis hispidus, sed non papillosus.

** *PHYLACTERIA. Pileus coriaceo-membraneus, submollis, effusus, imbricatus, aut passim infundibuliformis. Hymenium sublaeve, vel granulosum, et in fungo recenti setulis? brevissimis sporulosis? quaternatim aggregatis, distinctum. (Species terrestres, involventes, fusco-purpurascens, nigricantes, aut griseo-cinereae.)*

5. *elegans*, pileo integro subimbricato tomentoso albo, margine nigro - zonato, inferne fusco - purpurea.

Auricularia elegans. Sow. fung. v. 3. t. 412.

Prov. in Angliae sylvis. Pilei dilatati excentro communi divergentes et tomento candidissimo obtecti, ad marginem lineolis nigris pictis, subtus prima grisea adpersi. Sow.

6. *caryophyllea*, cespitosa fusco - purpurea, pileo infundibuliformi striato - fibroso, margine erecto albicante, saepius inciso. — Syn. fung. p. 565. Schaeff. t. 325.

Auricularia caryophyllea. Bull. t. 403. f. 6. (mediocris.)

β. *laciniata*, imbricata obscure fusca aut badia, pileo tenui laciniato crispo, subtus papillis congestis scabro. Syn. fung. p. 567.

Auricularia caryophyllea. Sowerb. t. 213. huc magis pertinet.

γ. *ambigua*, minor gregaria, pileo ut plurimum inciso, subtus laevi fusco, superne striato.

Obs. Mycol. I. p. 39. t. 6. f. 8 et 10.

Merulius radiatus. Holmsk. Fung. Dan. 2. p. 56. t. 29. Nees ab Esenbeck System. t. 34. f. 250.

Peziza radiata. Flor. Dan. t. 469. f. 2.

Hab. in sylvis frondosis, magis in pinetis, ad terram inter folia et stipitos, cum quibus

bis saepius connata. Pileus subflaccidus, convolutus, nunc integer, nunc uno latere incisus, aut ad marginem dissectus, plus minusve tamen verticalis. Stipes brevis.

7. *terrestris*, cespitosa fusco - purpurea, pileo depresso fibroso - strigoso dimidiato, margine albicante. *Syn. fung. p. 566. Nees System. f. 251.*

Agaricus tristis. Batsch Elench. fung. t. 24. f. 41.

Helvella caryophyllea. Bolt. fung. t. 175. ?

Etiam ad terram crescit, plerumque vero ad frustula lignea terra tecta, amatque, uti antecedens, solum arenosum. Pileus plerumque imbricatus aut cum vicinis margine connatus, subcrassus, horizontalis, hinc stipes, qui brevissimus, plerumque lateralis est, aliquando centralis, pileo tunc subintegro: *Thel. dubia. Syn. fung. l. c. — Priori valde affinis est.*

Obs. Thelephora mesenteriformis, substipitata imbricata floccoso - fasciata fusca, margine laciniato albido. *Willd. Berol. t. 7. f. 15.* ab auctoribus plerumque huc adducta, ad precedentem potius referri debet.

8. *palmata*, subcarnosa mollis, pileis imbricatis digitato incisis fusco - purpureis.

Clavaria anthocephala, coriaceo - molliuscula ferruginea, basi sublanata, apicibus albidis flabelliformibus umbellatis. *Bull. champ.*

p. 197. t. 451. f. 1.?, quae est Thel. *anthocephala* Swartz.

β. *mollissima*, subeffusa laciniata carnosae subtomentosa albida, inferne glabra fusco-purpurea. *Syn. fung.* p. 572.

Priorem varietatem in sylvis circa Parisios et alteram olim prope Gottingam, unica tamen vice, inveni, hinc incertus haereo, an fungi fuissent aut potius juveniles, ut conjicio. Cum *Merismate foetido* etiam multam habent affinitatem, sed superficies inferior laevis et hinc solummodo sporulifera est.

γ. *fusco cinerea*, late effusa serpens, margine demum subreflexo tomentoso.

α. *fusca*, tota obscure fusca, basi plicata.

Auricularia Phyllacteres. *Bull.* p. 286. t. 436. f. 2.

Thelephora umbrina, terrestris lata effusa mollis umbrina, margine subtomentoso albicante. *Syn. fung.* p. 578. eadem videtur species.

Thelephora crustacea, subcarnosa effusa glomerata fusco-umbrina subtuberculosa, papillis minutis sparsis obscurioribus. *Schum. Saell.* p. 396., quae in Daniae umbrosis ad terram crescit, non valde diversa est.

Hab. aestatis fine in sylvis plerumque ad basin arborum, aut truncorum, latitudine saepius ultra palmari serpendo sese extendit. Colore opaco, spadiceo, demum vero nigricat.

β. murina, late effusa grisea, margine subimbricato reflexo dilutiore. — *Bull. t. 436. f. 2.?* sed *flavescens picta*.

Thelephora caesia, suborbicularis terrestris laeviuscula caesio-cinerea. *Syn. fung. p. 579. Obs. Mycol. I. p. 15.* forte junior adhuc fungus, aut ob regionem magis borealem, ubi lectus erat, minus evolutus.

Crescit promiscue cum *α*, a qua vix specie differt, colore quoque obscurior evadit. De singulari setularum aut sporularum, quae hac in specie magis distincta apparet, confer Librum meum, cui titulus: *Traité sur les champignons comestibles, etc. à Paris, 1718. p. 70. et Bull. Hist. des champ. p. 44. t. 2. f. 9. h.*

*** *EPICRUS*. *Pileus membranaceus, simplex. Hymenium laevissimum, epapillosum. (Fungilli hujus generis (?) minuti et in muscis majoribus crescunt. Meruliis muscigenis affines, et plerumque albidii vel pallidi sunt).*

10. *vulgaris, α. candida*, gregaria minuta dimidiata candida. *Tab. VII. f. 5.*

Thelephora muscigena. Syn. fung. p. 572. Decand. Flor. Gall. 2. p. 105.

β. albido-grisea, pileo applanato subsericeo postice adnato albido-griseo, inferne subrufescente.

Hab. autumnno et hyeme leni in *Hypnis*,

agmine saepius denso; β . sat vulgaris. Vix lin.
2 lata.

11. *lutescens*, gregaria concava lutescens, ex-
terne albicans.

Rarius eam in muscis inveni, aestate. For-
mam habet magis regularem, fere uti *Peziza*.
Hymenium lutescens aut subochraceum.

**** STEREUM. *Pileus* coriaceo - suberosus.
Hymenium tuberculis aut papillis distinc-
tis, passim simul setulis gaudens. (Spe-
cies hujus sectionis verae *Thelephorae* sunt,
et plerumque in truncis aut ramis aridis pro-
veniunt.

† *Pileo imbricato horizontali, aut effuso reflexo,*
marginе libero. (Hymenio minus papilloso).

12. *hirsuta*, imbricata, pileo substrigoso-hir-
suto zonato pallido, inferne lutescente, fus-
co aut griseo. *Syn. fung. p. 570.*

Auricularia reflexa. Bull. champ. p. 202.
t. 483 f. 3. et 4. Sowerby fung. t. 27.

Boletus auriformis. Bolton fung. t. 82. f. a.
 β . *Cellularia cyathiformis. Flor. Dan. t. 1450.*
Sowerb. t. 91.

γ . *decipiens*, effusa resupinata glaberrima laevi-
gata aurantiaca. *Fries Obs. Mycol. I. p. 153.*

In hortis, sylvis et sepibus ad truncos,
ramos aut palos omnium vulgaris.

More congenerum, aetate juniore resupi-
nata oppressa et simplex, adulta uno latere re-

flectitur et horizontalis fit: lobis plus minusve numerosis imbricatis. Hymenium ejus laeve est et glabrum, sed colore mutabile nunc lacte flavum aut subaurantiacum, nunc fuscescens aut cinerascens, plerumque vero pallidum. Var. β . pileo integro infundibuliformi rarissime occurrit.

15. *sericea*, tenerior effuso, imbricata albicante-grisea, pileo sericeo villo brevi adpresso, subtus pallescente-incarnato. — *Schrad. Spicileg. Flor. Germ. p. 184.*

Thelephora nitida. Disp. meth. fung. p. 30.

β . *sanguinolenta*, imbricata dimidiata effusaque ad tactum cruentata, pileo albido-pallente hirsuto, margine subundulato, hymenio glabro griseo-fuscescente. *Conspect. fung. p. 274.*

Hab. ad truncos et ramos Pineos et Abietinos.

Antecedenti nimis affinis videtur et tanquam ejus varietas in *Synopsi* habita; interea tenerior et magis delicata, nec non pallidior et nitore quodam instructa. Varietatem β ., quoad specimina ab auctoribus mecum communicata, non ut speciem distinctum puto, et colore intensiore differt, sed characterem ejus particularem adducere placet: cruore, ad tactum fungi et praesertim ad pressuram, emissi copia pro aeris conditione diversa, tempore enim pluvio-

so creber prodiens, digitum et aquam, cui immersus fuit fungus, tingit."

14. *ochroleuca*, imbricata effuso-reflexa zonata pubescens subtus ochracea. *Fries Obs. Mycol.* 2. p. 276. (excluso synonymo Bulliard).

Auricularia pallida, imbricata, pil. reniformi plicato hirsuto pallido, zonis subimpres-
sis paullo obscurioribus, margine sublobato,
hymenio ochraceo. *Schum. Saelland.* 2. p. 398.
(Ex Fries).

In truncis Betulae et Fagi passim. Diver-
sa a Thell. *hirsuta* praecipue colore hymenii
ochraceo. Pubescentia etiam tenuis, nec stri-
gosa. *Fr.*

15. *tabacina*, pileo imbricato tomentoso ferru-
gineo nitido, margine undulato crispo albi-
cante, subtus glabro fulvo-livido.

Auricularia tabacina, *Sowerb. fung. t.* 25.

Auricularia reflexa, var. *variegata*. *Bull.*
champ. t. 1. p. 282. t. 483. f. 5.

Satis frequens occurrit, sero autumno, ad
truncos caesos quercinos. Species pulchra. Pi-
leus subtenuis, coriaceus, zonatus, basi minus
effusus, eleganter undulatus, margine ultimo,
aetate praesertim juniore, albicante, subtus lae-
vis et glaberrimus.

α. *chrysoloma*, imbricata tenuis, superne fus-
co-grisea, inferne ferruginea, margine re-
pando aureo.

Helvella nicotiana, parasitica ferruginea tenuis flexuosa, margine lobato crispato, nudo aureo, subtus venoso pallide ferrugineo. *Bolt. Fung. p. 174. t. 174.*

Hab. secundum Boltonium ad ramos et truncos *Syringae vulgaris*, supra quos serpit.

β. pulchella, parvula gregaria effuso-reflexa, pileo strigoso-hirto fuscescente, margine undulato albicante, subtus albido-livescente.

Eam his in regionibus inveni, hyeme leni (1817.) ad truncos caesos *Juglandis regiae*. Pilei lobi numerosi, verticales et inter se distant, 3 ad 4 lineas solummodo lati, non zonati, substantia tenues subcoriacei. Distincta forte species.

16. *variegata*, effuso-reflexa utrinque ferruginea, pileo tenui tomentoso fasciato, subtus setuloso.

Thelephora variegata, imbricata, supra sericea fasciata variegata, subtus pilosa ochracea. *Schrad. Spicil. Flor. German. p. 185. β. striata.*

Thelephora ferruginea. Syn. fung. p. 569. (exclus. Syn. Sowerbei).

Hab. in Germania ad truncos praesertim abietinos. Pileus unicolor, saltem meo in specimine, basi vero obscurior subspadiceus, et margine dilutius ferrugineus. Antecedenti tamen ita similis, ut setulis hymenii distingui

praecipue possit. — An forsan *Helvella nico-*
tiana Bolt. potius huc trahenda?

17. *rubiginosa*, imbricata zonata rubiginoso-
spadicea subglabra rigida, subtus papillis ma-
jusculis setulisque interjectis. — *Syn. fung.*
p. 567.

Thelephora ferruginea. *Decand. Syn. pl.*
Gall. p. 21.

Auricularia ferruginea. *Bull.* t. 378. *So-*
werb. t. 26.

Stereum rubiginosum. *Fries Obs. My-*
col. I. p. 155.

β. *Stereum corticiatum*, resupinatum effusum
papillosum confluens concentrice zonatum.
Fries l. c.

Nitida haec species, equidem non rara,
nascitur in nemoribus ad latera veterum trun-
corum, muscis tectorum. Plures pilei, sat elon-
gati, sed parum angusti, sibi invicem dense
impositi; juniores subtomentosi aut velutini,
margine laete ferrugineo; adulti denudati et de-
mum obscure badii fiunt. Substantia hac in spe-
cie, prae ceteris rigida, hinc nomen *Ehrhar-*
tianum Th. *fragilis* sat bonum.

Obs. Species *Sterei* ob setulas, quae po-
tius *paraphyses* quam verae *thecae* videntur,
a *Thelephoris* separatae, nullum in habitu prae
se ferunt characterem, quo primo obtutu distin-
gui possint; et aliquae species ita sunt consi-

miles colore et forma, ut hoc modo nata inter se differant.

18. *purpurea*, imbricata coriaceo-tremellosa tomentosa pallida: fasciis depressis fuscis, subtus purpurascens. *Syn. fung. p. 571.*

Auricularia amethystea. *Bull. champ. t. 483. f. 1.*

Auricularia faginea. *Schum. Saell. p. 399.*

- β . *lilacina*, subimbricata mollis a zona pallida, margine subtusque dilute purpurea.

Elvela lilacina. *Batsch Elench. fung. f. 131.*

- γ . *orbicularis*, resupinata simplex orbicularis, carnea, zonis concentricis.

Cresc. sero praesertim autumnio ad *Populi Tremulae*, *Betulae albae*, etiam *Alni vulgaris*, ubi magis effusa et coriacea apparet, truncos; β . ad quercus provenit, et γ . ad terram inveni particula lignea tamen adnatam. Pro diversis iis locis natalibus, colore plus minusve obscuriore variat. Hymenium laeve, in fungo exsiccato obscure fuscum; etiam cinereo-caerulescens *vorticosa*, imbricata coriacea zonata tomentosa vel strigoso-hirsuta, subtus purpurea. *Fries Obs. Mycol. 2. p. 275. +*

Auricularia persistens. *Sowerb. t. 388. f. 1.*
Auric. lilacina. *Schum. p. 398. (ex Fries).*

In Suecia ad truncos *Betulae*. Non nisi substantia ab antecedente distingui potest. An constans?

19. *hepatica*, pileo effuso reflexo imbricato glabro azono utrinque fusco - purpureo. — *Fries l. c. p. 276.*

Auricularia laevis. *Sowerb. t. 388. f. 2.*

Hab. in Anglia. Magnitudine communi; sed pileo ubique glabro et unicolore facile dignoscenda.

20. *bicolor*, imbricata fasciata subglabra spadicea subtus albe. *Syn. fung. p. 568.*

Thelephora fusca. *Schrad. Spicil. p. 187.*

Rarius sese offert, tamen bene distincta, consistentia tenuis, subflaccida, unc. 1-2 alta et totidem fere lata. Pileus parum tomentosus, arcte imbricatus.

21. *crispa*, effuso-reflexa, pileo undulato zonis crispo spadiceo, subtus cinereo pubescente. *Syn. fung. l. c.*

In Hercynia ad Pini Abietis truncos olim inveni hanc speciem parum ambiguam, quae substantia dura sicca, forte vetustate, distinguitur. Pileus basi sat late effusus, fasciis approximatis quasi crispus, fere glaber, sed hymenium, sub lente observatum, subvillosum apparet.

†† *Pileo resupinato seu lato effuso, margine libero demum in nonnullis parum subreflexo.*

22. *abietina*, coriaceo-membranacea, e fusco-cinerea subpurpurascens, hymenio umbrina pubescente. *Syn. fung. p. 176.*

Stereum abietinum, resupinatum subliberum, superne fusco-purpurascens, inferne umbrinum. *Fries Obs. Mycol.* 2. p. 274.

Hab. in Hercynia et Suecia ad ligna fabrefacta et ad truncos Pini *Abietis*, sed rare. Subtenuis, fragilis, 2-3 unc. latus, Hymenium subtomentosum quidem est, sed non setulis sparsis gaudet.

23. *Piceae*, resupinata inaequalis depressa glabra fusco-cinerea, papillis majusculis difformibus subrugosis.

In Vogesia ad ramos Pini *Piceae* invenit D. *Mougeot*.

Fungus junior suborbicularis, applanatus, margine parum pubescens; adultus irregularis, passim interruptus, margine liber, sed non reflexus, $\frac{1}{2}$ -1 unc. latus; colore et cinereo subpurpurascens. Papillae sat copiosae, in fungo juniore majores, sed tunc parcae.

24. *Pini*, tenuis glabra, globuloso-squamulosa subnitida spadiceo-cinerea nigricans.

Thelephora Pini. *Schleicher et Decand. Flor. Gall. Supplem.* vol. 6. p. 31.

Crescit in Helvetia et Jurae montibus ad cortices Pini sylvestris, pluribus individuis aggregatis: singulum vix 3 lin. excedit, et papillae vix distinctae sunt. Ab antecedente omnino distinctus, sed an *Thel. abietina* β . *pinæ* Albert. et Schwein. *Conspect.* p. 276. huc spec-

tes, in dubio est, cum auctores hymenium dicunt sordide album aut pallens.

25. *rufo-marginata*, majuscula suborbicularis tota glabra rugulosa cinerea, margine rufescente, papillis majusculis confertis laevibus.

Auricularia cinerea. Sowerb. fung. t. 388. f. 3.

Hab. in Anglia et in principatu Neocomensi, in Helvetia, ad ramos Tiliaceos.

Plantula satis pulchra, forma regulari distincta, unc. 1 fere lata, fusco-cinerea subnitida margine fuscescente. Substantia subrigida.

26. *quercina*, resupinata coriaceo-membranacea longitudinaliter effusa carnea, margine demum involuto subrusque umbrino. Syn. fung. p. 573. Nees Syst. der Schwämme t. 34. f. 233.

Thelephora corticalis. Decand. Syn. pl. Gall. p. 21.

Auricularia corticalis. Bull. champ. 1. p. 285. t. 436. f. 1.

Autumno vulgaris ad ramulos quercinos dejectas sed in prona parte seu quae terram respicit. Pileus ab initio totus depressus, sed demum margine involvitur. Humectatus subtrementellosus et rugosus fit. Recens laete carneus, vetustus fere nigricat. Nonnunquam spithamam longa, sed pro ratione angusta quod a latitudine ramulorum dependet.

27. *Cerasi*, parva suborbicularis flaccida (inferne) subtomentoso ferruginea, hymenio griseo - cervino setuloso.

Rarius provenit prope Nauchatel ad cortices Pruni *Cerasi*. Parvula, vix $\frac{1}{2}$ unc. lata (quoad specimen a cl. *Chaillet* transmissum); sed bene distincta species est.

* *Pruni*, coriaceo-membranacea effusa, primo subaurantiaca, demum umbrino-ferruginea rugulosa, margine byssino pallidiore. *Schumann*. *Saelland*, 2. p. 396.

In ramulis dejectis subputridis Pruni *spinosa*, Decembr. Pil. 1 - 2 poll. latus. *Schum.* — An a priore diversa et an hujus subdivisionis sit, ignoro.

28. *juratensis*, magna crassa cervina nitida, pileo tomentoso spadiceo. †

Hab. autumnno ad truncos Fagi *sylvaticae*, in summitate montis *Jurae*.

Obs. Specimen ab aestumatissimo D. *Chaillet* mecum communicatum, nondum satis evolutum erat, et verosimile sub maturitatem pileus reflexus est. Hoc hinc conjicio, quod hymenium, in specimine, fere absque papillis est, et fungi margo valde incrassatus coloreque dilutior est.

29. *Chailletii*, late effusa, tota glabra subcervina nitens, margine repando demum libero, papillis copiosis variis.

Species satis pulchra ad truncos *Taxi* nascitur prope Neuchatel. Subcrassa ferrugineo-nitida, 3 unc. longa, $1\frac{1}{2}$ lata.

30. *corylea*, late effusa ochracea crassiuscula, inferne glabra subspadicea, margine breviter reflexo. *Obs. Mycol.* 1. p. 35. *Syn. fung.* p. 569.

Circa Gottingam ad truncos *Coryli Avel-lanae* observata. Hymenium dilute ochraceum, saepe rimosum, papillis sparsis minutis obsitum. Fungus ceterum 3 - 4 unc. latus, et ad dimidiam unciam margine demum inflectitur. An duae sequentes tantum subvarietates sunt?

31. *peltata*, latissima lutescente-ochracea subrugulosa, margine demum lacera, inferne umbrina subtomentosa striis radiata.

Crescit prope Dax, in Gallia. *Thore.*

Truncis peltae in formam, uti *Gyrophorae*, affixa, spithamam fere longa. Papillae vix ullae.

32. *rudis*, latissima subcrassa pallida ochracea, superficie inaequali rudi, inferne strigoso-tomentosa, papillis (?) difformibus variis.

Hab. prope Pontuy, in Gallia. Ramos obducit ad latitudinem palmarem, 2 lin. est crassa, dura et forma quadam ingrata, quasi bullata distinguitur.

33. *pachyderma*, latissima applanata laevis pruinosa badia, subtus subvelutina spadicea.

Hab. ad truncos quercinos? prope Neocodium. Latitudine palmari, subregularis, ad latus albentior, margine crassiusculo subinflexo. Inferne obscura.

34. *rugosa*, effuso-reflexa, pileo duro flexuoso, glabro sordide fusco, hymenio rugoso laeviusculo, cinereo-pallido. *Syn. fung. p. 569. Swartz in Nov. Act. Suec. an. 1811. p. 77. †*

Ad truncos praesertim quercinos crescit. Pilei solitarii, effusi, marginibus reflexis, undulatis. Substantia fuliginea, rigidissima. Hymenium vel laeve, vel rugosum, demum fissile, papillis raris majusculis. *Siv.*

35. *discoidea*, crassa glabra subdifformis albidopallens concava, margine erecto undulato, inferne subsericea concolore. — *Decand. Flor. Gall. Supplem. p. 31. Moug. et Nees Stirpes cryptog. Fasc. 6. †*

Hab. in Germania, Gallia et Helvetia ad truncos quercuum. Forma huic speciei peculiaris, hinc facile dignoscenda. Unc. 1 lata subcrassa et duriuscula, extus pallide rufescens, disco quasi farinacea et laevis, sed demum saepius rimosa.

- β. compacta*, crassa dura tota glabra, pallide rufescens, disco depressa subfarinacea, margine undulato, subtus glabra nigricante.

Crescit ad truncos. Th. *discoideae* quo-

dammodo similis, sed multo crassior et durior et habitu minus grata, colore sordide fuliginosa, latitudine unciali.

36. *sinuans*, subrotunda inaequalis crassa, disco tuberculoso et inaequaliter impresso glabro pallida sublutescente rimoso, margine undulato-sinuato, subius spadicea, fasciis approximatis transversim sulcato.

Nascitur praesertim ad trabes quercinas, ad pontes destinatas, quarum sinus investit. *Moug.*

Priori non valde dissimilis, attamen diversa. Unc. 1 fere lata, 3 - 4 lin. crassa, valde flexuosa, superne glabra, non farinacea; inferne pileus distinctus adest, qui plures fascias seu zonas habet, sed non reflectitur. Fungus maturus rimoso-partitus ut in *Th. frustulata* a quo vero forma et statu perfectiore distinguenda.

***** CORTICIUM. Resupinatum effusum, margine totum adnatum, nec liberum. Papillae plerumque perspicuae. (Colore, plerumque dilutiore, varium).

† Colore incarnato, rufescente aut alutaceo.

37. *alutacea*, late effusa suborbicularis, disco rugosa alutaceo-carnea, margine latiusculo laevi albido.

Late investit ipsos arborum cortices, locis humidis jacentes, margine quidem mollis est, sed

sed non byssoidea et papillis destituta, quae discum occupant et majusculae sunt, forma variae, alutaceo-pallidae.

β. zeolitica, late effusa (humida) carneo-fuscescens tota glabra, papillis congestis majusculis subglobosis.

Passim occurrit hyeme ad sepimenta lignea. Spithamam longa, 2 unc. lata, margine tenuis, subpellucida, substantia coriaceo-tremellosa.

γ. gilva, tota fere glabra carneo-livida, papillis variis submarginalibus.

Sero autumno et hyemi leni ad truncos putrescentes nascitur. Longitudine antecedentis. Superficies ejus hinc inde laevis, hinc versus marginem papillae sparsae sitae sunt.

38. *concentrica*, effusa crassiuscula alutacea, margine subalbescente, papillis majusculis subhemisphaericis concentricis. *Alb. et Schwein. Consp. fung. p. 279.*

Ad sepes pineas se propagat, autumno et vere. Ultra spithamam saepe longa, 3-4 unc. lata; dum viget humida est laevis et alutacea, exsiccata vero aut senilis subrufescit et dura rimosaque evadit; papillis tunc collapsis et evanidis. *Consp.*

39. *amphibolia*, effusa tenuis fuscescens subalutacea, papillis sparsis majusculis. *Fries Obs. Mycol. I. p. 151.*

In hymenio Boleti suberosi tantum, in Suecia, inventa. Brevis, inodora et arcte adnata, pallescens. *Fries.*

40. *odorata*, suaveolens suberosa membranacea glabra laeviuscula confluens pallida.

Ad truncos abiegnos destructos. Glaberri-
ma tota, crassiuscula ex albido alutacea vel sub-
rufescens, confluens 4-5 unc. tunc lata, papil-
lis rarissimis. Odor fere Boleti odorati, cum
quo affinitatem habet. *Fr.*

41. *radiosa*, effusa confluens laevis glabra alu-
taceo-pallida, margine albo fimbriata. *Fries*
l. c. 2. p. 277.

Ad ligna dejecta faginea, quibus arcte ad-
nata; confluens 2-3 unc. lata. Papillae nul-
lae. *Fr.*

42. *laevis*, ex albido carneo pallens sublongi-
tudinaliter effusa laevis, margine byssino.
Syn. fung. l. c.

Hab. praecipue ad ramos populneos. Satis
longa, et aliquando tuberculosa: papillis sub-
globosis, verum plerumque superficie unita est.

43. *incarnata*, coriacea subcrassa laevis incar-
nata situ longitudinali, margine subbyssino.
Syn. fung. p. 573.

Non equidem communis ad ramos reperi-
tur. Papillae vix distinctae et parvae. Margo
etiam glaber.

β? *Corium*, magna crassiuscula carneo-pallida, margine sublibero, papillis obsoletis. —

Syn. fung. p. 574 †

Specimen hujus varietatis aut speciei mihi non magis ad manus est, ut certi quid de ea adferri possim.

44. *rosea*, dilatata subcorrugata tenuis rosea, margine tomentosa. *Syn. fung. p. 575.*

Eligit pulchella haec species, truncos aut assulas praesertim *Betulae albae*, passim quoque *Populi Tremulae*. Fere epapillosa est. Magis lata, quam longa. Satis nitida species.

45. *maculaeformis*, adpressa pallide rosea, junior subrotunda, pseudopapilla unica, mox confluens, margine glaberrimo. *Fries Obs. Mycol. I. p. 150.* †

Ad cortices in Suecia. Pileus crustaeformis, tenuissimus, membranaceus, dilute roseus, suborbicularis, vix lin. latus, sed mox confluens et difformis. *Fries* (Fungus videtur junior).

46. *fallax*, glabra arcte adnata laevis, ex alutaceo subincarnata, pseudopapillis majusculis angulatis. *Syn. fung. p. 574.*

Provenit in ramis minoribus, praecipue quercinis.

Obs. Nonnunquam veras papillas proferre videtur, quae tunc nimis regulares sibi

consimiles, ita ut a nodulis ramorum non ortae videantur.

47. *salicina*, late effusa subcarnosa mollis, primo albida, dein incarnata, superficie bullato-tuberculata pruinosa.

Autumno passim in salicibus cavis putrescentibus, prope Parisios lecta. Longitudine fere palmari, satis crassa, interrupta tuberculosa, ut in *Malachite*; exsiccata, ob substantiam molliorem, fuscescit et diminuitur.

48. *polygonia*, incarnata glabra opaca, papillis majusculis variis: subrotundis, angulatis et disco excavato-depressis. — *Syn. fung. l.c.*

Hab. satis rara et minus cognita species ad ramos *Populi Tremulae*, in Germania et Helvetia.

Juniores suborbiculares et papilla unica instructae, margine subbyssinae, dein plures in crustam uniformam confluunt. Exsiccata quasi pruinosa est, sed non rimulosa.

Obs. *Thelephora polygonia* in *Conspectu fungorum Lusat. superioris* p. 276. enumerata longe alius fungus videtur ob "marginem demum undique revolutum, duriusculum, subtus nigricantem."

49. *castantae*, unicolor incarnata tota glabra expallens sicca demum rimosa, papillis sat copiosis regularibus.

Hab. ad sepes in cortice ramorum *Castaneae vescae*.

Junior orbicularis, margine subradiato arcuato adpressa. Rimae, exsiccatione ortae, praesertim circa papillas, lirellas imitantur. Affinitate quadam gaudere videtur cum *Th. quercina*, sed tenuis est, et cum cortice ubique arcuato adnata, nec margine demum revoluta, etiam irregulariter effusa.

50. *dichroa*, longitudinaliter effusa tota glabra tenuis albido-cinerea subnitida, papillis sparsis rugosis subdifformibus rufo-carneis.

In ramis junioribus *Hippocastani*?

Primo intuitu *Th. polygoniae* similis, sed colore et substantia nec non forma papillarum ab illa differt. Margo regularis, in junioribus orbicularis, nam plura individua demum in crustam elongatam confluunt. In statu vetusto etiam fissuris sat magnis distinguitur.

†† Colore pallido, ochraceo, flavescente et subaurantiaco.

51. *fissilis*, longitudinaliter effusa, sordide alutacea, demum in particulas basi cohaerentes fissa, papillis copiosis magnis subaequalibus. †

Provenit parum dubia haec species, ob affinitatem cum *Thel. laevi* et *Thel. incarnata*, in corticibus *Tiliae* et *Carpini*, sed minus frequens. Crassiuscula est et carnosa, superficie

in particulas plus minusve multas, formaque varias fissa. Margo vix byssinus.

52. *agglutinata*, tenuis latissima glaberrima unita sordide rufescens fuscescensque subnita pseudopapillis variis rugulosa.

Hab. rarius in ipsa epidermide truncorum quercuum, iisque ita adnata, ac si agglutinata esset, latitudine palmari. Papillae verae non adesse videntur.

53. *corrugata*, effusa arctissima adnata tenuis rimosissimo-corrugata lurida, papillis nullis. *Fries Obs. Mycol. I. p. 154.*

In ligno decorticato in Suecia. Irregulariter effusa, glabra, 2-3 unc. lata. *Fr.*

54. *frustulata*, tota glabra laevis applanata durissima rimoso-partita: particulis sejunctis basi nigricantibus. *Syn. fung. p. 577. Decand. Flor. Gall. Suppl. p.*

Hab. ad ipsa ligna sicca; vix unc. 1 magna, sed subcrassa, demum in partes subaequales dilabitur.

β. *tessulata*, crassissima, tessulata cinnamomea, pulvere concolore obtecta. *Fries Obs. Mycol. I. p. 153.*

In Suecia, ad lignum dejectum. An species propria?

55. *Porioides*, parvula sublinearis, rimoso-porosa sordide pallida subvirescens. †

Passim ad cortices nascitur. *Chaillet.*

Longitudinaliter effusa, et in areolas numerosas subforaminulosas demum undique fissas. Color obscurus et fungus denuo observandus.

56. *mucida*, late effusa tenuissima undique glabra et incrustans expallida, papillis minutis sparsis.

In truncis caesis cariosis, quorum fissuras arcte investit. Papillae minutae lignis ortum suum forte debent, nam non ubique in superficie fungi adsunt. Color pallescente-subincarnatus, languidus.

57. *sebacea*, late et subaequale effusa incrustans laevis submollis pallida rufescens, papillis nullis. *Syn. fung. p. 577.*

Thelephora incrustans. *Obs. Mycol. I. p.*

39. *Syn. fung. l. c.*

Hab. aestatis fine post pluvias ad terram, supra gramina, folia sicca etc. Sebi ad instar effusa tunc pallida, tandem subrufa. Primo albidula, margine hinc illinc laciniata.

Obs. Haec et sequens species formam quidem habent *Thelephorarum*, sed crescendi modo et natura ad *Merismata* accedunt. *Th. incrustans* fungum adultum sistit.

58. *ericetorum*, compacta incrustans fulvo-grisea pulverulenta subtenuis. †

Semel eam inveni ad radices exsiccatas *Ericae cinercae*, commixta cum terra nigra *ericacea* (*terre de Bruyères*). Non equidem late est effusa. Color chrysophaeus.

59. *epidermea*, tenuis late effusa glabra prima alba dein pallescens, margine subbyssino, papillis (?) sparsis subangulatis.

β? *helvola*. *Hypochnus helvolus*; effusus tenuis; junior lacteus byssinus dein subglabratus *helvolus*, margine fimbriato. *Fries Obs. Mycol. 2. p. 281.*

In truncis dejectis *Betulae*. Margo latus albus. Papillae plane nullae. *Fr.*

Sero autumnno in sylvis ad corticem nudum ramorum quercuum; nec non *Betulae albae* observatur, 3-4 unc. magna.

60. *confluens*, subrotunda arcte adnata laevissima contigua glabra (alba aut laetissime flava), margine glabro adnato. *Fries l. c. p. 152. †*

Crescit in Suecia ad ramos *betulinos*. Vix uncialis, orbicularis, ramis arctissime adhaeret.

61. *citrina*, parce effusa laete flava ubique glabra laevis, papillis sparsis minutis.

Inventa prope Neuchatel ad truncos *abietinos*. Ob colorem citrinum pulchra species. In superficie (fungi transmissi) etiam tubercula majuscula sparsa observantur, sed hoc, uti vi

detur, ab subjacente ligno ejusque asperitatibus dependet; papillae verae parce adsunt et sat parvae.

62. *ochracea*, effusa latissima subtenuis ochracea, papillis sparsis irregularibus majusculis.

Fries l. c. p. 151.

In ligno abiegno putrido. Latitudine et longitudine pedali. Margo nunc glaber, nunc villosus. Papillae ab asperitatibus ligni ortae videntur. *Fr.* — *Thel. coryleae* affinis?

63. *decorticans*, oblonga laevissima tenuis glabra subcorticalis arcte adnata lutescens, demum fuscescens opaca.

Thelephora carnosa. Ehrenb. Sylv. Mycol. p. 30.

Thelephora comedens, e cortice erumpens, irregularis subcoriacea, lutescente - carnea, in ambitu albo-floccosa (?), hymenio laevi. *Nees System, 2. p. 65. t. 34. f. 255.* (icon. non adaequat).

Ubique fere ad ramulos exsiccatos quercinos, quorum cortices sejungit et destruit. Ultra spithamam nonnunquam longa, sed ob matricem vix uncia lata; et ita tenuis, laevis et concolor, ut ab ipso ligno oculo fugitivo, distinguere nequeat. Omnium minus carnosa est.

64. *Hydnoidea*, glaberrima decorticans flavo-aurantiaca, papillis sparsis elongatis subuli-

formibus. *Syn. fung. p. 576. Decand. Flor. Gall. vol. 2. p. 34.*

Nascitur quoque sub epidermide, dein soluta, ramorum Quercuum et Fagorum, autumnos sat vulgaris.

Obs. Papillae praesertim quae in medio sitae sunt, elongantur, forsitan ut epidermidem depellant, et dentes *Sistotrematum* omnino referunt, verum eae quae magis marginem fungi occupant, forma sunt papillosa, ceterum inter se distant, nec situ determinata neque numerosae, ut in *subulis Hydnorum* hoc obtinet.

65. *aurantia*, elongata nuda glabra aut margine subbyssina, aurantiaca, papillis varioribus. — *Syn. fung. p. 576.*

β. rimosa, elongata angusta aurantio - carnea, transversim rimosa.

Hab. ad ligna et ramulos decorticatos, praesertim hyeme, *β.* in horto botanico Parisiensi ad sepes. Caute distinguenda a *Thel. carnea*.

Nunc unc. 1 fere lata, nunc vix 3 lin. lata, etiam ubique glabra, aut margine subtomentoso distinguitur; etiam crassitudine variat. Color demum expallescit et tunc *Th. incarnatae* similis evadit haec species.

*** *Colore lateritio, croceo, cruento.*

66. *bolaris*, submagna orbicularis confluens cro-

ceo-lateritia (?), margine fibroso-sericea, papillis nullis?

Auricularia aurantiaca. *Sowerby Fung. 3. t. 291.*

Hab. in Anglia supra truncorum epidermidem.

Species elegans, formaque regulari distincta. Color quem autor aurantiacum pronunciat, secundum iconem potius lateritia aut bolaris apparet, nec flavedine tinctus.

67. *lateritia*, erumpens interrupte effusa, subconfluens dilute lateritia, papillis? densis rugosa. *Chaillet.*

Vere ad cortices quercinos, sub quibus se propagat, dein transversim erumpit, superficie costata, colore lateritio-alutacea. Habitu a ceteris speciebus differt hic fungus, nondum uti videtur plane adultus.

68. *crocea*, tenuis fragilis tota crocea. *Gmel. Syst. Nat. Lin. 2. p. 1441. no. 18. Schrad. Spicil. Flor. Germ. 138.* (Exclusis synonymis). †

Hab. in fissuris quercuum annosarum, aut ad Fagos emortuas. *Schrad.*

Obs. Specimen olim ab auctore datum a *Xylostromate giganteo* Tod. differt, nam substantia est crassum, compacto-pulveraceum, forte vetustate, ut vix hujus generis videatur.

69. *leproides*, indurata cohaerens opaca fusco-lateritia subaspera. †

Ad cortices exsiccatos, unica vice a me reperta. Primo intuitu, Leprarium aut Isidium refert, sed substantia, quamvis compacta, est coriacea et margine substriata. Forma irregulari, latitudine fere biunciali.

70. *cruenta* suborbicularis crassiuscula coriacea glabra rugoso-tuberculata sanguineo-rubra. *Syn. fung. p. 575. Decand. Synops. plant. Gall. p. 21. Mougeot et Nest. Stirp. cryptog. Vogeso-Rhenanae, no. 581.*

Hab. in subalpinis ad cortices Pini *Piceae*; aut ad ramos emortuos *Salicis Capreae* (Alb. et Schwein.)

β. roseo-rubra. Conspect. fung. p. 277.

Crescit ad ligna ramosque putridos.

"Var. *α. orbicularis* est, circumscripta, mediocris, vix uncialis, utrinque glabra, pulcherrime sanguinea, papillis instructa satis magnis subangulatis; tempore pluvioso, molliuscula, quasi ceracea adpressa, — sicco, firmula, subfragilis, semilibera: margine introflexo. — Var. *β. ad 2-3 unc. effusa*, laevis, subpruinata, constanter appressa, membranaceo-coriacea; *hymenium* dilutius roseum; papillae saturationes cruentae, congestae, subhemisphaericae." *Albertini et Schweinitz. Conspect. p. 278.*

71. *punicea*, subrotunda oblongaque tomentosa (pubera) subpulverulenta sordide punicea, demum brunescens. *Conspectus fung.* p. 278.

Prov. ad cortices fagineos et abietinos in Germania. Etiam in monte *Jura*. (Chaillet.)

"Colore ludit cinnabarino, sublateritio. Margine appressa vel subreflexa, subtus item tomentosa, colore terreo distincta. Papillae minutae, granulaeformes." *Alb. et Schw.*

Obs. Antecedenti ita similis, ut pubescentia tenui, sub lente conspicienda, ab illa distinguui solummodo possit.

†††† Colore ferrugineo, aut ferrugineo-grisea, flavo-fusco, fusco-purpurascente aut badio.

72. *ferruginea*, suborbicularis ubique tomentosa, medio pulveraceo-papillosa. *Syn. fung.* p. 578. *Obs. Mycol.* 2. p. 18.

Thelephora Persoonii. Decand. Flor. Gall. 2. p. 107.

Hab. in fissuris truncorum aut sub cortice ramorum. Late effusa, 3 unc. fere magna. Colore et substantia villosa, uti et forma *Hydno ferrugineo* simillima.

73. *cinnamomea*, late effusa tomentosa subcompacta, superficie piloso-setacea.

Prope Neuchatel inventa est haec species ad ligna quercina. In se conjungit caracte-

res genericos *Racodii*. *Herei* et *Hypochni* Fries *).

74. *aurea*, late effusa tomentosa aureo-fulva, superne granulis subvillosis vestita.

Hypochnus aureus. Fries *Obs. Mycol.* 2. p. 280. t. 6. f. 4. Micheli *Nov. Gen.* p. 124.

Hab. in Italia et Suecia in cortice dejecto *Betulae*, cui arcte adnata. Subtus glabra, laevis et subfusca. *Isabellina*, late effusa tomentosa isabellina (gilva aut rutilans), pseudopapillis regularibus. Fries, l. c.

Cresc. in cortice dejecto *Fagi sylvaticae*. Oriuntur papillae sparsae ex nodulis corticis aut sphaerulis. Fr.

75. *Padi*, griseo-ferruginea subfarinacea, pilis brevissimis obsita, fere epapillata, plicis oblongis subanastomosantibus.

In Vogesia ad ramos exsiccatos *Pruni Padi*. Substantia sicca, tenuis, non tomentosa. Color subcervinus, opacus. An plicae constantes et huic fungo naturales?

76. *fusco-purpurea*, tenuis glabra subcompacta purpurascente-fusca, undique setulis obsita.

*) *HYPOCHNUS*. Pileus pagina superiore adnatus, et in nonnullis exoticis reflexus horizontalis, submembranaceus, totus fere tomento contextus, vel pulvere in globulos villosiusculos compositus. Fries *Obs. Mycol.* 2. p. 278.

Rarissime ad ligna exsiccata crescit, longitudinaliter effusa, $1\frac{1}{2}$ unc. magna, tenuissima, vix supra lignum eminet.

77. *laxa*, orbicularis parva subconfluens mollis, subtus villosa alba, disco laevissima fusco-carnea.

Hab. in summitatibus montium Vogesiorum.

Subtenuis, vix 3 lin. lata, flaccida, subrimosa. Forte nondum bene evoluta. Affinitatem habere videtur cum *Peziza amorphae*.

78. *olivacea*, pulverulento-tomentosa olivacea, margine fimbriato albido. *Fries l. c. p. 282.* sub *Hypochno*.

In lignis abiegnis indeterminate effusa.

Obs. An cum tribus sequentibus *Coniophorae* species?

79. *marginata*, orbicularis subeffusa pruinato-tomentosa ex aurantio vel subrubente cinerascens, papillis majusculis crassis coacervatis, margine reticulato intricato discolore olivaceo. *Conspect. fung. p. 281. et 284.*

"*Corticium elegans*, fere biunciale, medio crassum, carnosum succulentum, margine siccum fibrosum; fibrae crassiusculae, ramosae, olivaceae, in reticulum inaequale intricatae. Species rarissima, in cellis, lignis pineis putrescentibus increscens, inventa."

80. *puteana*, suborbicularis carnosogelatino-sa olivascens, papillis ovalibus obscurioribus, maculisque umbrinis ex punctis elevatis, margine tenuissimo byssino. *Schum. Saell. 2. p. 397.*

In puteo quodam ad asseres reperta. Quinque pollices lata, olivaceo-virens, margine candidas, interne fusco-ochracea. — Affinitatem gaudere videtur cum *Comophora foetida*.

81. *Betulae*, membranacea rugoso-tuberculosa umbrino-badia, flavo-umbrino maculata, margine flexuoso luteo, papillis inaequalibus e fuligineo subnigricantibus. *Schum. Flor. Saell. 2. p. 396.*

In cortice *Betulae albae*, in Dania. Octobr. et Novembr. Duas uncias fere lata.

82. *chalybea*, tomentosa chalibea subcinerascens, papillis congestis pulverulentis. *Syn. fung. p. 579.*

Ad truncos provenit rarius, ubi ad 3-4 unc. effusa, textura ut in *Thel. ferruginea*.

Obs. Species sub hoc nomine in *Conspect. fung. Lusat. p. 281.* descripta, differt: "margine passim albido, etiam amare dicitur, vel ligna pinea fabrefacta *cellaria* tempore hyemali, vel ligna rudia sylvatica, aestate."

83. *bufonia*, subtomentosa e cinereo brunnea subpulverulenta, papillis congestis majusculis. *Syn. fung. p. 578.*

Semel tantum inventa, unc. 1. fere magna, suborbicularis, intus tomentosa.

84. *pusilla*, orbicularis parvula tomentosa obscure spadicea, demum confluens, papillis parvis (saepe unica) magnis solidis (non pulveraceis). †

Ad ramos siccos, in Helvetia, rarius, 3-4 lin. tantum lata. Subiculum totum tomentosum, margine (sub lente) parum fibratum. Papillae pro magnitudine fungi (forsan adhuc junioris) magnae, glabrae.

85. *monothelia*, subrufo-grisea glabra subfarinacea interrupta, papilla unica magna globosa aequali. †

Ad truncos rarissime. Fungus nondum evolutus videtur. Ad minores pertinet.

86. *fraxinea*, inaequaliter effusa fusco-cinereascens subtomentosa, sed compacta, papillis angulatis.

Thelephora cinerea. β . interrupta. *Syn. fung.*

Hab. rarius ad ramos exsiccatos *Fraxini excelsioris*. Subiculum interrupte effusum, cortici arcte adpressum, unciale, colore cinereo-spadiceum. Papillae subumbilicatae.

***** *Colore nigrescente.*

87. *fuliginosa*, magna coriacea fuliginosa seu nigrescente-spadicea, setulis densis obsita.

Hab. in Gallia ad ligna cariosa, mollia.

E longinquo *Dematium herbarum* quodammodo similis, sed revera hujus generis, ex membrana contigua coriacea est, licet subtenuis, et setulis copiosis, uti hoc in multis *Thelephoris* locum habet, instructum, papillis distinctis vero destituta. Ad 3-4 uncias effusa, $1\frac{1}{2}$ unc. lata.

88. *pithya*, nigricans interrupte effusa tota appressa glabra tenuis, demum rimosa, papillis subconfluentibus.

Hab. in Helvetia ad ligna fabrefacta *Pini sylvestris*? Crustam sistit late diffusam atram, subnitidam, matrici toti adnatam.

* *faginea*, glabra nigrescens, siccitate cinerea.

Schrad. Spicileg. Flor. germ. p. 186. Thelephora nigrescens.

Ad ramos dejectos et cortice destitutos *Fagi sylvaticae*, in Germania inventa. Membrana est tenuissima appressa, per aetatem in areolas polygonias dehiscens. *Schrad.*

89. *obscura*, tenuis irregularis subglebulosa nigro-fusca, papillis nullis. †

Supra ipsam epidermidem corticis arborum emortuarum, *Betulae* etc. inventa a D. *Chaillet*. Leprariam fuliginosam non male refert. Interrupte effusa, rimoso-glebulosa, opaca, glabra.

***** Colore caeruleo, caesio, cinereo vel griseo.

90. *caerulea*, longitudinaliter effusa subtomentosa amoene caerulea, margine demum sublibero, papillis parvis subacutis.

Thelephora caerulea. Decand. Flor. Gall. 2. p. 107. Th. fimbriata. Roth. Catal. bot. *Auricularia phosphorea*. Sowerb. t. 383.

Hab. pulchra haec species, hyemi leni, ad ligna et sepimenta subputrida, supra quae ad longitudinem spithamaeam serpit.

Rarius bene explicata et nonnunquam ut cespitulum villosum (an tunc *Byssus phosphorea* Lin.?) sese offert; perfecte vero evolutus expansus et margine, qui albicat, ad 2-3 lin. inflectitur, ubique vero tomento tenui oblecta, colore caesio-cinerea opaca, exsiccata rimosa et in aetate ulteriore grisea, subfuliginea evadit; hinc opinor sequentem fungum non specie distinctum.

* *fumosa*, effuso-diformis caeruleo-cinerea, villosa. Fries Obs. Mycol. 2. p. 279.

Cresc. supra folia dejecta *Betulae* in Sueciae sylvis montosis.

91. *Tiliae*, sublata effusa grisea, margine subtomentoso adpresso, papillis inaequalibus.

Hab. in Vogesia ad *Tiliae* ramos putrescentes. Forma est regularis, laevis, 2 unc. lata, fuscescenscineria, junior magis cinerea.

92. *cinerea*, lata adnata glabra cinerea, papillis parvis. *Syn. fung. p. 579. Conspect. fung. p. 282.*

Provenit plerumque ad ramos cortice denutatos, quos fere totus ambit, laevis, margine glaber.

93. *Lycii*, suborbicularis crassiuscula confluens glabra, subnitida, papillis minutis subcongestis.

Ad ramos exsiccatos *Lycii Barbari*. Singulum individuum $\frac{1}{2}$ unc. latum, crassiusculum, subrugosum, margine subrepandum, albidocinereascens, subpulverulenta, intus fuscescens.

94. *glebulosa*, glebuloso-partita glabra subparva cinereo-nitida: detrita intus fuscescens.

- β. *virgata*, parallelo-lineata conferta pulverulenta caesio-cinerea, papillis majusculis.

In hortis ad sepes oritur. Ab antecedente distinctam puto. Formam habet valde inconstantem: nunc enim linearis est et continua, nunc glebulosa, subrotunda (junior), applanata, interrupta aut confluens, demum rimosa, margine exciso.

95. *livida*, tota glabra subviscosa e caesio rufescens, papillis variis, nonnullis dilutioribus. *Obs. Mycol. I. p. 38. sub Corticio.*

Th. livida latissime effusa glaberrima subhumida livida caerulescensque, papillis sparsis irregularibus. *Fries Obs. Mycol.* 2. p. 276.

β. viscosa, late effusa subgelatinosa viscosa dilute caesia, papillis majusculis sparsis. *Syn. fung.* p. 580.

Hab. sero autumnno ad truncos et ligna putrida, quibus arcte adnata, margine glabro; exsiccata praesertim *β.* valde diminuitur et sordide fuscescit.

96. *viscosa*, late effusa gelatinosa subgyrosa tuberculosa caesio-cinerea, margine byssino albido, papillis minutis concoloribus. *Schumm. Saell.* 2. p. 397.

In trunco subputrido *Ulmi campestris* in Dania ab auctore observata, 3-6 unc. lata.

97. *grisea*, late effusa tota glabra mollis membraceacea (albido-caesia).

Reperi hunc fungum in Frigidario quodam ad ligna pinea fabrefacta subhumido crescentem. Spithamam longa, 2 unc. fere lata, 1 lin. crassa, coriaceo-tremellosa, sparse tuberculosa, quoad colorem sebo liquefacto similis, exsiccata vero sublutescit.

98. *cyclothelis*, magna membranacea, sicca rigida, undulato-rugosa cinerea, papillis regularibus orbiculatis aut hemisphaericis obscurioribus.

Obs. Singularem hanc speciem, generis forsitan proprii ad juglandes observata a cl. *Chaillet*. — Gyrophoram quamdam refert. Papillae subimmersae.

***** Colore toto albo.

99. *pergamenea*, magna membranacea subrigida laevissima glabra albido-pallens; recens glauca.

Hab. ad truncos pineos in Vogesia et Helvetia. Spithamam sublonga, 2 unc. lata. Substantia tenuis subpergamenea praecipue exsiccando. Papillae rarissimae. Margo liber, subinvolutus.

100. *gigantea*, maxima difformis sublibera glabra lactea. *Fries Obs, Mycol. 1. p. 152.*

Supra Pini *sylvestris* corticem in Suecia. Species dicit hic auctor, maxime singularis et alii forte generis. Membrana est 6-9 unc. longa et lata, valde difformis, truncis leviter adhaerens, contigua, flexilis, undulata, sine papillarum vestigio.

101. *cretacea*, lata submollis alba simularis, margine byssino, papillis confertis parvis rugulosa.

In frigidariis ad tabulas abietinas saepius humectatas; durante hyeme, gignitur. Saepe late effusa, subtenuis. Color in fungo recenti lacteus, in exsiccato pallidus. Margo byssinus brevis.

102. *leucocoma*, glabra laevissima albedo-pallens, margine angusto, fusco-nigro.

Ad ramos *Castaneae vescae* sepibus inser-vientes in Horto botanico Parisiensi inveni sat pulchram hanc speciem, margine nigricante subsinuatum facile distinguendam. 3 unc. longa.

103. *lactea*, suborbicularis ex albedo livida, medio subcarnosa: margine byssino. *Syn. fung. p. 581.*

Ad cortices *Pini Abietis* nascitur. Papillis vix ullis distincta.

104. *sera*, nivea effusa pruinato-subtomentosa tuberculosa mollis. *Syn. fung. p. 580. Consp. fung. p. 283.*

Occupat ligna putrescentia in sylvaticis udis, 2 unc. fere lata et longa. Octob. — Novembr.

105. *cariosa*, tenuis laevis alba, margine lato tomentoso.

Eam in interiore *Salicis albae* (excavatae) observavi, ligno carioso ubique adhaerentem; 2 unc. magna est. Ab initio nivea, dein medio pallescit.

β? *exigua*, candida parva in particulas divisa margine villosa, disco glabro pulverulento. †

Sero autumno ad truncos putrescentes passim occurrit. Singula particula 2-3 lin. lata, irregularis.

106. *Sambuci*, candida rugulosa pruinosa, margine glabra, papillis obsoletis. *Disp. Meth. fung. p. 31.*

Thelephora calcea var. γ . *Syn. fung. p. 581.*

Thelephora cretacea. *Fries Obs. Mycol. I. p. 153.*

Frequens aestate et autumno post pluvias ad truncos cariosos *Sambuci nigrae*. Pulverulenta, subinquinans (var. *Samb. racemosae* et glaberrima, unita), saepe rugoso-tuberculosa, sed non equidem papillas profert; interea non ut initium alterius fungi consideranda.

107. *acerina*, gregaria parvula subtenuis inaequalis glabra albido-caesio, papillis distinctis nullis. *Syn. fung. p. 581. Obs. Mycol. 2. p. 37.*

Communis ad truncos *Aceris Platanoidis* et *campestris*. Subnitida, argentea demum confluens.

108. *dryina*, crustaceo-membranacea rugulosa albido-pallens subpulverulenta.

Th. acerina. β . *quercina*. *Syn. fung. p. 182.*

Hab. ad cortices vetustos quercuum, quorum fissuras investit, forma hinc irregularis. Papillae distinctae nullae adsunt. Majorem affinitatem cum Th. *Sambuci* habet, sed multo tenuior est, nec ita pruinosa et farina detergibili oblecta.

109. *calcea*, gregaria parvula inaequalis confluens glabra albedo-fuscescens laevis. *Syn. fung. l. c.* †

Ad ligna dura exsiccata rarius crescit, ultra 2 unc. longa, $\frac{1}{2}$ unc. lata, compacta.

110. *aspera*, globulosa candida, setulis divergentibus subrigidis aspera. *Tab. VI. fig. 5.*

Semel sese mihi obtulit hic fungus generis dubii, nam non vera *Thelephora* videtur, nec bene ad *Hydna* referri potest. Aliud et curatius requirit examen, et hunc in finem icon est addita.

XLVI. CONIOPHORA. Resupinata, carnosomembranacea, mollis, rugoso-tuberculosa, pulvere manifesto oblecta.

1. *membranacea*, latissima subtenuis membranacea zonata, margine byssino pallido, disco badio. — *Decand. Flor. Gall. 6. p. 34.*

2. *leioplaca*, suborbicularis laevissima pallide fuscescens zonata: zonis concentricis pulveraceis.

Auricularia pulverulenta. Sow. t. 214.

Crescit passim in vaporariis ad ligna, vel in locis humidis ad trabes putrescentes; β . ad parietes humori expositos, toto fere anno, sed magis sero autumno.

Latitudine nonnunquam insigni, et e matrice facile evellenda. Superficie tuberculis difformibus magnis saepe notata, et sporulis globosis spadiceis oblecta, quae in β . magis in zonis coacervatae sunt; sed nullis venis, nec plicis distincta est haec species, quo differt a *Merulio destruente*, cum quo naturam participat septicam.

2. *foetida*, carnosio-tremellosa rugulosa fusca albo-pruinosa: zonis dilutioribus, margine lato albido, verrucis inaequalibus magnis.

Thelephora foetida. Ehrenb. Sylv. Mycol. p. 18. et 30.

Hyeme, prope Berolinum, in asseribus cloacam tegentibus inventa; late effusa, margine appressa. Sporidia minima, oblonga. Ehr.

3. *cuticularis*, subparva tenuis azona albido-pallida, margine byssino fibrilloso, pulvere fusco-olivaceo. †

Ad ligna exsiccata pinea sub ponte quodam inventa. Vix (in specimine a D. Chaillet transmissio) 2-3. unc. lata, ligno arcte

adhaeret; aetate juniore laevis glaberrima, de-
in pulvere subcompacto obscura obducta.

4. *cerebella*, undulato-gyrosa primo albicans
mollis pulvere rubiginoso. *Syn. fung. p.*
580. sub. *Thelephora*.

Hab. in Germania et Gallia in cellis ad
ligna. Substantia et superficie inaequali a prio-
ribus valde differt, hujus tamen generis est.

Obs. Nonnullae *Thelephorae*, praesertim
eae quae sub nn. 78 - 80. et 104. enumeratae sunt,
et a me ipso nondum visae, hujus quoque ge-
neris esse, videntur.

XLVII. MERISMA. Coriaceum digitato-
aut foliaceo-divisum ubique fructificans, lae-
ve, apicibus ut plurimum villosum. (Forma ut-
plurimum *clavariae* ramosae).

* *Decumbentia*, aut *adscendentia*, *subdifformia*.

1. *fastidiosum*, candidum late incrustans foe-
tidum e ramis laminosis constans. *Syn.*
fung. p. 582. *Hall. Stirp. Helv. ed. 2.*
no. 2251.

In Germania et Helvetia in umbrosis ad
terram magno saepe strato et cum vicinis cor-
poribus concrevit, praesertim post pluvias
autumnales.

2. *vermiculare*, decumbens albicans ramosis-
simum: ramis teretibus attenuatis rugulosis
subcarnosis.

Hab. prope Parisios, sed rare in sylvis, gramina et folia sicca subjacentia incrustans et ambiens, forma stalactidea. Rami sat longi sed tenues, vix 2 lin. crassi, ab initio subgelatinosi, dein subcartilaginei. Muscos allicit.

3. *serratum*, subascendens caespitosum viscosum fuligineo-albidum ramosissimum: ramis latiusculis serratis obtusis. *Comment. de Fung. clavaef. p. 106. t. 4. f. 7.*

Rarius in sylvis. Subcarnosum totum glabrum, affini *Thelephorae sebaceae*.

4. *cristatum*, adscendens subcoriaceum pallidum, ramis laciniatis subdifformibus membranaceis aut incrassatis rugosis. *Syn. fung. p. 583.*

β. *tuberculosum*, subdecumbens pallidum; ramis incrassatis obtusis. *Commentatio, p. 228. t. 2. f. 1.*

Clavaria laciniata. Bull. Champ. t. 415. f. 1. Sowerby t. 158.

Haec species apparet jam tempestive post pluvias aestivales in sylvis caesis, inter muscos. Primo fere albida, dein pallescit, substantia subramosa, sat pulchre divisa in ramulos stalactideos cristatos aut tuberculosos glabros.

γ? *penicillatum*, albedo-pallens basi incrustans, in ramos parcos obconicos multifidum. *Syn. fung. p. 583.*

In fagetis ad terram quoque crescit, sed postea non observare potui, an revera ut species differat. Rami magis incisi, quam illi *M. cristati* sunt.

Obs. *Merisma cristatum* β . *fuscum* Consp. fung. p. 284. *fusco-purpurascens*: apicibus subhirsutis albidis et pallidis neutiquam hujus speciei varietas videtur.

5. *cinereum*, griseo-cinereum basi incrustans, clavulis adscendentibus apicibus albidioribus incisis.

Prope Parisios autumnno crescit in sylvis locis umbrosis. *M. cristatum* quidem refert, sed dimidio fere minus, magis coriaceum et siccum est, et basi in crustam effusum.

†† *Erecta ramosa, ramis distinctis aequalibus.*

6. *palmatum*, fusco-purpurascens, ramis laevibus palmatis, apice albicantibus subnitidis (maleolens).

α . *Merisma foetidum*. *Syn. fung.* p. 584.

Clavaria palmata. *Scop. Carniol.* 2. p. 483.

β .? *angulatum*, caule basi tomentoso, ramississimo: ramis angulatis latiusculis, apicibus flabelliformibus. †

Ramaria palmata. *Holmsk. Coryph. ic.*

Clavaria palmata. *Nees System.* 2. p. 43. t. 66. f. 151. (exclus. synonymo.)

Hab. in Dania. Stipes tomentosus. Substantia durabilis. Sapor austerus, paulum subacris. Odor debilis, sed singularis. Color fungi mediae aetatis dilute badius, ramis rubescentibus; adulti spadiceus, apicibus murini coloris. *Holmsk.*

γ. *flabellare*, fusco-purpurascens tenerius, ramis linearibus subfastigiatis. *Syn. l. c.*

Clavellaria flabellaris. *Batsch El. fung. p. 227. t. 28. f. 159.*

Haecce varietas aut subspecies magis in sylvis frondosis oritur; formam minus evolutam exhibet. Odor etiam debilis est, qui in α est foetidus, illa insuper sylvas praefert abietinas, et 2-3 unc. alta est: ramis compressis dilatatis; β. quantum ex descriptione *Holmskioldi* dijudicari potest, ut species distinguimeretur.

7. *clavulare*, erectum coriaceum dilatato-ramosum inodorum pulverulentum subtuberosum. *Fries Obs. Myc. I. p. 156. t. I. f. I. †*

In sylvis abiegnis Sueciae. Stipes basi subteres, ceterum compressus, digitato-ramosus: ramis inaequalibus sublaciniatis concoloribus. *Fries.*

8. *pusillum*, gregarium minutum fusco-purpurascens, ramis rugosis e basi ortis divaricatis obtusis. †

Clavaria rugosa, Sowerby t. 404.

In Anglia observatum et parasiticum ad stipites vetustos viciae *Fabae*. An fungus adhuc juvenilis? vix 4-6 lin. magnus est et ramuli crassiusculi subfurcati, compressi, glabri. 9. *serratum*, caespitosum subadscendens viscosum fuligineo-albidum, ramis latiusculis serratis obtusis. *Comment. de fung. Clav. p. 106. t. 4. f. 4.*

Rarius in sylvis. Ramosissimum, subcarnosum, totum glabrum. Affinitas cum *Thel. sebacea* adest, sed habitu differt.

Species minus cognitae.

20. *strigosum*, caespitosum dilute rufum canescens, ramis congestis divergentibus striatis.

Comment. l. c. p. 94.

In fagetis et ericetis. Ramis brevibus, strigosis horridum.

21. *coralloideum*, fusco-nigricans submolle, ramis depressis striatis, apice fimbriatis albicantibus. — *Clav. coriacea Bull. p. 198. t. 452. f. 2.*

Ad terram et inter folia emortua, autumn.

* *Clavaria thermalis*, flavo-ferruginea, ramis cylindricis simplicibus vel irregulariter divis. *Decand. Syn. pl. p. 21.*

In thermalibus, trabibus humidis adnascens. Glabra, siccatione durissima.

* *Clavaria subterranea*, teres aut compressa subfusca furcatim aut inaequaliter divisa, tomento brevi vestita. *Decand. l. c. p. 21.*
In subterraneis.

Sed ubi pertineant Clavariae sic dictae *Dematiiformes* Schumacheri Flor. Saelland. p. 407. sine autopsia difficile dictu esset.

††† *HELVELLOIDEI* fungi sunt carnosii vel carnosio-membranacei (in pileo), formaque magis determinata gaudentes; plurimi glabri, stipitati et erecti ac pileolo quodam, sed pro generibus vario, vestito, nec non resupinati sunt. Hymenium laeve, ex thecis manifestis liberis compositum, quae sporulas continent plerumque octo, et in plurimis elastice avolantes.

† *Clavaeformes*, pileo perspicuo nullo.

XLVIII. *CLAVARIA*. *Pileus* (*clavula*) elongatus, teres, simplex aut digitato-ramosus, cum stipite confluens (glaber). *Sporulae* non elastice prosilientes, thecis inclusae. *Thecae* minutae, subcoalitae, omnem fere superficiem occupantes. (*Color albus, pallidus, flavescens. Substantia carnosa.*)

* *Coralloideae, ramosissimae.*

1. *crispa*, ramosissima flavescens, ramis planis crispis margine incisis. — Jacq. *Miscell. austr.* 2. p. 100. t. 14. f. 1. Schaeff. *fung.* t. 163? Alb. et Schwsin. *Consp.* p. 285.

Hab. in Carniolia, Silesia et in Lusatae pinetis. Vix hujus generis; *conf.* *Thelephora*, divis. prima.

"Statura non raro capitis puerilis. Caulis tuberosus. Rami numerosissimi, in foliorum formam compressi, margine argute serrati. Substantia est carnosa sapida; gulae blandiens, hinc fungum speciosum, *Clavariis* esse conge-

nerem, nulli prorsus dubio obnoxium." *Consp. fung. l. c.*

2. *Botrytis*, subdifformis, caul. crasso decumbente albido-pallido, ramis subbrevis in-curvis subrugosis; ramulis obtusis rubicundis. *Syn. fung. p. 587. Schaeff. fung. t. 176. Nees System. f. 150.*

Clavaria plebeja. Jacq. Miscell. 2. p. 101. t. 13.

Clavaria coralloides. Bolt. fung. 3. p. 113. t. 113.

Aestate et autumnò in sylvaticis et ericetis nascitur, sed minus frequens; rarius quoque bene evoluta reperitur, aëre pluvioso excepto, ubi rami magis elongantur et stricti sunt, eorum apices vero tunc minus colorati, Fungus edulis uti subsequentes.

5. *alba*, ramosissima candida, ramis rectiusculis dichotomis acutis.

Ramaria coralloides alba. Holmsk. Coryph. 1. p. 113. t. 12.

Clavaria coralloides. Sow. fung. t. 278. (fig. superior).

Clavaria Holmskoldiana. Fries Obs. Mycol. 2. p. 287.

Clavaria fastigiata. Villars Dauph. p. 1052.
Minus frequens provenit in sylvis abiegnis.

Tota glaberrima et candida, sed basi interdum violascens. Caul. subtenuis, interdum etiam distinctus nullus. Rami dichotomi: nunc recti, nunc flexuosi et teretes. *Fries.*

4. *flava*, caul. subcrasso pallescente ramosissimo: ramis ramulisque luteis. *Syn. fung. p. Schaeff. fung. t. 175.*

β. *divaricata*, magna, ramulis crassiusculis dichotomo-divaricatis.

Clavaria coralloides. Bull. Champ. p. 201. t. 213.

Clavaria muscoides. Bolt. fung. 3. p. 114. t. 114.?

Hab. jam mature post pluvias aestivales in fagetis et quercetis.

Obs. Species magna, minus communis, forma varia. Forte species diversae sunt sub nomine *coralloides* ab auctoribus descriptae, et a Mycologis, quibus occasio datur illas observandi, extricandae et iconibus illustrandae.

5. *formosa*, ramosissima, caul. crasso subdecumbente, ramis elongatis aurantio-roseis: ramulis obtusis flavescentibus. *Icones et Descr. fungor. minus cognitor. Fasc. 1. p. 11. t. 3. f. 5.*

Clavaria fastigiata. Batsch. El. fung. f. 48.?

Clavaria flava. Trattin. Esb. Schwaemm. f. 4. t. 2.

Provenit quoque in sylvis, jam mature. Ab antecedente, cujus tantum varietas?, forma elongata graciliore, ramis magis approximatis, apice aliter coloratis, praecipue differt.

6. *suecica*, gregaria ramosissima erecta ochraceo-incarnata opaca, caul. crassiusculo deorsum albido tomentoso, ramis multifidis acutis. *Fries Obs. Mycol.* 1. p. 256.

In pinetis muscosis Sueciae. Caul. 3-4 lin. crassus erectus, tomento fibrilloso albido radicans. Rami laxiusculi, flexibiles, 1 - 1½ unc. *Fries.*

7. *spinulosa*, ramosissima fuligineo-spadicea seu umbrina, caule crasso, ramis strictis subulato-spinulosis: oppositis brevioribus obtusis. *Obs. Mycol.* 2. p. 59. t. 3. f. 1. *Swartz Act. Suec.* 1811. p. 147.

In Germania et Suecia in sylvis. Species magna, rara. Rami breves, pulvere cinnamomeo obtecti.

8. *stricta*, ramosissima flavescente-fusca, caule tenui, fibrillis radicato, ramis ramulisque adpressis acutis strictis. *Comment. de fung. clavaef.* p. 45. t. 4. f. 1. *Schaeff. fung.* t. 286.

Obs. Meum fungus supra truncos aut ad eorum latera inveni, quo, et pulvere manifesto ramis adperso a consimilibus facile discerni potest. Hinc *Clavaria stricta* Schuma-

cheri (Saell. p. 401.) diversa est, illius enim stipes, secundum auctorem est decumbens, subcrassus; etiam nascitur inter folia dejecta faginea.

9. *Syringarum*, ramosissima helvola opaca, ramis approximatis strictis subglaucis, radice tuberoso-spongiosa.

Sero autumnino et initio hyemis saepius legi hanc clavariam sub Syringis ad terram. Cum praecedente specie, congruit forma, sed differt loco natali et basi tuberoso-spongiosa. Sapor Amygdali amari.

10. *decurrens*, flavo-fuscescens opaca ramosissima inaequalis, caul. tenui, radiculis fibrillosis albis.

In sylvaticis varius prope truncos aridos, seriatim saepius crescens.

Cespitem densum efficit; rami stricti adpressi; ramuli summi brevissimi, subacuti, nullo pulvere seu pruino obtecti. Sapor amarus.

11. *abietina*, gregaria ramosissima flavescens: contrita virescens, caule tenui basi et radiculis fibrillosis albis, ramulis subrugulosis.

Syn. fung. p. 589.

Hab. in Germania, Suecia et Helvetia in abietinis. Magnitudine mediocris, forma dendroidea, pulvere subcroceo oblecta, sapore amara.

12. *apiculata*, ramosissima subalutacea, caule subcrassiusculo, ramis brevissimis verticillatis acutissimis virescentibus. *Fries Obs. Mycol.* 2. p. 1288.

In Suecia ad truncos abiegnos vetustissimos.

Caul. adscendens, basi fibrillosus albidus, mox irregulariter divisus, pallide flavis; detrita non viret. *Fries.*

13. *pyxidata*, cespitosa expallido rufescens, ramis ramulisque subverticillatis, in tubum dilatatis. *Comment. de fung. clavaef.* p. 47. t. I. f. I.

Hab. in truncis in Germania et Suecia. Caul. tenuis, basi pilosus. Rami approximati, apice excavati. — cl. Schumacher (*Flor. Saell.* p. 400.), se hanc speciem etiam in sylvis ad terram nudam legisse, dicit.

14. *amethystea*, ramosissima violacea, ramulis teretiusculis obtusis. *Fries Obs. Mycol.* 2. p. 285. *Holmsk. Coryph. ic.* *Nees Syst. d. Schwamm.*

Hab. in Germania, Dania et Suecia in regionibus magis arenosis et ericetosis, ad terram. Colore facile a ceteris distingui potest.

- β. *brachycera*, ramosa violacea, caule subcrasso, ramis ramulisve brevibus crispis. *Comment. de fung. clav.* p. 61. *Barrel. Pl. icon.* 1261.

15. *fuliginea*, gregaria cinereo-rufescens ramossissima, caule tenui, ramis majoribus crassis compressis, lateralibus subincompletis, ramulis subfastigiatis brevibus acutis.

In sylvulis sat communis ad terram.

Caul. plusquam unc. longus, 2 lin. fere crassus, basi pro more subalbicans, dein in 5-7 ramos rudes divisus, quorum singulus trunco fere crassior est. Ramuli vero creberrimi, sed breves, nunc subulati acuti, nunc furcati obtusiusculi omnes subrugosi, roridi. Totus fungus 2-3 unc. altus et 2 latus. Sapor funginus, sed dein sub-amaricans. —

Obs. Sequentes fungi non diversi videntur

* *grisea*, cinerascens, caule subcrasso, ramis subrugosis, ramulis obtusis inaequalibus. *Comment. de fung. Clav. p. 44.*

* *grisea*, fuliginea, caule tenui, ramis rugosis, ramulis obtusis difformibus. *Fries Obs. Myc. 2. p. 289.*

* *cinerea*, terrestris, ramis remotis apice multifidis. *Villars Dauph. 3. p. 1050. Bull. p. 204. t. 554. (Figura mala).*

16. *cristata*, polymorpha ramosa fuliginea cinerascens albaque, ramulis tenuissimis fimbriatis obtusisque. *Syn. fung. p. 591. Holmsk. ic. Bull. t. 358. f. c.*

Clavaria nivea. Comm. de Clav. p. 48. t. 2. f. 4.

β. fallax, elongata subsimplex, indurata fuligineo-cinerascens, basi sphaerulis nigropunctata. *Comment. l. c. Sowerb. fung. 3. t. 307.?*

Frequens autumnno in sylvis frondosis crescit. Magnitudine mediocri. Singulare quod morificans non solum colore mutetur, sed saepe *sphaeria?* quadam parasitica vexetur; forma parum et substantia commutata.

17. *chionea*, ramosissima nivea, caule tenuissimo mox ramosissimo, ramulis elongatis tenuissimis inaequalibus acuminatis.

Prope Parisios, Septembr., ad terram inter muscos mihi obvia est elegans haec species, colore niveo distincta; caul. attamen parum rufescit et sub exsiccando tomento tenui obtegitur. Ramuli setacei. Biuncialis.

18. *gracilis*, ramosa ex albido pallescens, caul. tenui: radiculis fibrillosis, ramulis multifidis subincarnatis. *Syn. fung. p. 592.*

In Germania in pinetis ad terram. In omnibus partibus tenera et gracilis. Rami subangulati, recti; ramuli subsetacei.

19. *umbriata*, candida caule tenui villosa, ramis subcompressis elongatis: ramulis confertis setaceis. *Commentat. de fung. clav. p. 49.*

In abietinis hercynicis prius reperta. Caul. basi villosus. Rami satis crassi et longi, apice setaceo-incisi et hinc fimbriati.

Obs. Hae tres enumeratae species sibi subsimiles, et, forsitan, solis ex verbis difficillime extricandae, sed inter alia, loco natali et regione jam dissentiunt.

20. *trichopus*, nivea ramosissima, caule tenui hirsuto ramosissimo: ramulis furcatis palmatisque subfastigiatis. *Comment. l. c. t. 4. f. 3. Wallr. Ann. Bot.*

Hab. in sylvis ad terram. Magnitudo ejus, uti plerarumque antecedentium, media est. Rami subcrassi, non cristati, vel setacei, sed simpliciter furcati.

21. *macropus*, subramosa alba, caul. tenui elongato pallescente, ramis crassiusculis oblongis, ramulis furcato-subulatis. *Comment. p. 51. t. 1. f. 2 Swar tz Nov. Act. Suec. An. 1811. p. 152.*

In fagetis ad terram. Species subparva, sed caulis sat longus, uncialis, et lin. 1-2 crassus, subflexuosus, glaber. Rami stricti, elongati, teretes.

22. *grossa*, candida, caul. subtenui glabro ramosissimo, ramis crassis dilatatis subdifformibus, ramulis furcatis obtusis. *Comment. p. 50. t. 2. f. 2. †*

In sylvis ad terram. 3 unc. fere alta, 3 lin. crassa, fragilis.

23. *subtilis*, parvula subramosa alba aut citrina, ramis aequalibus elongatis subfastigiatis.

Comment. p. 51. t. 4. f. 2. Swartz l. c. p. 151.

In sylvaticis. Forma regulari distinguitur.

Unc. $1\frac{1}{2}$ alti, in ramos paucos divisa. — Rami dichotome furcati, et fastigiati, ramulis aequalibus attenuatis. Color albo flavoque variat. *Sw.*

24. *crocea*, minuta crocea ramosissima, ramis ramulisve conformibus subfuscatis. *Icon. et*

Descript. fung. p. 36. t. 11. f. 6.

Etiam ad terram nudam in sylvis, sed perraro occurrit, $\frac{1}{2}$ unc. vix magna, *Croco* similis.

25. *muscigena*, parva ramosa fusco-pallida, stipite brevissimo mox partito in ramos divaricatos: ramulis longiusculis subuliformibus. *Schum. Flor. Saell. p. 400. †*

In Dania inter muscos ad arborum truncos.

26. *digitata*, fusco-pallido, stipite angustiore subdigitatim ramoso: ramis incrassatis lobato palmatis obtusis. *Schum. l. c. p. 401. †*

In pratis sylvaticis. Octobr., in Dania.

27. *pratensis*, cespitosa parva flavescens, caule brevi, ramis geniculatis divaricatis: ramulis subfastigiatis obtusis. *Comment. de fung.*

Clav. p. 52. t. 4. f. 5. Bull. t. 358. fig. d. e.
Vaill. bot. t. 8. f. 4.

Clavaria muscoides. Schum. p. 402. Flor.
Dan. t. 836. f. 2.

Clavaria fastigiata. Decand. Flor. Gall.
2. p. 100.

In pratis aut collibus apricis, autumnosae frequens; sed inter gramina vel muscos delitescit. Vix unc. alta.

28. *corniculata*, subramosa helvola, caule gracili elongato, furcatim diviso, ramis acutis obfuscisve subbrevis. *Syn. fung. p. 589.*
Nees System t. 16. f. 152.

β . *Clavaria fastigiata. Schum. l. c. Flor.*
Dan. t. 775. f. 3.

In sylvis ad terram. Caul. basi pubescens, sat longus, sed tenuis. β . inter muscos, in pratis et pascuis; truncus hujus, secundum auctorem, est breviusculus, rami subcompressi dichotomi, et color luteo-ferrugineus aut fuscus.

29. *vitellina*, cespitosa laete flava, caule tenuissimo simpliciter ramoso, ramulis furcatis obtusis fuscis,

Hab. in sylvis umbrosis ad terram. Altitudine modo unciali, sed caespitem sublatum sistit.

30. *viscosa*, truncigena, vitellina viscosa tenax, ramis ramulisque furcatis; caule tenui radicato. *Comment. de Clav. p. 53. t. 1. f. 5. Schaeff. t. 174.*

Incolit truncos vetustos abietinos. Unc. 1-2 alta.

31. *tenacella*, fasciculata fuligineo-violascens subtenax, ramis congestis elongatis, ramulis furcatis obtusis. *Comment. p. 47. t. 3. f. 5. †*

In sylvis, ericetis ad terram rarius. Subcartilaginea quasi, demum cinerascit. Rami non valde divisi, approximati; 1-2 lin. crassi, 3 unc. circiter longi aequales.

32. *coriaria*, alba fragilis caespitosa ramosa, ramis erectis elongatis crassitie aequalibus subacutis. *Rebent. Flor. Neomarch. p. 378. Cl. subcaulescens.*

In pulvere coriariorum, magnos cespites format. Altitudine unciali ad $1\frac{2}{3}$. *Reb.*

Obs. Ob crescendi locum ulterius inquiri meretur, ita ut descriptio, nimis brevis, novis suppleatur observationibus.

33. *palmata*, pallescens carnosula subalutacea, caule tenui, ramis subcompressis palmato-digitatis acuminatis. *Comment. p. 45. †*

Passim in fagetis oritur. Caespitem 3 unc. latum et longum efficit. An forte var. depauperata Cl. *flavae*?

34. *mucida*, gregaria minuta subramosa albida, apice ochraceo, basi tomentulo cincta. *Comment. p. 55. t. 2. f. 3. Consp. fung. Lusat. p. 288. Swartz N. Act. Suec. p. 153.*

Ad latera truncorum putrescentium et mucosorum late se propagat pulchella haec species, longitudine trilineari; individua ad-sunt alia simplicia, alia furcatim divisa, tenuissima, apice flavescentia.

35. *Byssiseda*, varia subramosa parva rufescente pallida (albaque), radiculis longis byssoides insidens. *Syn. fung. p. 151. Fries Obs. Mycol. I. p. 157. Comment. de Clav. p. 55. t. 3. f. 7. (var. major).*

Ramaria fimbriata. Holmsk. Coryph. p. 98. ic.

In ramis querneis, aut ad Salicis *capreae* truncos sat rare provenit. Radix (subiculum?) magna, pro fungillo, nunc fibrillosa nunc ut Byssus alba plura specimina conjungit et corpuscula subjacentia obteggit. Ramii juniores sunt plerumque simplices et curvati, majores furcati, $\frac{1}{2}$ -1 unc. longi, etiam albi.

* *byssacea*, cespitosa minuta simplicissima, corniculis albis, basi unitis, lana candida, insidentibus. *Bolt. in Usteri Annal. I. p. 11. t. I. f. 5.*

Ad salicum annosarum praecipue cortices, hyeme crescit, dicit cl. Auctor. Cornicula

alba, acuminata, bilinearia fere, simplicissima, sed basi unita, parum incurva et lana tenuissima sed densa progerminantia, siccata in fuscum colorem tendunt, sed lana manet candida.

Obs. An clavulae semper simplicissimae? An a priore ut species sejungenda? Forsitan in duabus hisce anomalis clavariis, radiculae sic dictae modificatio sunt *pilei* obsoleti, et tunc hi fungi non male genus peculiare pone Odontias constituerent: clavulis pro *subulis* habendis.

*** Clavulis elongatis incrassatis obsolete ramosis.*

36. *rugosa*, gregaria submagna incrassata albidula *rugosa* simplex ramosaque: ramis parvis difformibus. *Bull. t. 448. f. 2. Flor. Dan. t. 1301. Bolt. fung. t. 115.*

Clavaria coralloides. *Sowerb. t. 278.* (Fig. inferiores.)

Vulgaris in sylvis, aut inter gramina ad margines sylvarum.

Mere variat forma et longitudine sic et tenuitate, qua de re nec non de pluribus synonymis videris *Comment. de fung. Clavaef. p. 57.*, quibus addi posset ob colorem: *β. flavescens*, subparva flavescens, ramis obsoletis divergentibus. — Inter Muscos, $1\frac{1}{2}$ unc. longa. *Merisma* refert.

37. *ceranoides*, caespitosa magna flavo-fusca, clavulis difformibus subramosis dentatis. *Syn. fung. p. 594.*

Clavaria rugosa. Sowerby fung. p. 235.

Hab. in Anglia. Species magna, cujus clavulae numerosae cespitem efformant latitudine palmari; substantia sunt subflaccidae, interne cavae, sursum incrassatae, subspadiceae.

38. *ignea*, caespitosa flavo rubra, deorsum ventricosa, ramulis paucis tenuibus brevibus. *Mich. Gen. pl. p. 209. t. 88. f. 2. †*

Hab. in Montis-Boni ericetis et sylvis caeduis, Octobr. *Mich.*

*** *Simplices, seu clavulis non divis.*

39. *pistillaris*, solitaria magna incrassata e flavo seu alutaceo subrufescens, apice rotundato. *Schaeff. fung. t. 169. Batsch Elench. fung. t. 2. f. 46. Comment. de Clav. p. 61-65. (Ob critica et observationes.)*

In sylvis frondosis, sed praecipue in regionibus calcareis nascitur. Iunior fere cylindrica et colore dilutior. *Americans. 4-5 unc. longa, farcta, subdura.*

β. *herculeana*, subspadicea rugosa, apice subdepresso. *Syn. fung. Holmsk. Coryph. ic. Schaeff. t. 169. f. 9. Sowerby fung. t. 277. Michel. Gen. t. 87. f. 2.*

In sylvis acerosis provenit. Major est et colore obscuriore fere spadiceo differt. — Substantia mollis gossipina. S.

40. *alba*, solitaria submagna tota alba incrassata subturbinata. *Traité sur les champ.* p. 255.

Clavaria major alba. *Micheli N. G. Pl.* p. 208. *Battar. Fung. Arimin.* p. 23 t. 3. f. a.

Hab. in Italia. Antecedente paullo minor, sed pro ratione parum crassa, extus intusque alba, solida. Esculenta ex Batana.

41. *Ligula*, subcaespitosa rufescente pallida opaca basi villosa attenuata, sursum teres subcompressaque. *Schæff. Fung.* t. 170. *Comment. de fung. clav.* p. 66.

Hab. in sylvis abietinis ramulis villo adhaerens. 2 unc. longa, 3 lin. crassa. Forma subvariabilis. Color ut in Cl. *pistillari*.

Obs. Species *Schumacheri* Saell. no. 2014. diversa? qui "fusco-flavescens, longa, radiculis penicillatis, et in pratis graminosis, aut ericetosis subhumidis insuper vegetat."

42. *cochleariformis*, caespitosa flavo-fusca, radiculis penicilliformibus e tubere ortis, clavula compresso-excavata apice subretusa. — *Schumach. l. c.* p. 407. †

In Daniae ericetosis aridis. Novembr. Color basi et in parte convexa clavulae dilutior. S.

43. *tuberosa*, flavescent, clavulis attenuatis in basi tuberosa hirta coalitis; intus cavis. —

Sowerby fung. t. 199. †

Hab. in Anglia ad truncos aridos, sub quorum cortice tuber absconditum est. 1 - 2 unc. longa, lin. 1. crassa, subflexuosa.

44. *bulbosa*, subgregaria minor flexuosa ferrugineo-aurantiaca, clavula subincrassata elongata obtusiuscula, basi bulbosa albido-tomentosa. *Schum. Saell. p. 405.*

Cresc. in Dania inter muscos in sylvaticis humidis, $1\frac{1}{2}$ unc. longa, interne carnosogelatinosa. *S.*

45. *junceae*, gregaria subcaespitosa tenuis argillacea rufescensque, basi villosa-fibrillosa. *Fries Obs. Myc. 2. p. 291. Conspect. fung. Lusat. p. 289.*

Clavaria rufescens, triuncialis, tenuissima. Michel. gen. t. 208. no. 11. Syn. fung. p. 600?

Clavaria hirta. Flor. Dan. t. 1257.

β? *pilipes*, rufescens solitaria gracilis, stipite inferne piloso. *Flor. Dan. t. 1100. f. 3.*

In nemoribus humidis, umbrosis; ad radices arborum, inter fol. decidua.

— "Clavula fistulosa, longissima, juncea, subfiliformis, apice acuto, nonnunquam subincurvo, 3-5 unc. longa, basi radiculis fibrosis. Niveis in orbem expansis fol. humentibus

bus infixa." *D'Alb. et de Schweinitz. l. c. p. 290.*

46. *fistulosa*, gregaria, flava, radice fibroso - barbata, clavula incrassata obtusa fistulosa longissima. *Syn. fung. p. 599. Holmsk. Cörryph. I. ic.*

Hab. in Dania ad truncos et ramos deciduos. Substantia mirae tenuitatis, ita ut fungus deficiente humore, contrahatur et digito Junci instar complicari queat. *Holmsk.*

47. *strigosa*, subgregaria, fusco-lutea aut pallida, clavula longissima incrassata fistulosa, caule inferne pilis strigosis obsito, basi nigricante. *Schum. Saell. 2. p. 405.*

Clavaria fistulosa. Flor. Dan. t. 1256.

In Daniae pinetis et alnetis, ramulis rejectis adnascens. Octob. — Novembr. Una cum stipite, qui plerumque est longissimus, 4 - 10 unc. longa, deorsum 1 - 2 lin. crassa. *Schum.*

48. *pilosa*, gregaria, fuligineo-umbrina, clavulis subincrassatis primo pilosis demum glabris. *Comment. p. 74.*

Clavaria fistulosa, subfuliginea teres fistulosa gracilis, pilis deciduis operta. *Decand. Syn. pl. gall. p. 21. Bull. champ. p. 213. t. 463. f. 2.*

Hab. in Gallia, ad folia semiputrida. Fragilis, lin. crassa et 1 - 2 unc. longa.

49. *Ardenia*, gregaria cinnamomea, clavula incrassata longissima cava, basi tomentosa.

Syn. fung. p. 599. Sowerb. t. 215. †

Hab. in Anglia ad ramos exsiccatos. Altitudine insignis, et pro longitudine sat crassa.

Clavariae fistulosae Holmsk. valde affinis videtur.

50. *macrorhiza*, solitaria simplicissima, clavula fistulosa flavo-fusca, basi contorta, radice longissima villosa. *Fries Obs. Mycol. 2.*

p. 292. Swartz in Act. Suec. an. 1811. p. 155. t. 4. f. 1.

Hab. in Suecia in locis declivis arenosis. In terram longissima descendit usque ad 3 uncias et ibi fibrillis densis albidis vestita. Stipes inferne subtorquatus, sulcatus. Clavula ipsa vix distincta, obtusa, leviter compressa. Color albo-caesius, demum fuscus. *Sw. l. c.*

51. *fusiformis*, caespitosa congesta aurea, clavulis attenuatis, basi subcohaerentibus subtenacibus. *Syn. fung. p. 601. Sowerb. t. 234.*

Clavaria fasciculata. Comment. p. 73.

In sylvis ericetosis subnudis habitat, autumnii initio. Clavulae plurimae fasciculum efficiunt basi cohaerentem, utrinque sunt attenuatae, 3-4 unc. longae, apice obscuriores.

52. *inaequalis*, caespitosa inaequalis fragilis lutea, clavulis acuminatis angulato-diformi-

bus. *Fries Obs. Mycol.* 2. p. 193. *Flor. Dan.* t. 836. f. 1. †

Clavaria vermiculata *Sowerb.* t. 253. ?

Autumni initio in campis et pratis siccis apparet post pluvias aestivales. Clavulae 2-4 unc. altae, forma variae: falcatae, contortae, compressae, angulatae. *Fries l. c.*

Obs. An a *Clavaria helvola* satis diversa?

β. albo-lutea, caespitosa fasciculata, clavulis ventricosus: aliis integris, aliis eroso-bifidis.

Sowerb. l. c. p. 253. (fig. superiores).

Provenit in graminosis. Majuscula, sub-turgida. Clavulae albae, apice citrinae ibique incisae. (*Quoad iconem*).

* *fasciculata*, caespitosa carnosae aurantiacae glabra, stipitibus ad basin fasciculatis, ad apices compressis et in senio laceris. *Decand. Syn. Gall.* p. 20.

Hab. in sylvis ad terram. Clavulae utrinque acuminatae. Est etiam clav. *fasciculata* *Villarsii*.

53. *luteola*, gregaria farcta alutacea oblongo-clavata teres. *Comment. de clav.* p. 66-67.

Clavaria caespitosa, caulibus simplicissimis, basi nonnunquam unitis oblongo-clavatis farctis aquose luteolis. *Jacq. Miscell.* 2. p. 89. t. 12. f. 2. *Schmiedel* t. 5. f. 1.

Nascitur ad terram in sylvis abiegnis, autumno. Clavulae cespitosae, aggregatae, carne alba compacta farctae, 1 - 2 unc. longae. *Wulf. l. c.*

54. *ericetorum*, cespitosa flavescens, clavulis incrassato-subcylindricis obtusis (passim divis) alutaceis, stipitibus subnitente flavis. *Obs. Mycol. 2. p. 60. Conspect. fung. p. 290. Schmiedel Ic. pl. p. 56 - 59. t. 15.?*

Hab. in ericetosis, sylvaticis frondosis.

Color recens alutaceo-flavescens, in fungo exsiccato fere argillaceus, opacus. Clavulae carnosio-ceraceae, in quibusdam ejusdem cespitis individuis, nonnunquam compresso-cornutae, adultae rugulosae, 1 - 2 unc. longae, 2 lin. crassae, farctae, basi equidem non angustae, ut in sequente.

55. *helvola (pascua)*, gregaria et subfasciculata, clavulis elongatis sensim incrassatis obtusis, flavo-aurantiis, stipite tenui dilutiore.

Clavaria helveola. Comment. de clav. p. 69.

Clavaria cylindrica. β. lutea. Bull. t. 463. f. 1. b. n. o.

Clavaria lutea. Decand. Flor. Gall. 2. p. 97.

Clavaria teres. Baumgart. Flor. Lips. t. 4. f. 2. a.

Clavaria angustata. Consp. fung. p. 290. (ut e descriptione patet).

Clavaria fragilis et β . *lutea* Schum. *Saeland.* p. 404.

Nascitur vulgaris haec species, in pratis siccis et pascuis, inter muscos ad margines, passim et in interiore sylvarum.

Obs. Pro aëris et soli conditione magnitudine variat, plerumque unc. 1 ad $1\frac{1}{2}$ - 2 unc. longa, lin. 1 et $\frac{1}{2}$ lata, fragilis, solida, laete flava, nonnunquam apice, sed non semper et ubique in locis, praesertim sub exsiccatione ferruginea vel cinnamomea fit, quod ex differentia specifica, ne difficultas in determinatione oriatur, excludendum videtur.

Sequentes fungi ut laeves varietates aestuandae, nec tamen negligendae sunt.

β . *dispar*, dilute flavescens minus alta, clavulis incrassatis: una alterave compressa subbicorni.

Clavaria fragilis, var. *lutea*. Holmsk. *Coryph. ic.*

Clavaria vermicularis. Sowerby t. 253.?
(Fig. inferiores, praesertim sinistrae; fig. intermedia ad individuum luxurians picta).

Clavaria pistillaris. Bolt. fung. 3. t. 113.?

In pascuis inter gramina brevia et muscos.
y. angustata, cespitosa flavescens, clavulis elongatis subadscendentibus, deorsum flexuosis. *Comment. de clav.* p. 72. t. 1. f. 3.

In pratis montosis muscosis subhumidis, quo clavulae longiores et multo angustiores factae sunt.

δ. *aurantia*, subcaespitosa parva aurantio-rubra, clavulis subrugosis aut canaliculatis, basi tomentosis. *Obs. Mycol.* 1. p. 32.

Magis in sylvis ad terram nudam adest, cui tomentulo adhaeret. Clavulae subdistant, unc. longae.

* *sylvatica*, dense caespitosa subargillacea, clavulis congestis longissimis angustis incrassatis. — *Comment.* p. 73.

In sylvis frondosis rarius. In eodem fasciculo clavulae numerosae, 40-50, sunt, basi fere cohaerentes, 4-5 unc. longae, apice subcompressae et obtusae, intus solidae, exsiccatae subcinerascunt. *Com.*

56. *flavipes*, caespitosa fragilis, clavulis falcatis albido-pallidis acuminatis, stipitibus nitente-flavis. *Comment.* p. 75. t. I. f. 4. — *Schumacher.* p. 404? apice fusco-umbrino.

In collibus graminosis apricis. *Stip.* subpellucidus, citrinus.

57. *fumosa*, caespitosa, fragilis cinerascens unicolor, clavulis congestis fumosis flexuosis. *Comment.* p. 76.

In sylvis ad terram, fasciculum multarum clavularum, basi subconnatarum offert. Fun-

gus recens aquose fuliginus, et fere hac nota ab *eburnea* modo differt.

β .? *striata*, caespitosa subfuliginea, clavulis longissimis flexuosis acuminatis, sparsim striatis. *Comment.* p. 75.

In sylvis.

58. *eburnea*, caespitosa fragilis nivea, clavulis congestis subfistulosis attenuatis, basi subpellucidis. *Syn. fung.* p. 603.

Clavaria fragilis. *Holmsk.* var. *alba*, icon. fig. f. 9. *Flor. Dan.* t. 735. f. 1. *Comment. de clav.* p. 77. cum synonymis.

β . *Clavaria gracilis.* *Bolt.* 3. t. III. et *Sowerb. fung.* t. 232. ? (minor minusque fasciculata in iconibus exhibentur et in hortis crescit),

δ . *tubiformis*, caespitosa candida, clavulis flexuosis deorsum incrassatis, intus fistulosis. — *Micheli*, p. 209. t. 87. f. 13.

Etiam haec species in nemoribus praesertim gignitur, locis umbrosis humidiusculis, fasciculum multarum clavularum basi subconnexarum aut separatarum sistens. Caulis basi hyalinus, non equidem tenuis, uti in sequente; β . ob cavitatem insignem, pro icone, forsitan distincta.

59. *pistilliforma*, gregaria candida, stipite elongato subtenui, clavulis incrassatis obtusis intus cavis.

Clavaria alba pistilliforma. Vaill. Bot. p. 39. t. 7. f. 5. — Mich. gen. pl. p. 209. t. 87. f. 10.?

Clavaria cylindrica. Bull. p. 212. t. 463. f. 1. Comment. p. 86. Sowerb. t. 90. (ubi nonnulla individua valde turgida picta sunt).

Clavaria fragilis. Holmsk. Coryph. ic. fig. a — c.

Hab. in pascuis apricis, autumnii initio. Clavulae equidem non cylindricae, saltem non in omnibus, apice obtusae, quo praesertim ab antecedente differt, qua etiam minor, nec basi unita. Ulterius tamen observari meretur.

60. *vermiculata*, cespitosa similis alba, clavulis farctis attenuatis subincurvis.

Clavaria solida. Comment. p. 78.

Clavaria vermiculata, non fistulosa candida. Micheli Gen. pl. p. 209. t. 87. f. 12.

Eam reperi, circa Parisios, mense Septembri, in sylvis fasciculatim crescentem; sed specimen non altum fuit, $1\frac{1}{2}$ unc.; basin versus non attenuata. Nonnullae clavulae subcompressae et apice subdivisae, quomodo icon Micheliana hoc etiam reddit.

β. *flexuosa.* Cl. *vermicularis*, cespitosa albissima cylindracea subulata flexuosa solida. Swartz in N. Act. Suec. p. 159. f. 2.

Hab. in Suecia. Caespitosa tripollicaris.

Stipites vix distincti. Clavulae cylindricae, crassitie pinnae columbinae, subflexuosae, apice subulatae, arcuatae, intus solidae, medio perforatae et humorem scatentes. *Swartz.* — A meo fungo, quem multo superat, forsan diversa.

61. *rosea*, subcespitosa obscure rosea fragilis, clavulis elongatis incrassatis farctis. — *Fries Obs. Mycol.* 2. p. 290. t. 5. f. 2.

β. *rubella*, subsolitaria recta incrassata aquose rubella. *Comment. de clavaef.* p. 81.

In duobus speciminibus hic, anno pluvioso 1816, a me collectis clavulae simul in minori copia, 3-4, aderant, sed etiam valde fragiles, 3-4 unc. longae. Apex secundum D. *Fries*, exsiccatione flavescit, quod non observavi.

62. *purpurea*, cespitosa subcompressa acuminata purpurea. *Syn. fung.* p. 600. *Flor. Dan.* t. 837. f. 2. *Fries*, l. c.

In Daniae et Sueciae herboris. Color rufo-fuscus, *rufa*, gregaria incrassata compressa rufa passim emarginata. *Flor. Dan.* t. 775. f. 1. *Syn. fung.* l. c.

In graminosis sylvaticis, in Dania.

63. *nigrita*, cespitosa nigra, clavulis longissimis fistulosis subreflexis. *Comment.* p. 79. *Swartz*, l. c. p. 159.

Rarissime in graminosis. Clavulae subcompressae, flexuosae attenuatae, 3-4 unc. longae,

in specimine Holmiae lecto, secundum Swartz, tantum unciales.

64. *falcata*, parva solitaria candida, clavula falcata obtusa, stipite brevi pellucido. *Comment. p. 81. t. 1. f. 3. — Mich. Gen. pl. t. 87. f. 6.* quoad iconem.

Occurrit satis frequens in umbrosis subhumidis ad terram nudam, sed solitaria et parca; unc. 1. longa, tenera.

Obs. Fungillus *Michelianus*, t. 87. f. 9., crescit in fictilibus hortorum terra stercoreata repletis.

65. *exilis*, subcespitosa minuta candida, clavula incrassata obtusa acutaque substricta, stipite pellucida.

Mense Novembri in pulvere coriario putrescente, prope Rouen inventa (*Havet.*), et ibidem satis profusa orta. Exigua, 2 - 3 lin. tantum longa, $\frac{3}{4}$ vix crassa, firmula.

66. *acuta*, subgregaria alba, clavula conica acuminata *farinacea*, stipite elongato. — *Sowerb. fung. t. 333. †*

In Anglia ad terram, $\frac{3}{4}$ unc. circiter alta. An *Isaria*? sed in insectis emortuis non parasitica.

67. *virgultorum*, gregaria conferta, primo albidula dein fuscescens, clavulis elongatis filiformibus, basi villosis.

Crescit in umbrosis ad stipites et folia sicca; eam semel quidem (prope Versailles) fine Octobris inveni, sed tunc magna in copia. Clavulae crassitudine fere aequales, medio tamen paullo crassiores, 2 unc. ad $2\frac{1}{2}$ longae; valde tenues, basi villositate albida ramulis adhaerent; recenti in statu aquosae candidae, exsiccatae pallide fuscae.

68. *filata*, gregaria longissima tenuis filiformis glabra albicans. †

In uliginosis inter muscos observavit D. Chaillet. Spithamam fere longa, sed vix dimidiam lineam crassa; basin versus paullo tenuior.

69. *tortilis*, subgregaria parva, clavulis elongatis teretibus aut subincrassatis candidis, stipite subtortili glabro spadiceo.

Hab. in Gallia meridionali in ramulis et foliis aridis. *Dufour*.

70. *pusilla*, gregaria minutissima alba, crassitudine ubique aequali. *Comment. p. 86. t. 3. f. 6. Conspect. fung. Lusat. p. 292.*

In folio quodam arido inventa. Serotina, nunc striata, nunc curvata, lin. 1. alta.

71. *diaphana*, epiphylla sparsa minutissima basi bulbo ovato flavescente pellucido, clavula attenuata incurva diaphana (?) candida. *Schum. Saell. p. 405. †*

Supra folia decidua faginea, Octobri, haud infrequens. 2 lin. fere longa. *Schum.* An forte eadem ac *Clav. gyrans*?

72. *pallida*, gregaria minuta diaphana, stipite longo flexuoso albido-caesio subbulboso, clavulis ovato-oblonga ex pallido subrosea. *Schum. l. c. p. 406.*

Hab. in terra humosa pineti. Sept. 2 - 3 lin. longa.

73. *muscicola*, gregaria minuta tota alba, clavula incrassata basi bulbosa. *Obs. Mycol. 2. p. 60. t. 3. f. 2.*

Clavaria muscicola, candida simplex elongata clavata, tuberculo radicali obscuriore. *Nees Syst. der Pilze und Schwaemme. 2. p. 43. t. 16. f. 154.*

74. *uliginosa*, simplex minuta nivea, basi nuda attenuata capillari in clavulam obtusam teretem abiens. *Wallroth Annus botan. p. 141.*

In uliginosis. Vix semiuncialis, fragilis. *W.*

**** *TYPHULAE*, *firmulae*, stipite magis distincto et sublongo instructae.

† Tuberculo radicali nullo.

75. *fuscipes*, clavula subincrassata flavo-cinnamomea, stipite longiusculo glabro nigro-fusco.

Rarius ad ramulos siccos prope Parisios. *Thuill.*

Inter minores major est, $\frac{3}{4}$ unc. fere longa, $\frac{1}{2}$ lin. crassa. Clavula subflexuosa, subsulcata, obtusa. Stip. dimidio fungi longitudine.

76. *chordostyla*, stipite elongato capillari albidoglabro, clavula sublineari flavo. †

Mitrula Todei. *Fries Obs. Mycol. 1. p. 160.*

Typhula Todei, pileo flavo, stipite albo glaberrimo, tuberculo nullo. *Fries Obs. 2. p. 298.*

Chordostylum Clavaria. *Tod. fung. Mecklenb. 1. p. 37. t. 6. f. 55.* Ex D. Fries, sed secundum b. auctorem capitulum coriaceum est, hinc potius ad fam. *Sclerotoideam* pertineret.

In stipitibus locis humentibus dejectis, praecipue in Filicibus. Aug.—Septemb. *Fr.*

77. *minuta*, gregaria minutissima, clavula lineari lanceolata cinnamomea, stipite pallido.

Clavaria minuta. *Sowerby fung. t. 391.*

Mitrula pusilla. *Fries Obs. Mycol. 2. p. 299.*

In Anglia ad involucra sicca Dipsaci pilosi, forte etiam ad caules aut folia carduorum crescit.

78. *ovata*, tota candida, clavula brevi obovata, stipite pellucido. *Comment. de fungor. clav. p. 85.*

Autumno in ramis Aridis, Rubi fruticosi, etc. Clavula subcava, passim subcompressa, stipite duplo brevior.

79. *obtusa*, gregaria minuta candida, clavula oblongo-ovata obtusa, stipite brevi concolore.

Clavaria obtusa. *Sowerb. fung. t. 334. f. 1.*

Clavaria quisquiliaris. *Fries Obs. Mycol. 2. p. 292.*

In stipitibus putrescentibus *Pteridis Aquilinae* in Anglia et Suecia, et in caulibus exsiccatis *Sonchi alpini* in Vogesia et Helvetia.

Clavula firmula, satis crassa, etiam subcompressa et ligulata, stipite longior. Totus fungillus 3 lin. altus. Varietatem stipite pubescente etiam delineavit cl. Sowerby; observare tamen licet, stipes in non paucis fungis exsiccando subtomentosus fuit.

80. *Filicina*, sparsa elongata subfiliformis pallide rufescens glabra.

Hab. in stipitibus aridis *Filicis Maris* in Vogesia (*Mougeot*) et circa Parisios in iisdem stipitibus, locis humidis dejectis (*Desvaux*).

Totus fungillus unc. 1 altus, $\frac{1}{2}$ lin. crassus; clavula ipsa stipite brevior eoque paullo crassior, teres et subacuta. Color in recenti forsitan albidus. Substantia sicca.

81. *peronata*, simplicissima albescens, clavula brevissima subacuta, stipite longiusculo pubescente.

In Vogesia rarius ad strobilos *Pini sylvestris*.

Habitu parum alieno distincta ob clavulam vix 2 lin longam, et stipitem membrana villosa quasi vestitum; hic $\frac{1}{2}$ - $\frac{3}{4}$ unc. longus et villositate sat densa matrici adnata.

†† *Stipite elongato tuberculo radicali insidente.*

82. *Erythropus*, clavula cylindrica brevi alba, stipite stricto elongato fusco-rubente. *Comment. p. 84. Syn. pl. t. 3. f. 4.*

Clavaria gyrans. Bolt. fung. t. 112.

Typhula Erythropus. Fries Obs. Mycol. 2. p. 297.

β ? *pinicola*. Cl. *Erythropus*, gregaria minutissima, stipite elongato capillari atro-rubente, apice albo villosa, clavula oblonga, subcylindrica vel ovata glabra albido-pallescente. *Schum. Saell. 2. p. 406.*

Non equidem (α) valde rara occurrit in ramulis aridis, sub quorum cortice tuberculum basitare latet, huc rugoso-sulcatum est et fusco-nigricans subdepressum. Stipes vix pubescens, subpurpureo-badius, apice albicans, tenuis. Clavula 2-3 lin. longa, subobtusula, alba. — Fungillus *Schumacheri* (β) etiam ob locum naturalem: in trunco putrido Pini *sylvestris* diversus videtur.

83. *gyrans*, gregaria candida, clavula cylindrica, stipite longo pubescente. *Comment. p.*

83. *Batsch El. fung. t. 164.*

Typhula gyrans, pileo stipiteque pubescente albis, tuberculo pallido, interdum nullo. *Fries l. c. p. 297.*

Hab. ad folia dejecta et ad stipites aridos. Valde debilis, minutissima. Tuberculum laeve, pallidum.

84. *granulata*, tota alba, clavula fistulosa, tuberculo granulato nigro. — *Willd. Prodrum. p. 705. f. 17.*

Hab. Berolinum.

85. *Rhacorrhiza*, clavula subulata alba, stipite glabro flexuoso subfusco, tuberculo lenticulari nigro. *Reichard in Act. Berol. I. p. 515. t. 9. f. 4-5. Sowerb. fung. t. 233.*

Hab. ad terram madidam. Filiformis, flexuosa, 2-3 unc. longa. Secundum D. Fries, Stipes est albus et tuberculum globosum.

Obs. Clavulae in speciebus hujus subdivisionis aut subgeneris, natura omnino eadem videntur antecedentium. Et nondum probatum est, an tubera illa radicalia ad hosce fungillos pertineant, aut an propria vegetabilia sint: *Sclerotia*, etc. Talia tubercula etiam ad basin stipitum nonnullorum agaricorum observantur.

XLIX. PHACORHIZA. *Tuberculum* radicale volvaceum, carnosum; primo includens *clavulam*, eamque per fissuram deinde protrudens.

1. *Scle-*

1. *Sclerotibides*, sparsa, clavula simplice alba, tuberculo badio. *Tab. XI. Fig. 1 et 2.*

Hab. in Vogesia in caulibus exsiccatis *Sonchi alpini* et *Cacaliae albifrontis*. (Mougeot.)

Tuberculum globosum, ab initio laeve, in statu vetusto et post emissionem clavulae col-
labitur et rugosum fit; insuper etiam ante ma-
turitatem, solum reperitur, quare facile pro
Sclerotione Serien haberi posset, cujus colore
et forma est. Clavula ipsa, (passim duae ad-
sunt) ex medio tuberculi, rarius ad latus, exit,
carnosa est et formam communem, et uti vi-
detur etiam structuram specierum simplicium
generis antecedentis participat; sed ab illis,
praesertim quae in ultima divisione enumeran-
tur, singulari evolutione discrepat, quare ut
genus peculiare hunc fungillum proponere vel-
lem.

Obs. Non confundi debet cum *Clavaria Sclerotioides* Decand. Flor. Gall. Suppl. p. 29, quae ob naturam totam *Sclerotii* ad Familiam *tuberaceam* pertinet, ejusque clavula continua-
tio est tuberculi, ac eodem colore.

L. GLOBOSUM. *Clavula* carnosa, sub-
compressa, margine subprominente a stipite di-
stincta. *Thecae* majores, disjunctae. (Color
obscurus: plerumque niger, fuscus, interdum
virescens).

1. *hirsutum*, subfasciculare hirsutum nigrum.
Nees System. t. 17. f. 157.

Clavaria ophioglossoides. Holmsk. Coryph. p. 18. ic. Schmiedel Ic. et Analys. pl. p. 92. t. 29. f. 1-19.

Crescit in sylvis, pratis; sed multo rarius est quam sequens, a quo praesertim hirsutiae differt. Clavula nunc compressa, nunc subrotunda plicata.

Obs. An tabula Sowerbei 85. ad hanc an ad sequentem speciem pertinet?

2. *glabrum*, totum glabrum ex spadiceo nigrum, stipite squamuloso basi albicante. *Syn. fung. p. 608.*

α. vulgare, fusco-nigrum, stipite levi, clavula subangusta, passim bifida.

Clavaria ophioglossoides. Bull. champ. 1. p. 196. t. 372. Bolt. fung. 3. t. III. (nimis incrassata picta).

Inter gramina, ad margines sylvarum autumno. Color etiam fusco-niger.

β. paludosum, glabrum nigrum, clavula subtereti, stipite laevi.

Geoglossum glabrum. Stirpes Crypt. Vogenso-Rhenanae. 1. p. 95. (Excl. Synonymo).

In pratis uliginosis, inter *Sphagna*. Magnitudine mediocri.

Ordo secundus.

296

γ. *uliginosum*, tenerius atrum stipite sublongo squamuloso. *Obs. Mycol.* 2. p. 62. t. 3. f. 4.

δ. *macropus*, stipite longo laevi.

Clavaria nigrita Schleicher. *Pl. Cryptog.*

In Helvetiae paludosis? Stip. 2 - 3 unc. longus. Clavul. unc. 1. longa, 2 lin. crassa.

3. *sphagnophilum*, glabrum, clavula olivaceo-atra, stipite tereti sublongo atro - purpureo.

— Ehrenberg. *Sylv. Mycol.* p. 30. †

Prope Berolinum, in Sphagnis, aestate. Proteus, magnum. Forma et magnitudine convenit cum icone Schmied. t. 25. f. 1 - 19. Ehrenb. An bene distincta species?

4. *atro - purpureum*, glabrum atro - purpureum.

Obs. Mycol. 2. p. 62. t. 3. f. 5.

Clavaria atro - purpurea. Batsch. *El. fung.* t. 2. f. 48.

Rarius in graminosis occurrit. Stipes satis crassus fibrilloso - squamulosus. Clavula passim divisa, et nonnunquam ob nimiam humiditatem aut congelatione valde difformis fit.

5. *viride*, gregarium subfasciculatum viride, stipite squamuloso. *Comment. de fung. clav.* p. 40.

G. *viride*, fasciculatum, clavula olivacea, stipite squamoso subaeruginoso, sporidiis fusiformibus: sporidiolis duobus. *Ditmar in Sturm. Fung. germ.* p. 97. t. 48.

Clavaria viridis. *Holmsk. l. c.*

α. aeruginosum. *Mycol. Hefte, I. p. 104.*

Hab. autumnno in sylvis, et hic locus ei est peculiaris. Pulchra ceteroquin species, magnitudine mediocris.

Obs. Geogl. viride *β. atro-virens.* *Mycol. Hefte l. c.* forsan ut species distinguenda, cum in pratis humidis crescat et stipes squamulis denudatus sit.

6. *olivaceum*, sparsum ex olivaceo subfuligineum clavula acuminata stipite sublongiore

Obs. Mycol. I. p. 46. t. 5. f. 7. Mycol. Hefte, l. c. p. 103.

Hab. in graminosis, pascuis, vel in pratis muscosis. Stipes glaber. Pileus compressus, subplicatus. Stipes flavo-fuscus, glaber, basi albicans. Color fungi junioris dilute fuscus, senioris atro-virens fere. *Mycol. Hefte p. 104.*

7. *difforme*, nigrum viscosum, clavula nitente compressa demum curvata difformi. *Fries*

Obs. Mycol. I. p. 159. †

Hab. in Sueciae pinetis, locis graminosis. Clavula junior subcanaliculata. An satis diversa a sequente?

8. *glutinosum*, subfasciculatum nigricans, clavula subelliptica, stipite viscosa. *Syn. fung. p. 609.*

In montosis graminosis apricis, rarius in sylvis.

β. lubricum, totum viscosum, clavula elliptica nigra, stipite atro-brunneo.

Hab. in Vogesia. Clavula etiam est viscosa.

9. *viscosum*, subparvum angustum viscosum nigrum, clavula tereti confluenta. — *Syn. fung. l. c.*

Hab. in pratis montosis et uliginosis in Germania et Suecia. A reliquis speciebus habitu valde differt, cum clavula a stipite vix distinguatur.

†† *HELVELLAE propriae*, pileo (mitra) distincto; submembranaceo a stipite separato.

LI. SPATULARIA. *Pileus* membranaceus, clavulaeformis, compressus, latere utrinque in stipitem decurvans. (*Sporae* elastice prosiliunt).

1. *flavida*, flavo-pallens. *Comment. de fung. clav. p. 34-36.*

* *S. flavida*, albido-lutescens, clava oblonga obtusa. *Nees Syst. 2. p. 44. t. 17. f. 156.*

Clavaria spathulata. Flor. Dan. t. 658.

Helvella spathulata. Sowerb. fung. t. 35. (optima).

Nascitur in sylvis, equidem non communis fungus est, et, uti videtur, solum in regionibus Europae septentrionalibus occurrit. Plerumque in seriem oblongam, vel circularem aggregatus, recens albidus aut aquose pallidus,

fragilis, et sporulas, uti sequentia genera, vi elastica emittit.

** *rufa*, clava obovata, margine undulata. *Nees l. c. f. 13. Schmied. ic. et Anal. pl. 50. f. 1.*

Prope Erlangam a b. Schmiedelio reperta. Pennaeformis rufa, magis compressa. Saltem insignis varietas.

LII. LEOTIA. *Pileus* ovatus aut orbicularis, margine libero stipitem ambiens.

* *Carnosae*, colore plerumque flavescentes aut rubicundae.

1. *circinans*, carnosae (majusculae) rufo-cinnamomeae, pileo orbiculari convexo. — *Icon. et Descr. fung. p. 16. t. 5. f. 5-7.*

Hab. in Germania et Helvetia in abiegnis, ad terram saepius in circulum crescens. Pileus juniorum subviscosus, pallide lividus; 3-4 lin. latus, margine subundulatus. Stipes fuliginæus, basi subnigrescens.

* *gracilis*, minor, pileo orbiculari rufo-cinnamomeo, stipite longo subpubescente aut farinaceo, dilute fuliginæo. †

Ad ramulos exsiccatos crescens prope Neocomium reperta a D. Chaillet.

Obs. Antecedente ultra dimidium minor, et stipitem multo longiorem et vix lin. 1. crassum habet. Color convenit. Nisi species, va-

rietas tamen memorabilis sine dubio est, antecedentem cum sequente connectens.

2. *Mitrula*, gregaria minuta, pileo ovato cinnamomeo. *Icon. pictae fung. Fasc. 4. p. 54. t. 22. f. 5.*

β. *Leotia pusilla*, cinnamomea, stipite gracili glabro, clavula ovata obscuriore. *Nees Syst. p. 44. t. 17. f. 160. Conspect. fung. var. γ. p. 295.*

Elvela cucullata. Batsch El. fung. p. 189. f. 132.

Mitrula Heyderi. Disp. meth. fung. p. 36. t. 3. f. 12.

In foliis aridis Pini Abietis. Sero autumn. Hujus generis forte minutissima, nonnullas lin. tantum alta.

γ. *strobilina*, submajor, capitulo subrotundo pallido unicolore. *Consp. fung. p. 296.*

Haec strobilo vetusto Pini *sylvestris* innascitur.

3. *truncorum*, gregaria carnosā parva, pileo hemisphaerico aureo, stipite albo squamuloso: squamulis glauco-nigricantibus. *Conspect. fung. p. 297. t. 3. f. 2.*

Cresc. in Lusatio superiore, rarius supra truncos lignaque madentia. Majo — Junio.

Gregaria, cespitosa, carnosā, subtenera. Pileolus 2 lin. latus, 1 altus, orbicularis, subtus flavidus vel albidus; exsiccatus aurantioru-

ber. Stipes subtenuis satis firmus, glaucescens, arefactus nigrescit 4 lin. altus. *Consp. l. c.*

4. *clavus*, sparsa, pileo majusculo haemisphaerico flavo-rubro, stipite fuligineo-virescente (subolivaceo), deorsum attenuato. *Tab.*

XI. Fig. XI - 12.

Hab. ad ligna pinea, in rivulis, in Vogesia. *Mougeot.*

Substantia firmula. Pileolus semiglobosus, pro ratione stipitis satis magnus. Stipes laevis vix squamulis obtectus, exsiccando nigricat.

Obs. *Helotium nigripes*, sparsum minutum, pil. primo plano, demum convexiusculo, margine subrepando, hymenioque convexiusculo flavo, stipite brevi cylindrico, apice subdilato fuligineo-nigricante. *Schum. Saell. 2. p. 412.* (ad asseres putridas, Novembr.) nostro fungo affinis videtur.

5. *uliginosa*, gregaria flavo-cinnamomea, pileoli margine cum stipite connato.

α *Bulliardii*, pileo ovato aut subrotundo. *Syn. fung. p. 612.*

Clavaria Phalloides. *Bull. p. 214. t. 463. f. 3.*

Clavaria epiphylla, clavata integerrima, capite obtuso cavo rubro. *Dicks. Pl. Crypt. 3. p. 32. t. 9. f. 10. Sowerb. fung. t. 293.*

β. *Ludwigii*, pileo subconico plicato elongato obscure flavescente, stipite deorsum nigrescente (?). *Syn. fung. p. 611. t. 3. f. 10.*

Hab. sero autumnio in paludosis, ad stipites et folia putrescentia saepius aquis pigris in-natantia. Majuscula unc. 1-2 longa, lin. 2-3 crassa. Affinitatem quandam prae se fert cum *Spathularia*. Var. β . in ipso loco natali de-nuo observari meretur.

* *aurantiaca*, pileo exiguo convexo aurantiaco carnosio-gelatinoso, stipite albo-rubescente fistuloso. *Cumino Fung. Vall. Pisii Specim. in Act. Acad. Scient. Taurin. 1805. p. 221.* sub *Helvella*.

Supra truncos et ramos putridos in fonti-bus aut rivulis degentes vere et aestate. Pileus magnitudine lentis, vel duplo major, subtus stipiti concolor. *Cumino*.

** *CUCULLARIA*. *Tremellosae aut carnosio-gela-tinosae, terrestres, colore obscuri, fuscescen-tes olivaceae aut virescentes, Pileo brevi sub-patulo.*

6. *lubrica*, flavo-virens tremellosa, pileo con-vexo-plano, margine undulato, stipite cras-siusculo cylindrico. *Tab. IX. fig. 4-7.* (var-ietates).

Clavaria tremula. Holmsk. Coryph. 1. p. 27.

Phallus lubricus. Flor. Dan. t. 719.

Hygromitra tremula. Nees System. t. 15. f. 144.

Helvella gelatinosa. Sowerb. t. 70. (Var. duplo major, subrufescens, stipite longo).

Aestate et autumnno non infrequens in sylvis, aut in graniosis ad latera sylvarum.

7. *marcida*, flavo-virens subtremellosa, pileo planiusculo repando, stipite deorsum attenuato. *Comment. de fung. Clavaef. p. 32. Syn. fung. l. c.*

Phallus marcidus. *Flor. Dan. t. 654. f. 1.*

Hab. in Norwegia. Descriptio adhuc desideratur.

8. *atro-virens*, gregaria parva atro-virens, pileolo orbiculari patulo, stipite aequali crassiusculo. *Tab. IX. fig. 1 - 3.*

Hab. in Vogesia in sylvis ad vias, locis humidis. *Mougeot. Subcarnosa.*

9. *platypoda*, minima tremellosa subtenax, pileolo fusco plicato-undulato, stipite compresso sordide albo. †

Helvella platypoda. Decand. Flor. Gall. Suppl. p. 29.

Hab. ad terram prope Grasse in Provincia. Sex lin. solummodo alta. Pil. irregulariter plicatus seu undulatus, margine subreflexus. *Dec. l. c.*

LIII. VERPA. *Pileus* submembranaceus, aut carnosus-membranaceus, a stipite distans. *Hy-menium* rugulosum, (non areolatum) vel laeve.

1. *digitaliformis*, gregaria, pileo campanulato subcylindrico, digitaliformi umbrino, stipi-

te transversim squamuloso. *Tab. VII. fig.*

1-3.

Hab. tempore vernali in sylvis prope Neocomium (Neufchatel in Helvetia). *Chaillet*, qui icones hujus speciei benevole communicavit.

Pileus, qui interdum in fungo valde adulto aut luxuriante, etiam hemisphaericus est, 3-4 lin. altus, totidemque fere latus, aqua emollitus, sub lente reticulato-rugulosus, sed non alveolatus, in sicco subasperatus, inferne laevis, pallidus aut subalutaceus. Stipes $1\frac{1}{2}$ - 2 unc. longus, 3 lin. crassus, cylindricus, intus cavus, flaccidus, transversim undulatus.

2. *Agaricoides*, subsolitaria, pileo campanulato subplicato fuscescente (dilute badio), stipite laevissimo pallido. *Tab. VII. fig. 4-5.*

Morchella agaricoides. *Decand. Syn. pl. Gall. 2. p. 213.*

Crescit passim, vere, in sylva Bolonien-si prope Parisios.

Pileus $\frac{1}{2}$ unc. altus et latus, demum corrugatus. Stipes ad 3 unc. longus, 2 lin. crassus, fistulosus, in adultis deorsum communiter incrassatus, pallidus.

3. *candida* (leucopus), pileo nigro, stipite albo. *Swartz in N. Act. suecicis. 1815. t. 4. Liljeblad Flor. suecica, edit. 3ta. Upsal. 1816.*

Hab. in Sueciae muscosis. Pileus campanulatus, margine revolutus, pallidiore. Stipes inflatus, fistulosus.

4. *conica*, pileo campanulato acuto, margine subsinuato fusco, inferne stipiteque flavo. †

Phallus conicus. *Flor. Dan. t. 654.*

Helvella Relhani. *Sowerby. Fung. t. 11.?*

Hab. in Dania, Germania et Anglia? Stipes cylindricus. Pileus sublaevis. Magnitudine antecedentium.

LIV. MORCHELLA. Pileus carnosus-membraceus campanulatus aut ovatus (passim teres) alveolis forma regularibus lacunosus.

* *Pileo basi a stipite libero s. patulo.*

1. *Gigas*, magna, pileo patulo, areolis oblongis rhomboidalibus, stipite squamuloso deorsum incrassato. — *Michel. N. Gen. pl. p. 202. t. 84. f. 1.*

Phallus Gigas. *Gmel. Syst. vegetab. 2. p. 1443. Ventinat. in Act. Instit. Gallici. 1. p. 508.*

Hab. in Italia, vere, in locis arenosis, etiam circa fossas. *Mich.*

Ex maximis. Stipes conicus, subcrassus, leucophaeus. Pil. 4 unc. altus, a stipite longe distat.

2. *undosa*, submagna, pileo conico, areolis obliquis amplis undulato-crispis, stipite reticulato. *Gmel. l. c. sub Phallo.*

Phallus crispus. *Ventin. l. c. p. 512.*

Phallo-Boletus exculentus, pileo ampliore, rufescente et crispo, pediculo leucophaeo fistuloso striis reticulato insignito. *Micheli p. 203, n. 2. t. 84. f. 2.*

Varius in locis crescit ad rudera, in Italia. Formam parum insoletam prae se fert. Stipes etiam alveolatus apparet.

3. *Rete*, subparva, pileo ovato obscure fulvo, areolis regularibus rhomboidalibus, stipite sublongo albo - pallido glabro. *Gmel. l. c. p. 1449.*

Phallo - Boletus pileolo parvo conico ex fulvo subobscurum, pediculo leucophaeo. *Mich. p. 203. no. 3. t. 84. f. 3.*

In Italiae populetis aut juxta fossas prope Mezzana. *M.*

Magnitudine media et forma quadam grata est. Stipes inverse clavatus. Pil. a stipite longe distat.

4. *fusca*, pileo subrotundo obscure fusco, areolis costis majoribus rectis subparallelis, stipite longo fuligineo-albido.

Unica tantum vice ad asseres, initio Aprilis, prope Parisios hunc fungum reperi, cujus pileus brevis est, fere membranaceus spadiceus. Stipes 3 unc. altus, 1 unc. crassus, laevis, subflaccidus.

5. *crassipes*, pileo brevi acute conico flavescente, areolis costis primariis parallelis, stipite longo crasso albido. †

Helvella hybrida. *Sowerb. fung. t. 238.*

Hab. in Anglia. Stipes, secundum iconem, 4 unc. longus, cylindricus, laevis, 1 unc. crassus. Pil. vix unc. $\frac{1}{2}$ altus et totidem fere crassus.

6. *mesomorpha*, flavescens, pileo campanulato basi a stipite longe distante, stipite subcrasso cylindrico, areolis rhomboidalibus. †

Morchella patula. *Syn. fung. p. 619.* (non *Gmelini*).

Helvella esculenta. *Sowerb. t. 51. fig. media*).

Hab. in Anglia. Magnitudine media. Pil. valde regularis, tenuis. Areolis conformibus. — Extra Angliam forsitan nondum detecta; saltem specimina, mihi sub hoc nomine missa, et a nonnullis autoribus germanicis ita descripta, ab *esculenta* s. vulgari specie, quo etiam *Phallus patulus* Gmel. et *Morchella patula* Grattinnick? referendus omnino videtur, haud diversa fuerint.

** *Pileo basi stipitem arcte ambiente.*

7. *esculenta* (vulgaris), pileo subrotundo seu ovato, basi contracto albido - pallido, areolis excavatis subrotundis. — *Synops. fung. p. 618.*

Phallus esculentus. Linn. Schaeff. t. 199.
Bull. t. 218. f. a - d. Sowerb. t. 51. (fig. sinistra).
Bolton. t. 91.

β. grisea, griseo-fuliginea subnigrescens, areolis subpolygoniis. — Sowerb. t. 51. fig. dextrae? Mich. t. 85. f. 2.? Batarra Fung. Arimin. t. 2. fig. f.

Utilis haec species in pluribus regionibus vulgaris est, praesertim *α*, cui stipes albus et basi subdilatus, et areolae subflexuosae, profunde excavatae sunt; an *β*. forte species distincta? areolae saltem in fig. *Sowerbeana* fere quadrangulares pictae.

Obs. Stipes in omnibus hujus generis speciebus, quae fungi *vernales* sunt, cavus, nec *fartus* quomodo hoc in *Synopsi* minus recte est adnotatum per inspectionem nimirum *fig. cl. Sowerby*, ubi cavitas, ratione aliarum specierum, vix perspicuus apparet.

* *pubescens*, minor pallida, pileo subovato, areolis magnis patulis, stipite pubescente.

Obs. Prope Neufchatel, in Helvetia, in sylvis Juratensibus communis, atque cum prior in usum oeconomicum colligitur. Stipes, in specimine a D. Chaillet transmissio, multo tenuior, vix 3 lin. crassus et pubescens, et areolae multo majores sunt quam in vulgari.

γ. conica, pileo tereti-conico sublongo umbrino, areolis costato-rhomboideis, stipite bre-

re albedo. *Traité sur les champ. comestibl. p. 257. Schaeff. t. 199. f. 1-3.*

Morchella contigua. Trattin. Fung. austr. t. 6. f. 11.

Hab. in Germania et Gallia.

Pileus unc. $1\frac{1}{2}$ altus, sed vix $\frac{1}{2}$ unc. crassus. Areolae costas seu plicas efficiunt longitudinales subparallelas, rugis transversalibus connexas. Stipes subscaber.

β . *metheformis*, pileo conico umbrino, areolis perpendicularibus costato-reticulatis subrhomboideis.

Morchella costata. Schmidt et Kunze, Fung. german. exsicc. Fasc. 8. no. 195. (exclusis synonymis).

Boletus esculentus in metam fastigiatus. Tournef. et Mich. Gen. pl. p. 203.

Phallus patulus Schrank et Gmelin? Conf. Commentarium in Schaeff. p. 81.

Obs. A sequente specie, qui duplo major et colore diversa est, omnino diversa videtur, licet forma areolarum non valde recedat. Sed stipes etiam subtenuis est et brevis, superficie farinaceus.

9. *costata*, magna, pileo longissimo conico, areolis costis rectis subparallelis anastomosantibus, stipite crasso cylindrico.

Phallus Anastomosis. Gmel. Syst. 2. p. 1449.

Bole-

Boletus esculentus compressus, dense foraminulatus et per longitudinem costatus, e cinereo virens, pediculo crassiore albo. *Mich. N. Gen. p. 203 no. 5. t. 35. f. 3.*

Provenit in Italiae agris. Spithamam fere longa et plus quam 2 unc. crassa. Stipes crassitudine ipsius pilei, albus.

10. *rimosipes*, pileo conico basi cubcontracto, stipite elongato rimoso. *Decand. Syn. pl. Gall. p. 43. et Flor. Gall. 2. p. 214.*

Hab. in sylvis circa Fontainebleau.

Spitham. fere alta. Stipes crassus et praesertim versus basin fissuris longitudinalibus, fere ut in *Helvella Mitra*, distinctus. *Decand.*

† *Species sequentes dubiae, vix Morchellae.*

* *crassipes*, stipite inferne dilatato, superne attenuato, pileo (fusco) brevi celluloso acuminato. *Ventenat, Mémoire de l'Institut. 1. p. 509. f. 2. Decand. l. c. p. 213.*

Hab., cum sequente, in Gallia, circa Pontchartrain.

* *tremelloides*, stipite brevi crassissimo, pileo (flavescente) celluloso lobato undulato. *Vent. l. c. f. 1. sub Phallo, uti antecedens.*

LV. *HELVELLA. Pileus* (stipitatus) membranaceus, subinflatus, deflexus, laevis, subtus farinaceus.

* *Stipite longitudinaliter sulcato.*

1. *Mitra* α *nigrescens*, pileo utrinque deflexo adnato, nigrescente livido, stipite farto albicante. *Schaeff. Fung. t. 154. Nees Fung. f. 163.*

Helvella lacunosa. Holmsk. Fung. Dan.

2. *p. 45. t. 24. Fries Obs. Mycol. 2. p. 301.*

β . *cinerascens*, minor, pileo pallide cinereo.

Helvella Mitra. Bull. champ. t. 190.

- γ . *albicans*, tota albida submajor (β), stipite incrassato, (pil. adnato).

Autumno in sylvis satis communis, praesertim in solo arenoso; interdum quoque ad truncos marcidos crescit. Magnitudine et colore pro loco et aëris conditione variat. Stipes vero profunde sulcatus et lacunosus, et pileus utrinque adnatus, huic speciei character est proprius, quae sub diversis nominibus adhuc descripta occurrit. *Helvella sulcata* Willd. et Afzel. non differt, et *H. Monacella* Schaeff. et nonnullorum botanicorum, junior aut non bene evolutus fungus est, stipite magis cylindrico.

2. *leucophaea*, pileo utrinque deflexo libero albo-pallescente, stipite sulcato-lacunoso candido. *Syn. fung. p. 616. Schaeff. t. 282. Michel. Gen. pl. t. 86. f. 7.*

Helvella Mitra. Sowerb. fung. t. 39.

Boletus leucophaeus. Batarra Fung. A-rim. p. 25. t. 3.

Cum priore, a qua pileo non stipiti adnato differt. Pileus superne albo-livescens, exsiccatus dilute rufescens, subtus dilute fuscescens, aliquando subcrispus. Stipes crassus, farctus. Haec species, uti, verosimile, reliquae hujus generis, edulis est.

3. *chrysophaea*, pileo expanso undulato irregulariter lobato fulvo-fusco, stipite albo ad medium sulcato.

Helvella reflexa. Cumino Fung. vallis Pissii, in Act. Acad. Reg. Taurin. 1805. p. 250. t. 2.

Hab. in montibus sub Fagis. Stipes basi attenuatus tubulosus, medium usque sulcatus. Pileus reflexus (uti fere in omnibus). *Cumino.*

4. *erythrophaea*, pileo stipiti adnato irregulariter lobato, stipite subsulcato (sulcis 1-2) dilute rubescente.

Helvella inflata, pil. stipiti adnato inflato fusco-rubro irregulariter lobato, stipite dilute rubescente fistuloso basi dilatato, 1-2 sulcato. *Cumino l. c. t. 3.*

Nascitur post pluvias circa areolas carbonarias in sylvis fagineis. C.

** *Stipite laevi, aut rariter lacunoso.*

5. *grandis*, pileo amplo 3-4-lobato fusco-pustulato, stipite albo laevi aut rariter sul-

cato (lacunoso?). *Cumino*, l. c. p. 250. t.

2.

"Primo vere, post pluvias in montosis sylvaticis crescit. Fungus magnus, nunc violaceus, nunc cinnamomeus, quandoque stramineus, aut fuscus, subtus albus. Stipes cavus. Edulis est, pretiosior *Boleto suillo*, aequiparatur *Phallo esculento*. Lin. et ejus loco venundatur sub nomine vulgari *Pungolo*." C.

6. *Infula*, magna, pileo inflato sublobato badio, stipite subtomentoso albo - rufescente. *Schaeff. t. 159.*

Provenit ad truncos, locis praesertim montosis et subalpinis, autumnno. Ex magnis. Stipes laevis aut sublacunosus, pallens, etiam subgriseus, farinaceus. Pileus subobliquus, stipiti adnatus, 2-3 unc. magnus, brunneus.

Obs. H. *Infula* β . *terrestris* in *Conspect. fung.* p. 299. ad sequentem referri videtur.

7. *esculenta*, gregaria, pileo vario subdifformi plicato-gyroso badio, stipite subbrevis albido rufescente. *Schaeff. t. 160. Flor. Dan. t. 116. Conspect. Fung. Lusat. p. 299.*

Helvelle comestible. *Traité sur les champignons comest. p. 260. t. 4.*

Nascitur iisdem in locis cum antecedente, sed, uti *Morchellae*, tempore vernali. Pileus adnatus, nunc depressus, suborbicularis, patu-

lus; nunc deflexus, emarginatus, sublobatus, et tunc minus plicatus. Plicae gyrosae, subintestiniformes. Stipes cavus, brevis, glaber, incarnato-dilute rufescens.

Obs. Frequens venum fertur fungus edulis: *gemeine Morchel, Stumpfinorchel.* *Consp. l. c.*

8. *leucopus*, pileo utrinque deflexo adnato, demum spadiceo, stipite glabro albo laevi.

Novam hanc speciem inveni, Aprilis initio ad aggeres sub arboribus, prope Parisios.

Habitu *Helvellae Mitrae*, paullo tamen minor. Pileus cum sub terra adhuc latet, est plumbeo-lividus, fungi adulti badius, exsiccati fere nigrescit. Stipes vix unc. altus, 3-4 lin. crassus, cylindricus, versus basin plerumque magna lacuna pertusa distinctus.

9. *albida*, subgregaria, albido-pallens, pileo liber subinflato subparvo, stipite longo farinoso. *Obs. Mycol. I. p. 71.*

Helvella fuliginosa, *Schaeff. t. 220. Sowby. t. 154.*

Helvella elastica. *Bull. t. 242. f. a — g.*

Jam ineunte autumnio in sylvis frondosis sese offert.

Etiam ex Calabriae ulterioris nemoribus resinosis eam habeo; stipite paullo crassiore et breviori subpubescente quasi diversa et quae

ibi nota est sub nomine vernaculo *Tripa di pecore*. Stipes longus, sed tenuis, deorsum crassior, cavus, farinaceo-pubescent, basi saepius lacunosus. Fungus recens albidus, pallens, demum subrufescit, passim subfuliginus fit.

Obs. Sequentes duae *Helvellae* ni fallor vix ut species, ab hac diversae, aestumari possunt.

* *pulla*, subgregaria cinereo-fuliginea, pileo libero bilobato: lobis semilunaribus deflexis, stipite breviusculo subundulato, sursum attenuato fistuloso. *Schumach. Saell. p. 411. Holmsk. Fung. Dan. 2. p. 49. t. 26.* (Figurae priores non differunt).

* *sublicia*, pileo convexo subplicato deflexo, stipite tereti subulato, basi subsulcato fistuloso. *Viborg - Holmsk. Fung. Dan. p. 57. t. 27.*

10. *nigricans*, parvula, e fuligineo nigricans, pileo libero utrinque compresso, stipite farinaceo cinereo. *Obs. Mycol. 1. p. 72. Michel. Gen. t. 86. f. g.* (vix differt).

Helvella atra. Afzel. in Act. Suec. 1783. p. 298. Flor. Dan. t. 504. f. 1.

Hab. jam mature in sylvis, locis umbrosis subhumidis, tamen minus commune. Pil. 3-4 lin. plerumque magnus, rarius $\frac{1}{2}$ unc. Stipes sat longus $1\frac{1}{2}$, hinc inde lacunosus, quasi pubescens, hinc inde lacunosus.

β? *Helvella atra*, pileo bilobato plicato deflexo libero, supra (?) hirto, inferne glabro, stipite basi subsulcato (lacunoso?) hirto solido. *Viborg in Holmsk. Fung. Dan. 2. p. 471. t. 25.*

*** *Species accuratius determinandae.*

* *flavo-virens*, mitra (pileus) subrotunda plana lacunosa, basi compressa flavo-olivacea.

Nees Fung. 2. p. 45. f. 162.

Cespitose in terra, locis subudis umbrosis crescit. Altitudo uncialis. Mitra 4-5 lin. aequat, basi undato-compressa, flavo-olivacea. *Martius Flor. Erlang. p. 402.*

Obs. Cum hi auctores nihil de fungi substantia commemorant, incertum manet, an idem ac *Leotia lubrica* sit, cujus habitum prae se fert.

* *fistulosa (incarnata)*, pileo libero planiusculo repando demum deflexo, supra livido-fuligineo, subtus incarnato-rosello, stipite longiusculo laevi subundulato fistuloso sursum incrassato carneo. *Conspect. Fung. Lusat. p. 299.*

In sylvaticis udis muscosis, Octobri.

"Ad *Pezizas* forsan translocanda. Pil. subdifformis, primo planiusculus, subumbilicatus, dein deflexus, uncialis fere. Stipes debilis, 2-3 unc. longus. Substantia tenera fragilis." *Conspect. l. c.* Affinis *H. albidae*?

* *quineloba*, pileo nigro regulari quinquelobo: lobis deflexis, stipite subcrasso albo.

Boletus albus, pileolo complicato nigro.
Batar. Fung. Arimin. p. 24. t. 2. f. h.

Pileus, secundum iconem, in lobos regulares partitur.

* *conformis*, pileo hemisphaerico regulari fulvo, stipite albo cylindrico.

Boletus pileolo Monachi. Batar. p. 25. t. 3. D.

Autumno ad ripas crescit. Stipes cavus.
Pileus membranaceus. *Bat.* Sensu systematico potius *Verpae* species esset.

LVI. RHIZINA. *Pileus* (acaulis) effusus, subbullatus, immarginatus, subtus radiculis fibrillosis sparsis.

1. *undulata*, effusa undulata, subtus pallida.

Fries Obs. Mycol. I. p. 161. Schaeff. fung. t. 153.

Helvella acaulis, undulata effusa spadicea, subtus fibrillosa pallida. *Syn. fung. p. 615.*

"Autumno humido in pinetis (etiam in sylvis frondosis): nunc loca adusta amat, nunc muscis pulvinatis, et graminibus, pileum passim perforantibus, superfusus occurrit hic fungus, substantia rigida, fragilis. Fibrae radicales crassiusculae, subsimplices albae, saepe copiosae." *Conspect. fung. p. 298.*

2. *laevigata*, convexa laeviuscula (orbicularis),
subtus pallida. *Fries l. c. p. 162.*

Peziza rhizophora, hemisphaerica convexa
fusca inferne albida concava, radicibus simpli-
cibus teretibus. *Reichard in Act. Nat. curios.*
Berol. 3. p. 214. t. 4. f. 4 - 6. Willd. Prodr.
Berol.

Octospora rhizophora. Hedw. Descr.
Musc. 2. p. 15. t. 5 f. 2.

Hab. rarius in sylvis. Priori valde affinis
videtur. Circa Berolinum et in Suecia inventa.

* *praetexta*, carnosa suborbicularis subconvexa
fusco - purpurea, limbo albo margine fibril-
loso praetexta, subtus flavescens, fulcris con-
coloribus. *Ehrb. Sylvae Mycol. (Diss.) p. 29.*

Prope Berolinum in arenosis ericetosis ubi
pauca specimina lecta ab auctore citato.

††† *Pezizoideae*, plerumque parvulae, minus sti-
pitatae, concavo-hemisphaericae, substantia
sat variae.

LVII. PEZIZA. *Pileus* (cupula) hemisphae-
ricus (plerumque), plerumque excavatus, sub-
stantia varius. Thecae in hymenium (saepius
discolor) aggregatae, sporulis 6-8 farctae.

* *Elvela* veteriorum (*Helvelloideae* Syn. fung.
Pruinatae Nees. *Aleuria* Fries). *Majores*,
ceraceae fragiles plerumque, glabrae, ex-
terne squamulis farinaceis adpersae.

* *Stipitibus crassis, sulcis longitudinalibus exaratis et sub cupulis ramificatis.*

1. *Acetabulum*, magna infundibuliformis fuliginosa, stipitis longi sulcis in cupula costato-ramosis. — *Vaill. Bat. Paris. p. 57. t. 13. f. 1. Bull. champ. p. . t. 485. f. 4. Se-werb. fung. t. 59.*

Hab. rarius in umbrosis. Vaillant omnium primus uti videtur, detexit hanc speciem Versalius, medio Aprilis. Etiam in Anglia crescit, et in Germania *Gleditsch* (Method. fung. p. 42.) sed diversas pto more commiscuit species. Stipes albus.

2. *contorta*, submagna, cupula imbricatim contorta stipite crasso simpliciter sulcato.

Boletus pileolo spiralibus plicis contorto. Batarra Fung. Arimin. hist. p. 24. t. 2. f. 9. Michel. N. Gen. pl. p. 204. no. 4.?

In Italiae nemoribus humidis provenit. Pedunculus albus, cavus et quasi ex funiculis compositus. Pileolus rufus et in plicas, in modum contortus. Substantia veluti cerea. *Batar.*

3. *caelata*, cupula hypocrateriformi fulva, stipite albo: costis subconnatis apice breviter ramosis. *Batar. l. c. p. 25. t. 3. c.*

Supra folia quercea putrida. Autumnalis. — Pedunculus albus, cavus, striatus. Pileolus calyciformis, fulvus. *Bat.*

4. *sulcata*, ex albido - pallens, minor, stipite sulcato, cupula subtus laevi. *Syn. fung. p. 643. t. 5. f. 1. Consp. fung. p. 313.?*

Cresc., vere, in sylvis fagineis, locis apricis. Stipes unc. 1. longus, 4 lin. crassus. Cupula hypocrateriformis, lin. 1 alta, sed 4 - 6 lin. lata.

5. *leucomelas*, solitaria minor tota alba, stipite crasso interrupte sulcato, cupulae disco nigrescente. †

Hab. in Gallia. Stip. terrae ultra dimidium infossus videtur, et ejus costae non usque ad cupulam adscendunt. Antecedenti ceterum similis, cujus an varietas: disco per exsiccationem aut aliam rationem colore mutato.

6. *ancilis*, submagna griseo - brunnea integra, stipite brevissimo apice costato - diviso: ramulis crassiusculis cum cupula coalitis.

- β. *clypeata*, parvula subplana nigrescens, stipite sub cupula subquadrifido.

Hisce in sylvis ad terram, Majo, inveni hanc speciem sat memorabilem.

Stipes subtuberosus, sub terra fere absconditus, nonnullas lineas longus, sub cupula ad medium in ramos sex praecipuos dividitur: ramis divergentibus tumidis, medio canaliculatis. Inter unamquamque divisionem seu costam, lacunae adsunt profundae et latae, plicato - rugo-

sae, albae. Cupula 2-3 unc. lata, valde fragilis, externe farinosa, basi albicans. Sequenti proxima: β . forte junior fungus, vix tres lin. latus.

7. *venosa*, subcontorta sessilis majuscula, superne umbrina, inferne alba, venis costato-ramosis. *Syn. fung. p. 618.* †

Elvela *cochleata*, coriacea acaulis hemisphaerica auriformis subcochleata, supra ex fusco nigra, subtus alba, venis costato-ramosis. *Wulfen in Jacq. Miscel. Austr. p. 112. t. 17. f. 1.*

Hab. in Austria. Substantia ceraceo-cartilaginea. Cupula primum subrotunda hians, sensim explanata, integra aut etiam lobata et veluti crispata, subtus ex centro venae nonnihil extantes. Ultra unc. lata. *Wulf.*

†† *Otidea*, cupula uno latere incisa saepius convoluta aut in formam auricularem producta, sessilis.

1. *umbrina*, magna caespitosa fragilis contorta obscure umbrina. *Syn. fung. p. 638. Bull. t. 154. f. a. Sowerby. t. 5. Schaeff. t. 274.*

Provenit in regionibus praecipue arenosis, tam in sylvis, quam in graminosis, imo ad latera fossarum, aestate et autumnii initio. Magna, fragilis.

2. *geochroa*, magna solitaria alutaceo-cinerea convoluta, interne basi tuberculosa albida, radice crassa horizontali.

In sylvis prope Parisios, sed unica vice hunc fungum reperi.

Cupula $1\frac{1}{2}$ unc. alta, $\frac{3}{4}$ unc. lata, externe squamulosa, deorsum terreo - fuliginea, fragilis, radicem oblongam subtuberosam terra tomentoque involutam exserit. Primo intuitu sequenti similis.

3. *alutacea*, caespitosa contorta, externe alba, intus pallescente - fuliginea, stipite vix ullo albo - tomentoso. *Syn. fung. p. 638. Bull. l. c. t. 154. B.*

P. cochleatae, var. *α*. *Batsch El. fung. cont. 1. p. 223. t. 28. f. 157.* sat bene convenit, sed integerrima depicta.

β. *Elvela ochracea. Schaeff. t. 155.*

Nascitur satis copiose autumnno in sylvis subumbrosis. *P. umbrina* minor, nec ita fragilis.

4. *fuliginea*, gregaria majuscula demum subconvoluta, interne plumbeo - fuliginea subnigrescente, externe grisea. †

Inter muscos mihi obvia, forma antecedentis, sed colore exacte *Helvellae Mitrae*, in cujus societate eam inveni, terrae subimmersa. Primo integra, auriculiformis seu obliqua, demum uno latere incisa subconvoluta, $\frac{1}{2}$ - $\frac{3}{4}$ unc. lata, 4 lin. alta.

5. *concinna*, caespitosa magna fragillina, exter-

ne citrina demum rugulosa, intus incarnato-pallida.

Helvella vesiculosa. *Bolt. fung. t. 175.*

Elvela scutellata *Schaeff. t. 150.* et *Elvela pyxidata*. *Schaeff. t. 236.*

Pulchra haec species mihi sese obtulit sub finem Julii, post pluvias continuas, inter congeriem foliorum quercuum, in sylva Vincennes prope Parisios, plures cespites sibi approximatos sistens. Individua majora latitudine erant palmari. Valde fragilis, exsiccando rugulosa, basi ubi terrae incumbit, albo-villosa. Non confundenda cum *Pez. onotica*.

6. *aurantiaca*, caespitosa contorta fragilis aurantiaca, externe albida. *Obs. Mycol. 2. p. 76.* *Flor. Dan. t. 657. f. 2.* *Schaeff. t. 148.*

Peziza coccinea. *Bull. t. 474.* *Sowerby, t. 78.*

Autumno praesertim in quercetis, in seriem aut in circulum crescit; 2 unc. magna, fragillina, diutius fumans, pulcherrima.

7. *onotica*, magna dimidiato-elongata pallescente-rosea, interne plicato-rugosa. *Syn. fung. p. 617.*

Peziza leporina *Sowerb. fung. t. 72.*

P. onotica, pileo substipitato auriformi buxo roseo, extus basi stipiteque tomentoso eborino (eburneo), sporidiis octo oviformibus;

sporidiolis duobus. *Ditmar in Sturm Fung. Germ. p. 33. t. 16.*

P. rosea, submagna subcespitosa auriformis convoluta, basi coarctata substipitata, intus carneo - rosea, extus pallida. *Schumacher Saell. p. 416.?*

Hab. in sylvis frondosis, sed rare et plerumque solitarie crescit. Cupula firma, oblique aperta, auriformis, passim quoque saciformis, substipitata. Magnitudine, colore, et loco natali praecipue a sequente differt, cum qua saepius confunditur.

8. *leporina*, gregaria cinnamomea dimidiato - elongata auriformis, intus laevis. — *Batsch El. fung. p. 117. Schaeff. t. 156. Flor. Dan. t. 1078.*

Peziza cochleata. Holmsk. Fung. Dan. 2. t. 6. f. e.

In pinetis non infrequens. Substipitata, etiam et subintegra occurrit, fragilis, vix unc. 1 - 1½ alta, basi tubiformis.

9. *felina*, solitaria dimidiato - elongata auriformis cinerea. †

Semel eam his in locis ad latera sylvae Meudon observavi, magnitudine intermedia inter duas antecedentes, ab utraque, colore differt griseo; forma magis ad *P. onoticam* accedit. Ex vivo denuo describenda.

10. *sepulchralis*, gregaria alutaceo-fusca, primo oblonga obtusa, verruccis acutis obsita, dein obconica, disco plano (uno latere) in cupulam dilatatam irregularem transeunte. *Rebent. Prodrom. Flor. Neomarch. p. 384. t. 4 f. 18.*

In caemeterii tumulis, prope Berolinum, mense Febuario singularis haec species detecta fuit ab auctore, qui sequentem, in libro cit. dat descriptionem:

Polymorpha: primo oblonga, obtusa, more *Lycoperdi* verrucis acutis obsita, dein obconica, extus lacunosa, disco plano marginato. Margo sensim expanditur in cupulam irregularem unc. circiter latam. Radiculae (sat multae) fibrosae longae albae.

††† *Cupula integra, sessilis aut stipitata stipite laevi. (Geopyxis).*

11. *lancicula*, acaulis cespitosa, cupulis externe dilute fuscis subolivaceis, ore contracto. *Rebent. l. c. p. 384.*

Hab., autumnno, ad fossarum latera. Cupulae 2 - 4 et ultra latae, 2 lin. altae, intus glabrae (laeves), extus rugosae, ore integro ad unum latus saepius hians. *Reb.*

12. *badia*, subacaulis integra, margine subinvoluta intus badia, externe subolivacea, stipite brevissimo albo-tomentoso. *Obs. Mycol.*

col. 2. p. 78. Syn. p. 639. Fries Obs. Mycol. 1. p. 16.

Peziza cochleata. Bolt. t. 90.

In sylvis, $1\frac{1}{2}$ unc. lata, sublacunosa, basi farinosa. Non confundi debet cum *Pez. umbrina*.

13. *bufonia*, magna spadicea, externe verrucosa, stipite brevi pallescente radicato.

Rarissime in sylvis subhumidis, sero autumno. Saepe fragilis, hinc incertus sum, an cupula semper integra, aut latere incisa sit. Ceterum magnitudine et colore cum *P. umbrina* concordat, verum externe non farinosa, sed verrucis conicis, submollihus ubique obsita. Externe colore opaca, terreo-fuliginea.

14. *granulosa*, solitaria submagna, primo hemisphaerico-concava, demum explanata subrugosa sessilis, intus umbrina, extus dilutior, margine granuloso crenata; *Schumacher* p. 415.

Rarius in faginetis ad terram, in Dania Septembr.

15. *papillosa*, acaulis subglobosa dilute fusca, externe papillis atro-fuscis obsita, margine crenato. *Reichard et Flor Wetteravia* 3. part. 2. p. 374. †

Crescit in Wetteravia in terra humosa hor-
torum, autumno.

16. *plicata*, acaulis majuscula subglobosa griseo-cinerea, externe subfurfuracea, interne livescente, ore subplicato.

Eam reperi, mense Novembri, ad terram in sylva Romainville procul Parisios. Quodatenus formam habet et colorem *Pezizae floccosae* Batsch. *El. fung. Cont. I. p. 223. t. 28. f. 156.*; sed individua juniora, meo in fungo, globosa, nec obovalia erant. Adulta semunciam magna, cerea, fragilis. Sequenti forte nimis affinis.

17. *spurcata*, gregaria majuscula sordide alba, seu subfuliginea, externe farinacea, cupula margine integerrimo.

Hab. in sylvis locis adustis, aestate.

Sessilis, $\frac{1}{2}$ unc. magna, fragilis, grisea, cupulae margo quasi striatus et magis farinaceus, sed non crenatus, uti in sequente, cui haud dissimilis.

18. *pustulata*, sessilis fusca, extus furfuraceo-pustulata. *Syn. fung. p. 646.* — Batsch. *El. fung. p. 223. f. 157. †*

Octospora pustulata. *Hedwig Musc. frond. t. 6.*

In pinetis et ad fossas. Unc. $\frac{1}{2}$ lata.

19. *crenata*, sessilis parvula subsolitaria albidopallens tenera, cupula subventricosa, margine crenato. *Obs. Mycol. 2. p. 79. Bull. t. 396. f. 3.*

Post pluvias aestivales mox apparet in sylvis umbrosis ad terram; magnitudine et forma cupulae glandis quercinae. Primo alba.

20. *ampliata*, solitaria glabra fuligineo-fusca ore amplo campanulata. *Icon. et Descr. fung. 2. p. 31. t. 8. f. 6.*

Hab. ad truncos. Externe fere glabra, carnosa, fragilis. Color internus subcinnamomeus externe pallidus.

21. *micropus*, majuscula pallescente-fuliginea, cupula subobliqua flexuosa, extus furfuraceo-granulosa, stipite brevi subcrasso. *Icon. et Descr. p. 30. t. 8. f. 3.*

In truncis nascitur perraro, aestate. Cupula subinfundibuliformis, latitudine circiter unciali, parum fragilis.

* *gemmata*, subgregaria sessilis sphaerico-depressa, ore repando subinvoluta, sordide pallida, extus granulis umbrinis obsita. *Schumacher. Saell. p. 421.*

Vere in trunco putrido Fagi, in Dania.

22. *lacera*, majuscula subcaespitosa sessilis, cupulis tenuibus (submembranaceis) saepius laceris, externe fusco-terrea (dilute umbrina) dense farinacea, intus albida.

Hab. ad ramos coryleos. Ehrberg. $\frac{1}{2}$ unc. et ultra lata, fragilis, stipite videtur carere;

sed externe uti in quibusdam *Lycoperdis* farina grossa oblecta.

23. *carbonaria*, stipitata subparva ochraceo-rufescens, cupula ventricosa margine farinoso crenato. *Conspect. fung. Lusat. p. 314. t. 4. f. 2.*

Occupat loca adusta sylvatica ibique carbonibus vetustioribus innascens. Statura variabilis, latitudine a 3-6 lin., intus viridius picta, deinde magis dilatata et unicolor, ob substantiam teneram saepe varie incisa. *Consp.*

24. *Marsupium*, subgregaria acaulis albido-glauca dein pallescens subglabra ore connivente flexuosa. *Syn. fung. p. 641.* (Exclus. Synonym.)

Elvela auriformis. Hoff. veg. crypt. 2. p. 10. t. 5. f. 3.?

Aestate post pluvias in sylvis frondosis ad terram nudam, in Germania; valde concava, hinc alta, basi intus tuberculosa, unc. fere magna.

25. *vesiculosa*, gregaria sessilis magna, primo globosa ore connivente dein campanulata pallide fuscescens, demum subincisa, ad basin albo-tomentosa. — *Sowerb. fung. t. 4.*

Peziza cerea et P. vesiculosa Bull. hist. 1. p. 270. t. 44. et 457. f. 1.

Oritur fine hyemis, in hortis ad fimeta

vetusta equina, dense plerumque caespitosa, et stipitem quemdam tomento albo involutum intra fumum protrudit.

Obs. Sequentes tres forma et loco nimis affines ut tantum varietates videantur.

β. coriaria, magna subgregaria albo-fuliginea, externe grumoso-granulosa subcaulescens basi albo-tomentosa, margine primo involuta repando-diffusa et fuscescens.

In ipsis hybernaculis plantarum, lignis pulverem coriariarium includentibus increscens, primo vere. Latitudine propemodum est palmari, ob locum forsitan magis faventem, magisque explicata.

γ. isochroa, minor caespitosa sessilis tota alba glabra, cupula ventricosa margine integerrimo.

Provenit quoque in Caldariis sed supra ipsum pulverem coriarium.

Forma gaudet quadam grata, regulari, quae vero in valde adultis deprimitur. Si plura individua sociata sunt, tunc ob mutuam compressionem angulata sunt. Substantia crassiuscula, sed fragilis, latitudine 4 lin. et altitudine 2-3 lin.

δ. turbinata, acaulis caespitosa magna turbinata albo-grisea, ore subcontracto crenato.

Syn. fung. p. 636. Nees System. t. 38. f. 257.

P. coronata, inverse ovata ventricosa albo-grisea margine erecto dentato. *Batsch. El. fung. p. 121.*

Fungoides pyri inversi forma ore lacero et veluti coronata ex albo griseum. *Michel. Gen. p. 207. t. 86. f. 2.*

Cresc. in fimo equino tempore brumali. *Mich.* Haecce distinguitur forma obconica, cujus margo dentibus magnis erectis munitus est.

26. *incisa*, magna gregaria subcyathiformis substipitata, margine laciniato: laciniis erectis et reflexis inaequalibus, intus umbrina, extus dilutior. *Schumach. Saell. 416. P. vesiculosa.*

In fagetis, autumnno, in vicinitate arborum caesarum. *S.*

27. *sphacelata*, solitaria aut gregaria subventricosa fusco-pallida, basi plicata, margine erecto duplicato-crenulato: crenulis sphacelatis, stipite brevi bulboso. *Schum. Saell. p. 419.*

Hab. in Dania ad terram rarius, in graminosis et in vicinitate truncorum. *S.*

Obs. Auctor citavit ut synonymem sequentem fungum, sed hujus figura cum ejus phra-

si non respondet. *P. crenata*, quam etiam allegant, aliis fungus est.

* *Catinus*, substipitata globoso-campanulata, limbo repando, externe pallide ochraceo punctisque fuscis notata, interne fusco-ochraceo Viborg. — *Holmsk. Fung. Dan. 2. p. 22. t. 8.*

Hab. in Dania ad ligna putrefacta.

28. *coronata*, sessilis magna cyathiformis seu campanulata fusca utrinque glabra, limbo laciniato erecto. *Jacq. Misc. Austr. 2. p. 140. t. 10.*

Hab. in Austria, locis umbrosis, ligno putrido adhaerens, 2 - 3 unc. lata, inaequaliter laciniata, intus fusca (badia?) foris pallida.

29. *repanda*, cespitosa magna intus lutescens, externe albicans sinuata, margine repando, stipite brevissimo demum obsoleto. *Ic. pictae fung. rar. Fasc. 4. p. 49. t. 22. f. 3.*

In sylvis rarior, in Germania.

Obs. Affinitate quamdam cum *Pez. alutacea* gaudere videtur, etiam margine interdum incisa sed non ad basin usque fissa nec convoluta.

30. *Crucibulum*, solitaria maxima infundibuliformis sessilis subalbida. *Syn. fung. p. 642.*

Fungoides maximum, vasis fusorii forma,

subalbidum. *Mich. Gen. pl. p. 207. no. 7. t. 86. f. 1.*

In Italiae sylvis olim a Michelio detecta magna haec species, ore lato aperta, omnium facile altissima, ad basin sensim angustior.

31. *Sowerbea*, caespitosa globoso - hemisphaerica, intus flavescens externe subreticulata albida, radice crassa villosa alba. *Syn. fung. p. 642. Sowerb. fung. t. 114. Pez. radiculata.*

Hab. in Anglia ad terram in hortis. Submagna, primo globosa seu ore connivente, dein dilatata, subrepanda, extus subtomentosa, sessilis, sed radicem fusi formam sat magnam, intra terram et stramina absconditam habet.

32. *cerea*, caespitosa substipitata lutescens, cupula repanda subinfundibuliformi extus cum stipite crasso albicante. — *Sowerb. t. 3. †*

Cresc. in areolis corticis coriarii.

Obs. Species magna, habitu quodam peculiari subdifformi distincta, hinc eam, quamvis a me nondum repertam, a *Pez. vesiculosa* uti et a *priore*, cum quibus propagationis locum communem habet, diversam omnino puto. Caul. $\frac{1}{2}$ unc. longus, apice valde dilatata et villosus? Descriptio auctoris, pro more, nimis brevis et vix ulla.

33. *abietina*, gregaria cinnamomea subflexuosa

campanulata, externe demum ruguloso-lacunosa, basi tomentosa. *Syn. fung. p. 637.*
Conspect. fung. p. 309.

In sylvis abietinis. Unc. 1 lata, basi saepe albicans.

β. *grandis*, magna acaulis flexuosa campanulata olivaceo-umbrina (spadicea?), basi lacunosa alboque tomentosa. *Obs. Mycol. i. p. 27.* Var. hujus luxurians potius, nec *Pez. leporinae*.

γ. *rubiginosa*, gregaria majuscula rubiginosa externe dense farinosa, stipite brevissimo subtomentoso.

Hab. in Vogesia, praesertim ad margines viarum sylvaticarum. Unc. lata, oblecta pulvere farinaceo concolore in verrucas acutas, uti in quibusdam lycoperdis, compacta. Forsitan distincta species.

* *porosa*, sessilis orbicularis cinnamomea subrufescens, disco demum foraminulosa. *Syn. fung. p. 640.*

Quoque in Pinetis observatur. *P. abietina* dimidio minor. Hymenium punctato-lacunarum, quod tamen olim in diversis individuis et diverso tempore in Hercynia vidi (etiam hujus Fungi cl. Fries, tanquam Sueciae indigenae mentionem facit) modo fortuito ortum vi-

detur, cum simile quid in aliis speciebus non occurrat.

34. *verrucosa*, magna infundibuliformis (?) pallescente livida, tuberculis inaequalibus obsita, stipite difformi brevissimo. *Syn. fung.* p. 646. †

Olim prope Gottingam, ad truncos inventa, externe, pro more, albicans aut pallidior, et protuberantiis inaequalibus rugosa. Stipes subtuberosus quasi. *l. c.*

35. *Rapula*, gregaria, cupula infundibuliformi, primo subflavescens dein umbrina, stipite sublongo deorsum fibrillis longis vestita. *Syn. fung.* p. 658. *Nees System.* t. 38. f. 297.

Peziza Rapulum, cerea tenuis fragilis glabro, in stipitem longum radiculis instructum desinens, cratera cupulari. *Bull. Hist.* I. p. 265. t. 485. f. 2.

P. radicata, cyathiformis, intus pallide testacea, extus cinereo-gilva, stipite glabro tortuoso, radice simplici villosa. *Viborg-Holmsk. Fung.* 2. p. 24. t. 9.

Humo profunde infixi occurrunt. Stipes tortuosus aliquando fibrillis radicalibus destitutus. *Bull.* — Radix filiformis, vellere fusco undique vestita sub muscis et fol. putrescentibus proserpit. Stipes albidus. *Vib.*

36. *tuberosa*, cupula parvula campanulata, sti-

pite longo abscondito, tuberi magno subdif-
formi nigro innato.

P. tuberosa, capitulo subcampanulato ex-
tus fusco intus pallidiore, stipite basi tubere
nigricante fungosa adnato. *Dicks. Pl. Crypt. 2.*
p. 25. Sowerb. t. 60. Bull. champ. 2. p. 266.
t. 485. f. 3.

Octospora tuberosa. Hedwig. Musc. frond.
2. p. 31. t. 10. f. 2.

Pez. radicata, fusco - spadicea hemisphae-
rica, radice longa simplici. *Reich. in Act. Be-*
rol. 3. t. 4. f. 4 - 6?

Hab. in graminosis sylvaticis, aut in pra-
tis montosis. Vere.

Stipes ad capitulum usque intra terram de-
git. *Dicks.* — Humo profunde infixi reperi-
tur. Ex colore albo, dilute fuligineo aut sub-
fulvo in lateritio - fuscescentem transit. *Bull.*

Cupula $\frac{1}{2}$ unc. lata et alta. Stip. 1 - 2 unc.
longus, tenuis.

Obs. Tuberculum radicale subglobosum
nigrum, intus album, 3 - 4 lin. magnum est
vera hujus speciei pars, nec pro *sclerotio*,
tubere quodam, aut alius plantae perfectioris
radice habenda, nam substantia interna et cor-
tex exterior tuberculi et stipitis hujus fungi,
eadem sunt natura.

37. *Tuba*, stipitata cespitosa magna, cupula in-

fundibuliformi fulva, radice tuberosa subdepressa. *Syn. fung. p. 644. Pez. perennis.*

Peziza Tuba. Batsch. El. fung. p. 121.

Fungoides cespitosum infundibuliforme fulvum, radice nigra tuberosa perenni. Michel. Gen. pl. p. 205. no. 5. t. 86. f. 10.

In Italiae castanearum sylvis, mense Aprilii. Hujus plantae radix perennis est et e tribus plerumque tuberculis constat, quorum medium fungos profert, alterum effoetum, et tertium instanti anno fertile. *M.*

38. *bulbosa*, cupula hemisphaerica cinerascens, intus badia aut pulla, stipite sublongo, radicis tubere subrotundo. — *Nees System. t. 38. f. 289. Hedw. l. c. p. 34. t. 10. f. 3.*

Autumno ad fossas sylvarum. *Hedw. Sequenti subsimilis!*

39. *Macropus*, gregaria magna, cupula hemisphaerica utrinque cinerascens, stipite longissimo attenuato. *Obs. Mycol. 1. p. 26. t. 1. f. 2. Schaeff. t. 167. Flor. Dan. t. 1200. f. 2.*

Peziza stipitata. Bull. champ. 2. p.

Peziza sublicia. Holmsk. Fung. 2. p. 26. t. 10.

Haec species, stipite longo notabilis, non equidem rara est in sylvis inprimis frondosis, et aestate praesertim humida. Stipes nunc lae-

vis, nunc hinc inde lacunosus. Cupulam semper regularem, orbicularem inveni, ab aliquibus convexa est delineata: fungo forsitan nimis maturo, aut in locis humentibus invento.

β. *hirta*, cupula stipiteque hirto vel pubescente. *Obs. Mycol. I. p. 26. t. I. f. 3. Sowerb. fung. t. 38.*

Octospora *villosa*, stipite mediocri elongato patellaque convexo-marginata villosa: villis connexis sursum spectantibus. *Hedw. Musc. frond. 2. p. 54. t. 19. b. ?*

γ. *erosa*, pallescente-cinerea, cupula valde dilatata saepiusque erosa, stipite mediocri laevi subcylindrico albicante.

Elvela hypocrateriformis. *Schaeff. t. 152.* satis congruit.

Cupula multo latior et altior quam in α., et margine, forsitan ab insectis aut limacibus, fere semper lacera. Stipes vero multo brevior, vix unc. 1 longus, basi subtuberosus, saltem tumidus. Rarior in sylvis occurrit.

δ. *tephromelas*, dimidio minor grisea, disco nigro.

Fungoides parvum pyxidatum externe ex obscuro griseum, interne nigrum et pellucidum. *Mich. Gen. pl. p. 205. t. 86. f. 11.*

Crescit vere in Italia, in Scandicci territorio. *Mich.* Verosimilis diversa.

40. *melastoma*, majuscula cespitosa, cupula urceolata externe fuscescente pulvere rubiginosa adpersa, intus nigra, stipite brevi radiculis byssoideis densis nigris.

Peziza melastoma. Sowerb. fung. t. 149.
(Nomen antiquius).

Pez. rhizopus, arcte cespitosa subconnata, cupulis e globoso dein hemisphaericis intus nigris, extus lanato - hirtis (?) rubigine conspersis, capillitio radiceformi undique tectis. *Conspect. fung. p. 317.*

Hab. rarior in Gallia, Anglia et Germania, ad plantarum radices terra denudatas aut ad truncos muscosos arborum.

Fungus sane peculiaris, habitu *Lycoperdi*. — Stipes inter jubam atram, saepe longissimam absconditus, crassiusculus.

Cupula juniorum globoso-obcordata, margine lato inflexo, dein ore patulo ventricosohemisphaerico. *Conspect.*

Obs. Specimina circa Parisios et in Vogesia detecta, nullo tomento, basi excepta, vestita. Sequens non valde diversa videtur.

* *Stenostoma*, gregaria clavato-cylindrica superne (?) villosa aurantio - miniata, cupula concava, margine inflexo ore angusto, stipite nigro fibroso - subradicato. *Martius Flor. p. 469.*

Hab. in ligno putrido. Trés lin. alta, turbinata. Stipes totus niger, floccis concoloribus in ligno radicans. *M.*

41. *melania*, *α. Vogesiaca*, gregaria submagna, cupula campanulata, demum subrepanda, nigrescente umbrina subrugulosa glabra, stipite breviusculo radiculis byssoideis nigris. — *Moug. et Nestl. Stirp. Crypt. Fasc. 6. no. 584.*

Hab. vere in Abiegnis Vogesorum, ad ligna putrescentia, sub terra abscondita, quibus adhaeret radiculis byssoideis, nonnunquam late expansis.

Cupula junior subventricosa, adulta campanulata, margine saepius crenulato-lacera, ultra unc. 1 lata, externe vix pruinosa, sed exsiccatione corrugata est. Stipes brevis, 3 - 4 lin. altus, 2 - 3 crassus, passim lacunosus.

β. *fusco-cana*, stipitata subtruncalis, extus fusco-nigricans laevis, intus incano-olivascens, cupula lata, stipite breviusculo compresso-plicato sublacunoso. *Conspect. fung. p. 312. t. 5. f. 2.*

Incolit sylvas abietinas muscosas, vere. Stipes $\frac{1}{3}$ - $\frac{1}{2}$ unc. longus, una cum cupula basi exteriore rugoso-lacunosus. Cupula margine interdum crenata, intus canescens, aut in olivaceum vergens, 1 - 2 unc. lata. *Conspect.*

δ. *Sphagnophila*, submagna stipitata, cupula campanulata intus atra, extus rugosa subumbrina, stipite subcrasso, fibrillis radicato.

Hab. in Germaniae paludosis. *Ehrenb.* Cupula in specimine transmissio, farinacea et valde plicata, ultra unc. lata. Stip. 2 lin. altus, fibrillis muscos obtegit.

42. *Nigrella*, sessilis hemisphaerica atra, extus deorsum byssina. *Syn. fung. p. 648.*
Conspect. fung. p. 318. Decand. Flor. Gall. 6. p. 2.

Elvela hemisphaerica. *Wulfen in Act. Soc. Nat. Curios. Berol. vol. 8. p. 141.*

Vere et autumnio; Gregaria aut cespitosa, terrestris, rarius truncigena.

Intus atra glaberrima lucida, externe vestita tegmine fusco-nigro crassiusculo, e filiis byssinis intertextis compacto; $\frac{1}{2}$ - 1 unc. lata.
Conspect.

43. *Diadema*, nigra globosa extrinsecus rugoso-costata, vertice late aperto, ore contracto marginato-crenato. *Gmel. Syst. vegetab. 2. p. 1455. Mich. Gen. t. 86. f. 18.*

Hab. quoad Michelium, vere, in Scandici alti sylvis, verosimile ad terram. Species est nitida, magnitudine et habitu *Pezizae crenatae*, etiam radiculis fibrillosis donata.

44. *arctaria*, acutis fusca radicans; junior globoso-campanulata; margine integro, senior dilatata lacera. *Syn. fung. p. 641.*
Act. Suec. 1763. p. 295.

Crescit in Hallandia meridionali (provincia Suecica) in arena mobili, quam figere dicitur radice longissima fibrosa.

45. *fulva*, stipitata magna hypocrateriformis
fulva, inferne alba, stipite incrassato subbrevis. *Micheli Nov. Gen. pl. p. 205. t. 86. f. 6.*

Hab. in Italia per Boboli nemora, Novembri. Unc. 1 lata. Stipes $\frac{1}{2}$ unc. longus.

46. *fulgens*, magna sessilis campanulata integra, superne rubra, inferne alba.

Apparet Majo — Junio, in pinetis prope Neuchatel, colore nitido fulget. Non confundenda cum *P. aurantiaca*, quae contorta et autumnalis est.

47. *ericetorum*, subgregaria minor integra, ut plurimum depressa, intus fulvo-rubra, externe albo-flavescens.

Peziza leucoloma *β. major. Consp. fung. pag. 335?*

Aestate, post imbres, in ericetis passim occurrit. Praesertim explanata, tamen quoque campanulata est, sed non contorta, $\frac{1}{2}$ unc. lata.

48. *purpurascens*, subgregaria sessilis livido-purpurascens, cupula campanulata, ore lato subobliquo. *Tab. XII. fig. 10.*

Rarius ad terram nudam provenit. Magnitudine media, forma grata, passim incisa.

49. *violacea*, subsessilis orbicularis depressa purpureo - caerulea. *Syn. fung. p. 629. Consp. pag. 312.*

Ad terram et interdum in truncis cariosis nascitur. Primo subcampanulata, dein dilatata, margine subinvoluta, latitudine semiuncialis.

50. *depressa*, sessilis orbicularis depressa rufo cinnamomea. *Obs. Mycol. I. p. 40. t. 6. f. 7.*

In agris argillosis, locis humentibus, plerumque solitaria provenit, $\frac{1}{2}$ unc. lata, inferne carnea.

51. *applanata*, gregaria subconfluens orbicularis depressa, primo luteo-ferruginea, demum badio-umbrina, disco tuberculata, ut plurimum sessilis. *Schum. Saelland. p. 428.*

Ad terram in fol. dejectis Pini *Abietis* lecta, Octobri. Lin. 2 — 5 et ultra lata. Nonnunquam obsolete stipitata. Caro umbrina. S.

Obs. Auctor suam ad speciem adduxit: Octospora *applanata*, expansa plana rugosa cinnamomea subtus pallidior. *Hedw. Musc. frond. II. p. 22. t. 5. C.*

52. *hepatica*, atro-purpurea sessilis reflexa, disco saepius scrobiculo notata. *Batsch. El. fung. Cont. I. p. 199. t. 26. f. 138.*

In terra nudiore inter muscos se propagat. *Gregaria subdepressa*, 2 lin. lata. *B.*

53. *Schenkii*, sessilis convexo-depressa patens, utrinque atra nitida, margine brunneo. *Batsch. Elench. fung. p. 125. t. 12. f. 52.*

Prope Jenam in terra muscosam, vallecula umbrosa inventa, mense Aprili; 2-3 lin. lata, regularis, concava, margine integra.

Obs. Notavi hunc fungum, dicit b. Auctor, nomine cl. I. H. C. Schenke, Med. D., qui me, eodem die quo eum legebamus, excursionem prima botanica florum delicias, postea nunquam neglectis, quasi adduxit. Anatomicus diligentissimus fuit florumque et minerarum scrutator egregius.

54. *versiformis*, gregaria parvula forma varia, cupula substipitata subcontorta olivacea, externe subpurpurascens. *Icon. et Descr. fung. p. 25. t. 7. f. 7. Consp. fung. p. 314. var. β. livido-purpurascens.*

Hab. in truncis putrescentibus abietinis in Hercynia, et *β.* in squamis putrescentibus strobilorum Pini *Abietis*, in Lusatia superiore. Cup. carnosa, 2-3 lin. lata, subcrassa, demum planiuscula, margine saepius incisa.

** LACHNEA. (Lachnum Fries.) *Cupulae piloso-strigosae aut tomentosae, (plerumque sessiles.)*

* *Cupula pilis rigidiusculis hirta.*

55. *fasciculata*, gregaria magna hemisphaerica sessilis, intus albida, extus umbrina, pilis densis fasciculatis. — *Schrad. Bot. Journ.* 2. p. 60.

Peziza hemisphaerica. Hoffm. *vegetab. cryptog.* 2. p. 23. t. 7. f. 6. *Syn. fung.* p. 640.

Peziza hispida. Sowerby. t. 147.

Peziza lanuginosa. Bull. t. 396. f. 2.

Peziza hirsuta. Holmsk. *Fung. Dan.* 2. p. 57. t. 19.

Frequens aestate et autumnio in sylvaticis umbrosis ad terram vegetat, interdum etiam in truncis reperitur. Unc. 1 magna. Junior globulosa, fere spadicea, adultior hemisphaerica, nonnunquam subeffusa, intus albo-glauca. Pili per fasciculos dispositi, concolores.

Obs. Inter figuras, quae in opere b. *Holmskioldi* hac de specie exstant, una adest, quae cupulam repraesentat velo membranaceo tectam, quod sane notabile, et a me nondum observatum est.

* *livida*, gregaria subconfluens minor, primo hemisphaerica dein explanata, medio subelevata, intus glauco-livida, extus fuliginea, pilis longis strigosis concoloribus.

Schum. Saell. p. 422. Batsch. El. fung. Cont. I. p. 221. t. 28. f. 154. quoad fig. 8.

In trunco putrido Fagi sylvaticae. Aprili.

Obs. Auctor magnitudinem hujus fungi non indicavit, quem vix ab antecedente diversum suspicor, saltem icon Batschiana laudata eadem species, sed junior videtur.

56. *brunnea*, majuscula subcaespitosa utrinque spadicea, cupula subconcava et subflexuosa, pilis minutis fasciculatis hirta. *Consp. fung. p. 317. t. 9. f. 8.*

In carbonariis sylvaticis humidis vegetat.

"Differt a priore forma et colore. Junior quidem subhemisphaerica, adulta vero constanter margine inflexa et depresso-plana. Tota semipellucens, unicolor (discus in figuris pallidus est pictus) et eleganter brunnea, ac 3 lin. lata." *Consp.*

57. *hybrida*, gregaria majuscula concavo-hemisphaerica, externe pallide brunnea hispida, intus subcoccinea (?). *Sowerb. fung. t. 369. f. 1. †*

Hab. in Anglia. Quodammodo similis *P. hispidae* (fasciculatae), sed multo dilutior et minor (ex icone 4-6 lin. lata est), et interne participat ruborem *Pez. scutellatae*. *Sow.*

58. *hirta*, solitaria submedia, cupula globosa, margine subtumido subconnivente, intus miniata, extus fusco-hirta. *Schum. p. 422.*

In sylvis Danicis rarissime ad terram nudam occurrit. Octob.

59. *turfosa*, gregaria subhemisphaerica, demum explanata, margine acuto ciliato intus coccinea, extus flava, pilis fasciculatis viridescentibus. *Schum. l. c. p. 423.* (P. Polythrichi).

Nascitur in sepibus turfosis, autumnno, inter Polythricha. Stipes obsoletus, dilatatus.

60. *marginata*, subconfluens minuta, primo hemisphaerica, dein plana, extus flavo-aurantiaca, margine tumido luteo, extus pallida, pilis strigosis sparsis. *Schum. l. c.*

Supra fol. Pini *Abietis*. Octobr.

61. *Theleboloides*, subgregaria sessilis urceolata: ore ab initio clauso lutescente, basi albidia, setis sparsis strictis concoloribus. *Consp. fung. p. 321. t. 12. f. 4.*

Nascitur ad terram pinguem in stercoratis. Forma Thelebolis simillima, sed thecis gaudet octosporis. Junior tota alba, subparabolica.

62. *stercorea*, gregaria flavo-fuscescens, setis badiis erectiusculis. *Obs. Mycol. 2. p. 89.*

Peziza equina. *Flor. Dan. t. 779. f. 3. Sowerb. t. 382.*

Octospora scutellata. *Hedw. Musc. 2. p. 11. t. 3. f. 1.*

Aestate et autumnno sat vulgaris in pratis

et sylvis supra finum. Plerumque connivet,
1-2 lin. lata.

β. equina, gregaria densa, cupulis obconicis
subtruncata flavo-succina, pilis strictis.

Autumni initio ad stercora equina obvenit.

Obs. Non *Ascoboli* species videtur, cu-
jus thecae dejectae fuerunt, nam hujus gene-
ris species nondum strigosas vidi.

Ascob. ciliatus, infra describendus, mar-
gine tantummodo ciliatus est.

63. *strigosa*, gregaria sessilis parva, cupula glo-
boso-hemisphaerica, externe setosa atra, in-
tus subochracea. *Syn. fung. p. 648. †*

P. fusco-atra, sparsa sessilis fusco-atra
tomentoso-hirta, cupulis subglobosis concavis,
intus sordide pallidis. *Rebent. Flor. Neom.*
Prodr. p. 385?

Cresc. in umbrosis ad terram subhumi-
dam. Lin. 1 lata, sicca connivet.

Obs. Rebentisch de suo fungo, forsitan
distincto, qui etiam in umbrosis supra terram
viget, dicit, quod primo oblongus sit, et Cy-
atho *striato* adhuc clauso similis, ejusque dis-
cus quidem pallidus, sed exsiccatus nigrescit;
insuper 2-3 lin. latus est.

64. *hispidula*, concava extus hispidula nigra,
intus laevis albida. *Schrad. Bot. Journ. 2.*
p. 64.

Pez. discolor, gregaria sessilis urceolata, extus atra nitens, setisque crebris patentibus ubique hispida, intus caesio - alba. *Mart. Flor. Cryptog. p. 464.?*

In ramulis dejectis et in truncis frequentius occurrit. Interne albida vel cinerea, extus setis nitentibus oblecta. *Schr.*

65. *corvina*, suberumpens utrinque atra subdepressa, externe pilosa.

Passim ad ramos et ligna. Pusilla, late aperta.

Subiculum in aliquibus adest maculiforme, pariter nigrum.

66. *variabilis*, gregaria densa sessilis, cupulis urceolato-subcylindricis flavo-virentibus setis sparsis strictis aut subnullis.

Peziza erecta. Sowerb. t. 369. f. 10.

β : *denudata*, cupulis magis cylindricis glabris. *Sowerb. l. c. f. 12.*

Hab. in locis umbrosis, suffocatis, supra muscos. Bilinearis, subobconica, margine stricto. Variare dicitur e colore obscure fulvo ad flavo-virentem aut pallidum. An β . revera eadem species? cum omnino glabra et muscigena sit.

67. *crucibuliformis*, solitaria submedia turbinata, primo epiphragmate tenuissima clausa, margine tumido involuto, albido-pallida vil-

lōsa, stipite brevi crassiusculo subsulcato.
Schum. p. 424.

Rarissima in ligno putrido Fagi *sylvaticae*.
lio. Sex lin. alta, 5 lata. — Cupula si-
ili *velo*, quod Holmskiold in *Pez. fascicula-*
observavit, ab initio clausa videtur!

1. *lasiopus*, gregaria olivaceo-umbrina, cu-
pula orbiculari planiuscula, pilis brevibus
fasciculatim sparsis, stipite sublongo tomen-
toso. *Schum. Saell. p. 426.* *Peziza tomen-*
tosa.

Cresc. in Dania, ad ligna putrida. Cupu-
4 lin. lata. Stipes a dimidia ad unc. 1. lon-
is, sursum incrassatus. *Sch.*

1. *Berberidis*, sparsa minuta rufo-fuscescens
hirta, basi setis strigosis subradiantibus cinc-
ta. *Syn. fung. p. 649.*

Perraro ad siccos *Berberidis* ramos oritur.
ipulae villis longioribus, subiculum tamen
in efficientibus, matrici adnatae, et lineares.

1. *Atropae*, gregaria minuta obconica badia
substipitata, pilis densis subrigidis erectis.
Syn. fung. p. 649.

1. *Cacaliae*. Moug. et Nestl. Stirp. Cryptog.
Vogeso-Rhenan. v. 7. n. 686.

Plantulam pulchellam in caulibus *Atropae*
elladomiae, in locis apricis montosis sylv-
cis, primo vere, prope Gottingam inveni.

Specimina vero paullatim diversa, minus obconica pilisque uti videtur crassioribus obsita, in *Soncho alpino* et *Cacalia albifronde* crescentia, in Vogesia et in monte Jura detecta sunt.

71. *Nidulus*, minuta sessilis orbicularis subdepressa substrigosa umbrina. *Kunze Fung. Germ. exsicc. Fasc. 3. no. 72.*

Cresc. in Germania, Vogesia et Helvetia, praesertim in caulibus *Convallariae multiflorae*. Minuta, subtenax, villositatem demum amittere videtur.

72. *sulphurea*, sessilis sparsa strigosa tomentosa sulphurea intus albida. *Syn. fung. Ic. et Descr. fung. p. 31. t. 8. f. 4. et 5. (P. albo-lutea).*

Ad caules herbarum; Vere Parvula; junior et exsiccata globosa, colore aliquando fuscescens, opaca.

β. *caesio-sulphurea*, subconfluens, hemisphaerica depressa, disco albo-caesia, subtus pallida, pilis longis sulphureis obsita, basi umbrina. *Schum. Saell. p. 432. (In ramulo dejecto. Octobr.)*

γ. *leucophaea*, sessilis depressa strigoso-hirta fuscescens, disco plano albido et subincarnato.

Autumno, circa Parisios, ad caul. excicatos *Urticae dioicae* nascitur. Substantia ce-

rea, mollis. Colore subbadia, sed simul dantur individua fere citrina.

73. *flavo-fuliginea*, sessilis planiuscula dilatata, extus strigoso-hirta laete sulphurea, intus olivaceo fuliginea. *Consp. fung. p. 319. t. II. f. 7.*

Ab antecedente; forma, colore, substantia subtremellosa fere, nimis recedit, quam ut conjungi possit. Gregaria, initio cupularis, dein flexuosa et applanata, etiam oblonga. Ad lignorum frustula nascitur et folia, in paludosis sylvarum, autumno. *Consp.*

74. *Martini*, gregaria majuscula depressa sessilis, cupulis orbicularibus, externe cervinis, disco applanato lutescente.

Ad ligna prope Taurinum inventa, et nonnullis abhinc annis mecum communicata fuit a b. *Martin*, clerico et strenuo quondam plantarum cryptogamicarum indagatore. Species sat pulchra, 2 lin. fere lata. Affinis *P. sulphureae*, sed multo major.

75. *rufo-olivacea*, subsolitaria sessilis orbicularis applanata, extus villosa-pulverulenta rufo-ferruginea, disco virescente-olivaceo. *Consp. p. 320. t. II. f. 4. Decand. Flor. Gall. 6. p. 25.*

Passim in ramis Rubi *fruticosi* suffocatis, aestate. Plerumque exacte circularis, disco

plano in vetustis fere nigrescente. Substantia carnosio-ceracea, molliuscula.

76. *spadicea*, sparsa sessilis subgloboso-hemisphaerica: ore connivente, tomento denso subcompacto spadiceo vestita.

Ad ramos Populi Tremulae, Aprili. Chaillet. Majuscula, substantia sicca. Juniores orbiculares, nitide badiae; adultae expallescent, opacae.

77. *flammea*, sparsa sessilis subglobosa hirta, externe rufo-ferruginea, intus nigra.

Peziza *flammea*, sessilis subglobosa strigoso-villosa unicolor flammea rubiginosa. *Consp. fung. p. 319. t. I. f. 6.*

Hab. in Lusatia, etiam prope Neuchatel, ad ligna vetusta carpinea. Primo connivens, dein subpatens, hemisphaerica. Substantia nigra, sed hymenium tegumento prorsus concolor est rubiginosum. *l. c.*

78. *caerulescens*, subsparsa minuta hemisphaerica, disco plano caeruleo, margine tumido elevato, subtus tomento-fusco obsita. *Schumacher. Saell. p. 434.*

Nascitur, in Dania, ad asseres semiputridos, autumno. Caro albida.

79. *leucocoma*, subsessilis ore lato aperta albida, basi virescens, villis densis erectis.

Rarius provenit ad ligna cariosa subvires-

centia, unde basis cupulae tali colore tincta videtur. Ex hemisphaerico subinfundibuliformis est stipite brevissimo subnigrescente.

Discus videtur olivaceo-nigrescens. Vilus non contextus, sed pilos strictos sistit.

80. *leucotricha*, majuscula sessilis hemisphaerica concava, extus intricato-strigosa candida. *Consp. fung. p. 322. t. 7. f. 5.*

Terrestris est in umbrosis humidis; Septemb. Initio globosa et clausa, dein patens, regularis. Color disci ex albo subglauescit. 3-4 lin. lata. *Consp.*

81. *ciliaris*, cyathiformis nivea, externe pilis sparsis longis obsita. *Schrad. Journ. Bot. 2. p. 63. †*

Hab. in Germania. Folia incolit varia, sed imprimis quercina. Cupula recens et humida aperta est. Ciliis longis sparsis plus minusve patulis a *Pez. virginea* differt. *Consp. p. 324.*

* *globulifera*, epiphylla sparsa exigua alba, primo clavata dein turbinata, tandem explanata, margine ciliis longis globuliferis ornata, extus stipiteque breviusculo villosis. *Schum. Saelland. p. 427. Peziza citiaria.*

Hab. in Dania, in fol. dejectis fagineis. Octob.

82. *Platani*, minutissima sessilis orbicularis

plana albida, extus nigra setosa, setis margine strictus. *Obs. Mycol. 2. pag. 87.*

Rarissime in foliis deciduis *Aceris Platanini*, in Germania, reperta. Cupula sicca conivertitur, humida folio adpressa. Setae marginem praecipue occupant.

83. *papillata*, gregaria minutissima subincarnata, disco papillata, margine ciliata. *Syn. fung. p. 650. Consp. fung. p. 321.*

Autumno in fimo sicco vaccino satis frequens.

* *vaccinea*, tota alba obconica, disco plano (papillato?), externe setis longis albis strictis.

Supra stercus vaccinum, circa Parisios, inventa, *P. papillata* multo major.

* *cervina*, gregaria minutissima subluteferruginea, cupula convexa obconica (papillata?), margine acutiusculo, extus setis longis fusco-luteis strigosa. *Schum. l. c. (P. papillata).*

In fimo cervino. Junio. Caro pallida. S.

84. *chlorina*, gregaria minutissima viridis subobconica, disco convexo, margine acutiusculo, extus setis fusco-umbrinis strigosa. *Schum. l. c. Pez. strigosa.*

In fimo vaccino sub finem aestatis. Caro viridis. S.

85. *pilosa*, solitaria minuta subhemisphaerica

depressa, disco convexiusculo carneo subtus pallidior, pilis longis erectiusculis fuscis dense vestita. *Schum. p. 433.*

Inter rariores et in ligno abietino subputrido crescit Septembri. Intus carnea. *S.*

86. *setosa*, excavata patellaeformis aurantiaca, extus setis longis erectis fuscis vestita. *Nees System. 2. p. 68. t. 37. f. 275.*

Autumno ad ligna putrida.

87. *hirtella*, gregaria subconfluens parvula, cupula hemisphaerica depressa, margine acuto, aurantiaco-lutea, subtus pallide lutea, pilis brevibus fuscis sparsis. *Schum. p. 433. (P. subhirsuta).*

Supra materiem viridem in locis humidis crescit, Majo-Junio, circa domos. Caro albo.

* *cinnabarina*, subconfluens demum hemisphaerica plana rubro-cinnabarina, margine acuto, subtus flavo-pallida hirta, margine rubescens. *Schum. p. 432.*

Eodem in loco eodemque tempore crescit uti antecedens, a qua vix differt.

88. *crinita*, exigua sessilis hemisphaerica ciliata, subtus cinerea, intus rubro-purpurea. *Bull. Champ. p. 249. t. 476. f. 2.*

Hab. in Gallia supra ligna putrida, inferne, sed praesertim ad marginem, pilis longis nigris hirta. Caro alba. *Bull.*

* *chermesina*, substipitata planiuscula hemisphaerica chermesina, extus pilosa. †

Peziza ciliata. Viborg - Holmsk. Fung. Dan. 2. p. 36. t. 18.

Hab. in ligno putrido. A *P. scutellata* forsitan non diversa, habitu tamen parum aliena est et substipitata dicitur.

89. *scutellata*, majuscula subgregaria sparsaque applanata aurantiaco - rubra, margine setis nigris hispida. Syn. fung. p. 650. Sowerby fung. t. 27. Schaeff. fung. t. 284.

Peziza aurantiaca. Bull. Champ. I. p. 247. t. 10.

Aestate nascitur ad palos prope rivulos, aut ad Salices excavatas putridas; sed semper amat loca humentia. 2 - 4 lin. lata, sessilis. Bene distincta et jam a veterioribus botanicis cognita species.

90. *melaloma*, gregaria subparva sessilis concava demum planiuscula sordide aurantiaca, margine pilis subtilissimis atris obsita. Consp. fung. p. 336. t. 2. f. 5.

Prov. in Lusatia, in carbonariis sylvaticis vetustis, passim. Lin. 1 - 2 lata, nunc orbicularis, nunc plus minusve difformis, margine varie flexuosa. Subtus aut prorsus laevis est, aut pilis atris, sparsis et obsoletis hirtiusscula, qui vero in margine constantes discum cingulo circumdant nigro. Consp.

91. *vitellina*, terrestris majuscula subcaespitosa flexuosa laete flava, ad oras setosa.

Rarissime, mense Augusto, circa Parisios, in prato quodam locisque subhumidis inventa. Carnosa, fragilis, undulata, 4 lin. circiter lata, versus marginem fimbriis erectis sparsis obsita.

- * *sabulosa*, terrestris subplana, regularis, disco cinnabarina, externe setis badiis brevibus hirta.

Hab. in Vogesia ad terram sabulosam aut in pratis siccis, Junio, rarius. (*Mougeot.*) Crassa, subdepressa, sessilis, 2-5 lin. lata, exsiccata flavescens.

92. *caerulea*, plana sessilis caerulea, margine obtuso ciliari. *Bolt. fung. t. 108. f. 2.*

Secundum auctorem in sylvis *Abietinis* ad truncos, locis humidis, oritur. Disco caerulescit, inferne nigrescit et laevis. Pili pallidi, molles. Conformatione est *Pezizae scutellatae*.

93. *episphaeria*, minuta gregaria conferta sessilis, cupulis planis, disco lutescente, externe pilis longis albis sparsis. *Martii Flor. Erlang. p. 465. †*

Autumno circa Erlangam in *Sphaeria destucta* lecta.

94. *carneo-rufa*, gregaria, cupulis urceolatis subpellucidis carneo-rufis, extus margine-

que] setis rigidiusculis albidioribus, in ore erectis, hirsutis. *Mart. Flor. Erlang. Cryptog. p. 464.*

In arena humida prope Röthelheim, in Franconia, Majo, inventa. 1-2 lin. lata.

*** Cupula tomento molliore subimbricato vestita.*

95. *coccinea*, magna stipitata turbinata (aut hemisphaerica) coccinea, externe albida subto mentosa, margine crenulato vel nudo. *Obs. Mycol. 2. p. 75. (cum Synonymis). Jacq. Austr. t. 163.*

Peziza epidendra. Bull. t. 467. Sowerby t. 13.

Peziza dichroa. Holmsk. Fung. 2. p. 21. t. 7? (Descriptio commixta videtur.)

Peziza poculiformis. Hoffm. veget. Crypt. 2. p. 27. t. 7. f. 5.

Octospora Craterella. Hedw. Musc. frond. p. 55. t. 19. f. c.? (Non valde differt, nisi cupula hypocrateriformi et stipite longiore).

Provenit species haecce elegans, initio veris, mense plerumque Martio, in sylvaticis ad terram, sed supra fragmenta ramulorum, terra tectorum, plerumque solitaria; verum ad rariores pertinet, et non ubique crescit. Magnitudine pro aëris conditione variat, ab $\frac{1}{2}$ ad 1 unc. lata. Stipes subincrassatus, longitudine quoque diversus.

Obs. Ejus nitor fuit qui Batarram ad legendos fungos excitavit; vid. Fung. Arimin. p. 27. Idem respectu mei de *Peziza aurantiaca* dicere possem.

96. *pithya*, gregaria majuscula flavo - rubra, cupula demum plana, basi cum stipite brevi villosa, villo evanescente. *Syn. fung. p. 652. Ic. et Descr. fung. fasc. 2. p. 43. t. II. f. 2. Consp. fung. p. 323. Nees Syst. 2. f. 287.*

Hab. in caudicibus, ramis, acubusque languidis Pini *Piceae*. Vigoris ejus tempus est primum ver, inter ipsam nivem deliquescentem. *Consp.* Etiam in ramulis dejectis Pini *Abietis*, Majo - Junio, in Dania. *Schum. p. 435.*

Cupula carnosula, applanata, latitudine varia, 3 - 4 lin., vix marginata (secundum cl. Schumacher margine acuto luteo - ferrugineo gaudet). Stipes brevis, nonnunquam subglobosus. An hujus subdivisionis? cum adulta fere glabra evadat.

97. *chrysophthalma*, erumpens parvula substipitata, cupula submembranacea fulva aut dilute flava, externe tomentosa pallida.

Peziza calycina. *Decand. Suppl. Flor. Gall. p. 25. excl. synonymi.*

Peziza calycina. *Schum. Saell. p. 424?* (qui fungillum suum in *strobilo abietino invenit*).

In Vogesia et Helvetia ad Pini *Abietis* ramos et cortices nascitur. Pulchella, 2 lin. et ultra lata, subfasciculata aut solitaria, valde aperta, demum fere glabrescit; recens colore est aureo-vitellina. Stipes sub cortice absconditus, crassiusculus, basi nigrescens.

98. *quercina*, erumpens sparsa minuta subsessilis, externe villosa nivea, disco subaperto laete flavo. *Syn. fung. p. 653.* P. *calycina*.

Peziza bicolor. Sowerb. fung. t. 17.

Peziza transparens. Obs. Mycol. 2. p. 87.

Octospora calyciformis. Hedw. Musc. 2. p. 78. t. 22. f. B.?

Autumno non infrequens ad ramulos dejectos quercinos observatur. Vix lineam magna, plerumque clausa, aëre praesertim sicco.

99. *pulchella*, sparsa subsessilis villosa candida, disco subaperto coccineo. *Syn. fung. p. 653.*

Peziza bicolor. Bull. Champ. p. 243. t. 410. f. 3.

Peziza Oxyacanthae. Obs. Mycol. 1. p. 41.

Vere, in ramis praesertim *Mespili Oxyacanthae*, sed minus frequens provenit. Etiam minuta et fere clausa, interne aurantio-coccinea, extus albicans.

β. flavo-coccinea, intus coccinea, extus

amoene sulphurea denseque villosa, villo subpulverulento. *Consp. fung. p. 323.*

Ab auctoribus in cortice abietino inventa. Decemb. An potius varietas. *P. chrysophthalmae?*

γ. bicolor, subconferta, primo globosa, dein explanata concaviuscula, intus flavo-aurantiaca, extus cum stipite brevi fusco-olivacea. *Schum. Saell. p. 425.*

In ligno putrido, autumn. An species?

100. *tricolor*, sparsa cupula hemisphaerica, margine tumido, intus pallide lutea, externe obscure grisea, stipite brevi albido. — *Sowerby fung. t. 369. f. 6.*

Hab. ad ligna, in Anglia. An cupula tomentosa?

101. *sericea*, sparsa stipitata majuscula, cupula campanulata ochraceo-ferruginea, extus villo viridi sericeo adpresso tecta. *Consp. fung. p. 325. t. 3. f. 7.*

Cresc. in truncis lignisque vetustis, vire dine tinctis, locis madidis, sylvaticis. Ab initio tota pulchre viridis sive psittacina, adulta ochraceo-ferruginea, tomenti simul apices cum margine cupulae introflexo albescunt nitore quodam holosericeo. *Consp.*

102. *capillaris*, solitaria fusco-virens cupula primum clavata, dein subhemisphaerica, ex-

tus stipiteque longo capillari villosis. *Schumacher. p. 425.*

Hab. in fol. dejectis fagineis. Cupula 1 lin. lata. Stip. 4 lin. longus, flexuosus, basi umbrinus. *S.*

103. *patula*, epiphylla stipitata villosa alba, disco plano ochraceo (primum albo). *Obs. Mycol. I. p. 42. Nees System. p. 70. t. 38. f. 285.*

Vere item et autumnno in fol. exsiccatis, praesertim quercinis, non equidem rara. Valde parva, sub lente observanda plantula.

104. *virginea*, gregaria tota candida, stipite sublongo, cupula turbinato-hemisphaerica villosa. *Obs. Mycol. I. p. 28. Sowerb. fung. t. 65. Holmsk. fung. 2. p. 31. t. 14.*

Peziza parvula. Flor. Dan. t. 1016. f. 4.

Peziza clandestina. Bull. t. 416. f. 5. (quoad figuram.)

Satis communis, autumnno, ad ligna et ramulos, locis praesertim umbrosis, subudis.

- β . *carpophila*, minor turbinata, densius villosa, ore connivente. *Obs. Myc. I. p. 27. Consp. fung. p. 324. In pericarpis Fagi.*

105. *clandestina*, stipitata gregaria turbinata tomentosa opace cinerea, ore connivente. *Obs. Myc. I. p. 41. Bull. Champ. I. p. 251. (quoad descriptionem).*

Nascitur in ramis siccis praesertim *Sambucinis* et *Rubi Idaei*. Sparsa, sed nonnunquam in agmine denso obvia. Cupula turbinata: villo evanescente dilute umbrina fit.

β. discoidea, cupulis patulis planiusculis, intus albidis, extus cum stipite tomento-villoso cinereo-umbrinis.

P. clandestina, *β. patula*. *Consp. fung.*

Cresc. inter rejectamenta hortorum et in sylvis.

106. *Serratulae*, sparsa stipitata turbinata grisea, cupula pubescenti-farinosa, stipite brevi glabro.

Detecta a D. Chaillot in caule exsiccatō *Serratulae arvensis*. Forma antecedentis, sed magis grisea et vix villosa est.

107. *vernalis*, subsolitaria parvula cinereo-pallida, cupula hemisphaerica, margine primo subinvolutō demum erecto, extus stipiteque elongato filiformi flexuoso villosis. *Schum. p. 425.*

In stipitibus dejectis. Aprili. 2 lin. lata.

108. *cerina*, stipitata hemisphaerica (sicca difformis) tomentosa subpulverulenta, lutescente-grisea, disco ochraceo. *Syn. fung. p. 651.* *Nees System. 2. f. 283.* *Moug. et Nestl. Stirp. Cryptog. Fasc. VII. n. 637.*

Haecce species, quae satis vulgaris, sed

minus cognita est, in lignis praesertim induratis originem sumit. Cupula globoso-turbinata, aut hemisphaerica, substantia firma, crassiuscula, lin. 1 vel $1\frac{1}{2}$ lata. Stipes brevissimus, sed crassus. Color subnitens, flavo-cerinus, aut lutescente-olivaceus.

109. *grisea*, sparsa minuta sessilis, lutescente-grisea, cupulis globosis duriusculis pubescentibus, ore minuto apertis.

Etiam ad ligna provenit, sed multo rarior. Primo intuitu Sphaeriae cuiusdam similis.

110. *pellita*, gregaria sessilis, cupulis subglobosis dense tomentosis e griseo dilute gilvis.

Hab. in ramulis Lonicerae *Caprifoliae*, in Vogesia. Vere. Molle villosa, primo clausa. Lin. 1-2 lata, forma grata.

111. *araneosa*, sparsa majuscula subferruginea opaca, cupulis brevissime stipitatis hemisphaericis late apertis glabriusculis, margine tumido subtomentoso. *Sowerby. fung. t. 369. f. 5.* (excluso Bulliardi Synonymo?) †

Ad vetustos truncos *Salicis*, in Anglia locis suffocatis crescit, 2 lin. lata.

112. *squamosa*, gregaria, cupula hemisphaerica lutea, margine tumido elevato, subtus pallide lutea, basi albida, villo squamuloso conferta vestita. *Schum. Saell. p. 434.*

Oritur in materia viridi circa domos, locis humidis. Majo-Junio. Caro pallida. S.

113. *alboimarginata*, gregaria minuta dilute umbrina villosa, cupula primo clavata, dein subhemisphaerica, dein convexa, margine tumido subinvoluta, stipite brevi cylindrico. *Schum. p. 436.*

Hab. in caulibus plantarum. Junio. Similis dicitur, quoad formam, *Pez. cyathoidi. Bull. t. 416. f. 3.*

114. *Cyathula*, sparsa subferruginea, cupula infundibuliformi tomentosa, stipite sublongo glabro.

Eam inveni, his in regionibus, ad stipites plantarum. Forma grata distincta, ore lato patula. Stipes fuscus.

115. *fuscescens*, epiphylla minuta stipitata tota brunnea, pilosa, primo globosa, dein hemisphaerica. *Syn. fung. p. 654. Consp. fung. p. 325.*

Rarius occurrit in fol. fagineis (non amplius a me observata). Secundum auctores *Conspectus*, cupula, aëre sicco, globosa, clausa; humido, semipatula et praesertim circa marginem pilis badiis vestita est.

116. *dryophila*, gregaria minutissima epiphylla sessilis orbicularis villosa badia, ut plurimum connivens.

β. varicolor, sparsa minutissima alba, pallens aut caerulescens, etiam grisea.

Copiose, saltem circa Parisios, vere et aestate in fol. *quercinis* aggregatis, supra quae ut puncta fusco-nigrescentia apparent cupulae, plerumque contractae, aliter hemisphaericae, utrinque spadiceae, excepta var. *β.* quae rarior in iisdem foliis et nonnunquam cum *α.* socialis vegetat.

117. *eruciformis*, substipitata? procumbens hirsuta nivea oblonga, ore subconstricto truncatulo. *Batsch. El. fung. p. 126. Mich. Nov. Gen. pl. p. 208. t. 86. f. 16.*

Mense Januario, in Etruria, ad truncos olim a Michelio detecta est singularis et vermiculiformis haecce species, quae decumbens, varie flexa, subteres aut oblonga, quasi truncata, sed vix lin 1. magna.

118. *fallax*, conferta subsessilis sicca, cupulis subhemisphaericis dense villosis, plerumque clausis, intus pallescens (passim prolifera).

β. albo-violascens, intus caesia vel violascens. *Consp. fung. p. 322. t. 8. f. 4.*

In variis regionibus et locis obvenit, et sub variis nominibus mihi transmissa fuit. Exsiccata enim formam habet alienam, vix Pezizoidream, ejusque discus saepe quoque proliferus. Stipite brevi est, cortici inserta. Ceterum minuta, et colore alba.

119. *pulveracea*, gregaria stipitata subindurata cinerascens, cupula subturbinata pubescente-farinacea. — *Consp. fung.* p. 342. t. 3. f. 2.

In greges satis magnos supra cortices *Betulae albae* effusa, pulvere cinereo dense oblecta, basi nigra, lin. 1 lata. Stipes validus, incrassatus.

Var. *racemosa*, stipite prolifero distincta, D. Ehrenberg mecum communicata.

120. *corticalis*, gregaria subglobosa sessilis pulveraceo-pubescent, ex cinereo subrufescens. *Syn. fung.* p. 651. *Consp. fung.* p. 322.

Equidem non rara supra cortices truncorum caesorum inveniri potest. Recens patula et magis colorata, vix lin. lata.

121. *granuliformis*, gregaria villosa globosa minuta|utrinque alba, recens obconica. *Syn. fung.* p. 651.

Peziza sessilis. Sowerby t. 389. f. 1.?

- β. *Peziza villosa.* *Syn. p.* 655. *Obs. Myc.* 2. p. 84.
γ. *incarnata*, sessilis globosa hirsuta albida, intus carnea. *Obs. Mycol. l. c.*

Peziza granuliformis. *Nees Syst.* 2. p. 67. f. 283.

Satis frequens in caulibus herbarum, praesertim *Solani tuberosi*. Sowerbeus suum fungillum ad radices putrescentes arborum inve-

nit, qui hemisphaerico-obconicus, late apertus est, et forte specie distinctus.

122. *Schumacheri*, gregaria minutissima nivea dein fuscescens globosa, subtomentosa, ore primo epiphragmate (velo?) membranaceo clauso, demum aperto, stipite obsoleto nigrescente.

Peziza granulaeformis. *Schum. Saell. p. 435.*

In ligno subputrido. Septemb.

123. *micacea*, congesta minutissima sessilis, cupulis tenerrimis farinoso-subtomentosis pallide helvolis.

In caule *Cardui nutantis* a D. Chaillet et sub nomine *chrysocoma* Bull. missa. Late disseminata, dilute cinnamomea, subnitens, nonnunquam tota pallida, arenuliformis.

124. *armeniaca*, gregaria substipitata, cupulis globoso-subturbinate pubescente-farinaceis subcinnamomeis.

Rarissime ad ligna exsiccata. Juniores subglobosae, villosae, albicant; adultae magis pulveraceae, subfulvae.

125. *cinnamomea*, erumpens fasciculata, cupulis carnosocoriaceis pubescente-farinaceis rubiginoso-lateritiis, disco superficiali. — *Decand. Flor. Gall. Suppl. p. 23.*

Hab. in principatu Neocomensi ad quer-

tuum caudices. Majo. Furfure lateritio quasi oblecta, 1 - 2 lin. lata, margine undulata.

126. *Piceae*, sparsa minuta suburceolata pulveraceo-pubescent grisea aut dilute cinerea, disco plano nigrescente.

Hab. in Pini *Piceae* foliis in monte Juratensi. Chaillet.

127. *tumida*, sparsa, cupulis subcoriaceis depresso-ventricosis, disco tumido nigrescente, externe tenue-tomentosis opacis gilvo-pallidis.

In sarmentis *Vitis viniferae* rarissime inventa a cl. Chaillet.

Ab initio, fere ut in sequente, tota clausa videtur, saltem tomento involuta, in medio dein aperitur ore circulari, discus tandem conspicitur qui est superficialis seu prominens, rugulosus quasi et immarginatus. Latitudo fungilli linearis.

128. *amorpha*, sparsa et subconfluens majuscula, cupulis subcoriaceis depressis subflexuosis tomentosis rufescente-pallidis. Syn. fung. p. 657. Decand. l. c. p. 25.

Ad caudices *Abietis*, in regionibus magis montosis, alpestribus provenit.

Aetate juniore tuberculiformis et tota villosa; adulta in cupulam expanditur saepe irregularem et passim cum vicina confluentem, latitudine 2 - 3 linearem.

Obs. Haecce species, quoad structuram melius examinata, ut genus proprium ad familiam *Thelephorarum* referri forsitan deberet et cui adnumeranda esset *Thelephora evolvens*, junior subrotunda clausa, dum evolvens cupulaeformis, extus margineque pallida tomentosa, disco glabro obscuriore (in ramis *Betulinis* crescens). *Fries Obs. Mycol.* 1. p. 154. t. 10.

**** *TAPESIA.* *Cupulae* in subiculo villosi aut submembranaceo insidentes.

* *Cupulis* tomentosis.

129. *aurelia*, sparsa sessilis subventricosa aureo-rubiginosa, subiculo tenui niveo.

Species pulchella innascitur foliis deciduis fagineis, vere in Vogesia ubi a D. Mougeot reperta. Color in fungillo recente aurorae. Cupulae majusculae, submembranaceae, subcrispatae, 2 lin. latae.

130. *stipata*, substipitata turbinata incrustans tomentosa ferruginea, exsiccata crispa subiculo tenui.

Peziza anomala. var. 1. *Obs. Mycol.* 1. p. 29. *P. rugosa.* *Sowerb. fung.* t. 369. f. 3. ?

Satis frequens in lignis, ramis et in fissuris corticum sese propagat. In statu sicco ejus habitus ita mutatur, ut alius generis speciem putares.

131. *domestica*, gregaria sessilis rufo-ferruginea? cupulis teretiusculis obovatis subtomentosis, subiculo tenui albicante. — *Sowerby* t. 351.

Ad parietes humentes calce illinitos in Anglia observata. Subiculum gossypinum, vel submembranaceum.

132. *rufiberbis*, gregaria sessilis, tomento spadiceo insidens, intus ex albido carnea, extus barba laete rufa hirsuta. *Consp. fung.* p. 329.

In ligno duro, humi jacente, Martio, in Lusatia. Cupulae concavo-planiusculae, subdifformes et subconfluunt, lin. 1 latae. *Consp.*

133. *tephrosia*, sessilis, cupulis confertis teretiusculis tomentosis cinereis, subiculo subdenso concolore.

Peziza anomala, var. *poriaeformis*. *Syn. fung.* p. 656. *Decand. Flor. Gall.* 6. p. 26.

In salice cava, olim prope Gottingam, a me inventa. Subiculum habet a 4 lin. ad unc. $1\frac{1}{2}$ latum, irregulariter effusum. Cupulae etiam subdense conglomeratae, sed formam regularem conservant. Color subumbrino-cinereus, in subiculo dilutior.

134. *conspersa*, cinerascens pulvere albicante adpersa, cupulis subhemisphaericis concavis parum distantibus.

Thelebolus hirsutus. Decand. Flor. Gall. 2. p. 272. et *Ejusd.* Syn. pl. p. 56.

Rarius progerminat in cortice arborum. Cupulae breves, subcrispae. Species nitida.

135. *caesia*, conferta, cupulis villosis applanatis basi in membranam villosam cohaerentibus, disco subtremelloso caesio. Syn. fung. p. 657. Decand. Flor. Gall. 6. p. 25.

Peziza Lichenoides. Ic. et Descr. fung. p. 29. t. 7. f. 1.

P. caesia, gregaria, pileo sessili acetabuliformi intus caesio, extus candido villosa, basi pilis longissimis constructo, sporidiis globosis. Ditmar. Fung. Germ. p. 63. t. 51.

Ad ligna quercina, in Germania et Gallia, sed rara.

136. *Rosae*, cupulis sparsis orbicularibus concavis subtomentosis spadiceis, subiculo concolore. Obs. Mycol. 2. p. 82. Decand. Flor. Gall. Suppl. p. 18.

Ad cortices ramorum *Rosae* praesertim *caninae*. Cupulae suburceolatae margine inflexo, ore minuto hiant, demum fere glabrae fiunt. Sequenti valde affinis species.

** *Cupulis glabris.*

137. *fusca*, cupulis glabris sparsis subapplanatis cinereis, subiculo latissimo fusco. Obs. Mycol. 1. p. 29. Decand. l. c. p. 18.

In

In truncis et ramis aridis crescit. Cupulae marginatae, saepius applanatae et *P. cinereae* similes, cum qua non confundenda est haec species. Subiculum tenue, sed compactum, 2-3 unc. latum.

138. *Pruni avium* (obscura), subiculo lato nigro, cupulis minutis sparsis concoloribus: ore subconnivente albicante. *Obs. Mycol.* 2. p. 82.

Peziza fusca, β . obscura. *Decand. l. c.* p. 18.

Hab. minus frequens supra corticem laevem *Pruni avium*, quem obtegit tomento tenui nigrescente. Cupulae profundius excavatae, subflexuosae, margine contractae.

139. *lateritia*, cupulis nigris majusculis glabris, subiculo denso lateritio immersis.

Rarissime circa Parisios ad ligna reperta. *Racodium* refert ob tomentum crassum, cui vero cupulae, quae planae aut subconcavae sunt, sat profunde immersae observantur.

140. *sanguinea*, cupulis minutis nigris glabris, subiculo compacto subtenui rubro. *Syn. fung. p. 657. Nees Syst. 2. p. 67. t. 37. f. 271.*

Exoritur praesertim in assulis ligneis, locis sylvaticis apertis. Crustam non latam sis-

tit, ex tomento tenui constructam. Cupulae patellaeformes.

141. *ferruginea*, cupulis hemisphaericis ferrugineis majusculis elevatis, tomento tenui subcapillari adnexis. *Tab. 1. f. 7. et 8.*

Ad ramulos quercinos, in umbrosis dejectos. Cupulae campanulatae, margine tumido. Subiculum, an potius radiculae? fibrillosum. Species pulchella, sed semel modo reperta, curatius examen hinc meretur.

142. *testacea*, subiculo latissimo granuloso-villoso argillaceo, cupulis testaceis margine granulosus.

Peziza byssiseda. *Fries Obs. Mycol. 2. p. 309.*

In Sueciae lignis pineis vegetat. Cupul. minutae sparsae, plano-concavae.

143. *flexuosa*, subiculo contexto albido, cupulis subtumidis flexuosis carneo-pallidis.

In sylvis supra terram, inter stipites et muscos rarior provenit. Faciem habet *Lichenis* cujusdam exempl. gr. *Lichenis tartarei*. *Lin.*

144. *confluens*, cupulis aggregato-subconfluentibus aurantiacis, tomento albido evanescente insidentibus. *Obs. Mycol. 2. p. 81. t. 5. f. 6. Consp. fung. p. 337.*

Oritur tam in sylvis ad loca, ubi ligna combusta fuerunt, tum alibi ad terram post ae-

difficiorum infausta incendia, passim quoque in caldariis observatur supra tubos fumiferos lutarios. Cupulae minutae, ceraceae, flavo - aurantiae, etiam miniatae.

β. rosella, sessilis glabra sparsa incarnato - rosea, cupulis carnosius minutis concavis, subiculo tenui fibrilloso. *Ehrenb. Sylv. Mycol. p. 29.*

Ab auctore in loco quoque adusto, cum *P. confluenta* reperta, a qua ut speciem separavit.

δ. lilacina, cupulis congestis vinoso - lilacinis.

Subspeciem hanc, autumnis, in locis non adustis, sed in ambulacris sub Ulmis vidi. Color ejus est plerumque vini rubri aqua diluti.

145. *xanthosia*, subiculo intertextò pallescente, cupulis? tenuissimis appianatis vitellinis. †

Ad ligna exsiccata semel inventa. *Myrothecium* refert, verum receptacula subtremellosa quidem videntur et immarginata, sed non materia soluta obiecta.

146. *porioides*, late effusa nivea, cupulis hemisphaericis glabris dense congestis, membranae tomentosae semi immersis. *Consp. fung. p. 327. t. 6. f. 5.*

In Lusatia superiore, in corticibus lignisque abiegnis, humi jacentibus, detectus pecu-

liaris hic fungus, a ceteris hujus generis speciebus parum aberrans.

Genuinam tamen sistit *Pezizam* *), dicunt cl. Auctores. Cupulae minutae, carnosae, valde regulares, extus filis byssinis membranae communi passim affixae sunt, supra quam plus minusve prominent. Membrana (subiculum) haecce tenuis quidem, sed tamen firmiuscula, candida, ejusque margo nunc subaequalis, nunc fimbriato-laciniatus, ultra 3 unc. lata. *Consp.*

†††† *PHIALEA*. Cupulae carnosae - ceraceae passim subtremellosae glaberrimae, saepius stipitatae.

147. *Phiala*, majuscula stipitata cyathiformis sordide lutea, margine integerrimo. *Flor. Dan. t. 1078. Fries, l. c. i. p. 166.*

Hab. in Dania et Suecia ad ramos alneos Cupula 3-4 lin. lata. Stipes brevis.

*) Cl. Fries proprium ex hac specie constituit genus, *Porotheleum* ab eo dictum et sequenti modo definitum:

Pileus resupinatus, expansus, papillosus: papillis distinctis superficialibus, mox depresso-porosis.

Et speciem appellavit *Porotheleum lace-rum*. Vid. *Ejus Observ. Mycol. 2. p. 272.* Ad novum suum genus? etiam retulit *Boletum fimbriatum*. *Syn. fung. p. 546.* qui vera *Poly-pori* species est.

147. *Pocula*, magna hemisphaerica subspadicea, stipite sublongo tenui. *Syn. fung. p. 659.*

Hab. in Suecia. Cupula $\frac{1}{2}$ unc. lata. Stipes 1-2 lin. longus.

148. *Ciborian*, stipitata magna digitaliformis spadicea aut cinnamomea. *Flor. Dan. t. 1078.*

f. 1. *Fries Obs. 1. p. 166.*

Prov. in uliginosis. Cupula primo ventricosa, dein campanulata, $\frac{1}{2}$ unc. lata. Stipes unc. 1. longus, basi fibrillosus.

149. *Friesii*, gregaria rufescens, cupula infundibuliformi, stipite longissimo flexuoso spadiceo. *Fries Obs. Mycol. 2. p. 307.* Peziza *Ciborioides*.

Ad culmos in uliginosis primo vere. Cupula subtenax erecta, demum subcrenulata, 3 lin. circiter lata. *Fr.*

150. *firma*, submagna dilute umbrina, cupula infundibuliformi, dein dilatato-repanda, stipite sublongo basi nigrescente. *Syn. fung. p. 658.*

Peziza ochroleuca. *Bolt. fung. t. 105. f. 1. Sowerb. t. 115.*

In umbrosis ad truncos et ramos. $\frac{1}{2}$ unc. lata, subtenuis. Stipes 3-4 lin. altus.

* *flavescens*, stipitata glabra, cupula planiuscula difformi, stipite basi attenuato. *Viborg in Holmsk. Fung. Dan. 2. p. 27. t. 11.*

Hab. in Dania ad ramos putrescentes. Antecedenti forma ac magnitudine similis.

151. *virgultorum*, subcaespitosa media, primo globosa alba, demum plano-convexa pallida, margine acuto, stipite longo flexuoso, basi lignoso umbrino. — *Schum. Saell. p. 431.*

Ad caules plant. majorum, etiam in stipitibus et ramis dejectis, non infrequens. *P. ochroleucae* Bolt. (*firmae*) affinis. *S.*

152. *Tuba*, stipitata tota lutea, cupula limbo plano, stipite filiformi. *Bolton fung. p. 106. t. 106. f. 1.*

Hab. in Angliae uliginosis, ad ligna putrescentia. Exacte tubiformis, $\frac{1}{2}$ unc. lata.

153. *explanata*, stipitata demum glabra subfusca, cupula planiuscula, margine denticulato. *Holmsk. fung. Dan. 2. p. 35. t. 17. †*

Hab. in Dania, autumnis, in ramis putridis. Ab initio tenuissime pilosa, dein villositatem exuit, et explanatur, latitudine 4 linearum. *l. c.*

154. *echinophila*, majuscula badia, ab initio subplana, dein infundibuliformi, stipite subtomentoso dilutior.

Peziza echinophila. Bull. champ. 2. p. 235. t. 500. f. 1.

Hab. ad vetusta (anni praeteriti) involucria *Castaneae vescae*, sed sat rara. Stipes lon-

gitudine varius pro fungi situ. Cupula 2 lin. fere lata.

155. *Infundibulum*, stipitata semiglobata farcta, parum excavata, fusco-ferruginea, stipite sursum incrassato cum cupula confluyente. *Batsch El. fung. p. 211. t. 27. f. 147. †*

Peziza calyculus. Sowerb. t. 116. Syn. p. 660.?

Circa Jenam in ramulo demortuo, orta. 3 lin. magna. Quoad formam antecedenti exacte similis; sed color et locus differunt.

156. *Buccina*, gregaria majuscula infundibuliformis obscure flava, stipite incrassato subincurvo. *Syn. fung. p. 659.*

Rarius ad ligna et ramulos abietinos. Pro diversa aetate, forma valde varia. Juniores incurvo-clavatae, adultae campanulatae.

157. *merulina*, carnosio-tremellosa vitellina, cupula infundibuliformi, externe plicata: plicis longitudinalibus.

Rarissime ad Salices inventa. A latere conspecta *Merulium* refert ob plicas in stipitem decurrentes. Latitudine 2 lineae.

158. *bolaris*, majuscula subsolitaria, cupula hemisphaerica subinfundibuliformi subochracea, disco badia, externe rugoso-venosa, stipite breviusculo nigricante. *Syn. fung. p. 658. Batsch El. fung. p. 221. t. 28. f. 155.*

Passim ad ramulos aridos. Stipes firmus, $1\frac{1}{2}$ lin. longus. Cupula totidem lata, in sicco margine subinvoluta, valde regularis.

159. *Digitalis*, stipitata tenuis submembranacea, cupula pendula digitaliformi, intus ex albo caesia, extus umbrina (cervina). *Consp. fung. p. 315. t. 5. f. 1.*

P. cervina, cespitosa cervina, cupula tenui subfarinacea, margine inflexa, intus pallida. *Syn. fung. p. 647.?*

Hab. in Germania, Helvetia et Vogesia ad corticem Pini *Piceae*.

"Stipes 1 - 2 lin. longus, incurvus, tenuis, umbrinus. Cupula $\frac{1}{2}$ unc. fere alta, 2 lin. lata, margine erecto, basi obliqua, hinc subventricosa, nunc laevis, nunc fibrillis parvis oppressis striata. E cupulis farina albida (sporulae conglomeratae) copiose saepius effusa simul adest in cortice." *Consp.*

160. *Perula*, sparsa parvula, cupula submembranacea elongata infundibuliformi albida (exsiccata nigrescente), stipite brevi nigrescente.

Semel inveni in umbrosis uliginosis ad caules aridos *Angelicae sylvestris*, autumnis. Forma marsupii. Cupula ore lato aperta, 2 lin. alta, substantia tenuissima.

161. *lacera*, stipitata membranacea parva albida, cupula campanulata, intus rugoso - ve-

nosa, demum multifido-lacera: margine sub-revoluta. *Consp. fung. p. 316.* (P. membranacea.) †

Cresc. ad stipites semiputridos humidos, praesertim *vitis*, autumno. Cupula albida, aut flava, extus cum stipite tenui brevi cinereo-nigricans, fibrillis densis striata, 3-4 lin. alta. An *Meruliis* affinior. *Consp. l. c.*

162. *Capula*, stipitata campanulata, membranacea subdimidiata? nivea, limbo difformi subsinuato, stipite tenui. *Viborg in Holmsk. Fung. Dan. 2. p. 41. t. 22.* †

β. *cernua*, sparsa albido-pallida, cupula cyathiformi cernua, margine ut plurimum laciniato: laciniis subinvolutis, stipite breviusculo demum umbrino. *Schum. Saell. p. 421.*

"In lignis putrescentibus *Sambuci nigrae*, in Dania, mense Augusto. Infundibulum minimum refert, adolescens vero forma est globosa ore rotundo, et interdum usque ad stipitem fisso; tandem adulta oblonge campanulata, modo dimidiata, modo obliqua." *Viborg.*

Obs. Ni fallor, potius ad familiam *Thelephoream* spectat et forsitan species subgeneris *Epibryi*.

163. *laciniata*, cupula hemisphaerica subventricosa umbrina, margine stellatim fisso, stipite brevi fusco. *Consp. fung. p. 333. t. 7. f. 6.*

Nasc. in foliis Salicis *Capreae*, locis humentibus Aestate. 2 lin. lata, venis eleganter reticulata. Stipes $\frac{1}{2}$ lin. longus subincurvus. Fungillus patulus Didermatis *stellaris* peridium non male refert, sed fructificat more *Pezizarum* fumo semineo crebro avolante. *Consp.*

164. *cernua*, parva stipitata, cupulis infundibuliformibus coccineis cernuis. — *Mich. N. Gen. pl. p. 205. t. 86. f. 12.*

Tempore brumali, per montis Senarii saltum, in Italia supra emortua ligna olim inventa a Michelio.

165. *Calicium*, gregaria stipitata glabra disco flavo, extus stipiteque subadscendente albis. *Fries Obs. Mycol. 2. p. 305. (P. bicolor).*

Peziza bicolor, var. *Sowerby fung. t. 369. f. 1.*

Hab. in Anglia et Suecia ad ligna, 2-3 lin. alta. Elegans species *Calicium* refert. Nomen *bicolor* jam plures gerunt species, hinc mutandum esse mihi videtur.

166. *Subulipes*, cupula hypocrateriformi lateritia, stipite longissimo tenui.

Peziza subularis. Bull. champ. I. p. 236. t. 500. f. 2. Decand. Syn. pl. Gall. p. 17.

Super semina semiputrida *Helianthi annui* et *Bidentis tripartitae* oritur. *B.*

167. *fructigena*, gregaria pallida, cupula carnosa subinfundibuliformi hemisphaerica, sti-

pite longo subflexuoso deorsum attenuato saepiusque incurvo. *Obs. Mycol. I. p. 42. Bull. t. 228. Sowerby t. 117.*

P. fructigena, patellaeformis lutescens, disco fuscescente, pedicello longo flexuoso, basi floccosa adhaerente, sursum incrassato. *Nees ab Esenbeck System. 2. p. 71. f. 272.*

Hab. in sylvis frequens ad glandes quercinas, aut ad ramulos siccos.

Obs. Pez. Calyculus Batsch. El. fung. t. 12. f. 57. et Carpinì Batsch El. p. 215. f. 150. nec non Pez. convexa Holmsk, Fung. 2. p. 54. t. 16. non differre videntur. Etiam sequens hujus tantum varietas videtur.

* *salicina*, ochracea suberumpens, cupula flexuosa, stipite longiusculo tenui recto. *Syn. fung. p. 663. (Ad ramos Salicinos).*

168. *Caucus*, unicolor pallide fuscescens, cupula vasculiformi margine erecto, stipite crassiusculo cupula brevior. *Rebent. Flor. Neomarch. Prodr. p. 386. t. 4. f. 17.*

Peziza amentalis. Schum. Saell. p. 418.

Hab. in Germania et Dania, vere, in amentis deciduis putrescentibus *Populi albae*.

Quoad *iconem*, Cupula est majuscula, vasculiformis, spadicea; et stipes filiformis, saltem ratione cupulae, valde tenuis.

169. *pyriformis*, cupula globoso-turbinata sor-

dide flavescente, stipite elongato filiformi al-
bido. *Fries Obs. Mycol.* 2. p. 308.

β. *Octospora pyriformis.* *Hedw. Musc.*
p. 32. t. 10. f. 8.

Ex alis foliorum quorundam muscorum
(*Bryi ligulati* et *palustris* praesertim) propul-
lulat. Cupula subdepressa, flavo-fuscescens
margine concolore Fr. (ex *Hedwigio* margine
albido).

170. *Campanula*, parva, cupula membranacea
campanulata flava, stipite filiformi subbre-
Nees Syst. 2. p. 71. t. 38. f. 295.

Hab. . . . Color pallide citrinus. Cupula
erecta ore lato apperta.

171. *Cyathoides*, sparsa parva albido-pallens,
cupula subcarnosa globoso-cyathiformi, sti-
pите longiusculo, deorsum subcrassiorc. *Syn.*
fung. p. 662. *Bull.* p. 250. t. 416. f. 3.

Peziza Solani. *Obs. Mycol.* 2. p. 80.

Non infrequens in caulibus herbarum. Mi-
nuta, stipite pro ratione longo donata, colore
passim incarnata. Quomodo sequens differat,
non perspicio.

* *tenella*, infundibuliformis tenuis candi-
da demum lutescens, pedicello filiformi. *Nees*
l. c. p. 71. f. 294. *Batsch El.* p. 150. f. 151.

172. *Scutula*, gregaria major luteo-rufescens,
cupula orbiculari, utrinque planiuscula, mar-
gine prominente, stipite sublongo tereti.

Peziza herbarum. Nees. t. 38. f. 232. cupulae formam et colorem non male refert.

Etiam ad caules plantarum crescit. Antecedente multo major est, et ab ea cupula hypocrateriformi, dilatata, nec non colore opaco, obscure helvolo differt, vix tamen ut species distincta salutanda.

173. *julacea*, majuscula brunnea, cupula hemisphaerico-patula, stipite firmo longitudine vario. †

Peziza amentacea. Balb. in *Act. Acad. Taurin. vol. 2. t. 2.*

In sylvis vineae *Bianze* prope Taurinum inventa. In solis amentis putrefactis *Betulae Alni* interdum sub tellurem (?) degit, et tunc stipitata, modo autem iisdem telluri solum instratis amentis insidet, tuncque sessilis est. B. — Cupula, quae cerea dicitur, quoad iconem 2-3 lin. lata, semiorbicularis, et stipes $\frac{1}{2}$ unc. longus.

174. *Cacaliae*, sparsa turbinata firma glabra fuscescens (dilute fumosa), ore albidior subconnivente, stipite sublongo crassiusculo.

Cresc. in caulibus *Cacaliae albifrondis* in *Vogesia*. Forma *P. Serratulae*, sed glaberrima, subnitens. Substantia duriuscula. Juniores albae; adultae subinfundibuliformes.

175. *clavata*, majuscula pallide fuscescens gla-

bra, stipite crassiusculo cum cupula in clavam confluyente.

Aestatis initio ad caules herbarum majorum rare provenit. Cupula in valde adultis etiam ventricosa, plerumque infundibuliformis, colore griseo-cervina, opaca.

176. *Urticae*, stipitata sparsa, cupula submembranacea hrmsphaerica basi protracta, ex albo dilute fuscesciente, stipite longiusculo.

Majo reperta ad caulas aridos *Urticae dioicae*, locis umbrosis secus muros. Calyculum refert ore subcontracto; exsiccata crispa aut subplicata fit et quasi tenui pulvere conspersa; magnitudine communi minorum.

177. *convivalis*, stipitata glabra, cupula infundibuliformi albicante. *Batsch Elench. fung. p. 122. no. 16. Mich. Gen. pl. p. 205. t. 86. f. 14.*

Hederae foliis innascens olim observata a Michelio. Poculiformis, 2-3 lin. alta, stipite longiusculo instructa.

178. *subulata*, minuta sparsa albido-fusca, cupula primo turbinata, demum cyathiformi, margine tunido, fauce (?) dentibus subulatis primo conniventibus coronata, stipite breviusculo incurvo. *Schum. p. 418. †*

Fungillum hunc rarissime in stipitibus, loco umbroso humido dejectis, legit Auctor, m. Octobri.

179. *radiata*, parva stipitata albido-pallens, margine dentibus setaceis erectis coronata, stipite longiusculo incurvo. *Obs. Mycol.* 2. p. 86. *Syn. fung.* p. 662.

Peziza denticulata. *Flor. Dan.* t. 1016.

Peziza coronata. *Bull. Champ.* p. 251. t. 416. f. 4. (Color male dicitur ferrugineo-fus-
ligineus).

Aestatis fine ad caules herbarum majorum, passim ad ramulos crescit. Color primo albus, dein pallescit. Stipes 1 - 2 lin. longus.

180. *coronilla*, stipitata glabra alba, cupulis subhemisphaericis, margine dentibus subtriangularibus cincto.

Peziza inflexa. *Sowerb.* t. 306. non multum differt. *Bolton fung.* t. 106. f. 2. dimidio major est.

Autumno rarius ad caulem *Urticae dioicae* hanc speciem inveni, quae antecedente paullo major est, et ab ea forma dentium praecipue differt. Stipes sat longus, subincurvus.

181. *Cyathus*, sparsa, cupulis subcylindricis fuscis, intus pallidioribus, ore denticulato connivente, disco albido. *Nees in Martii Flor. Erlang.* p. 463.

In ramulis *Betulae albae* reperta. Cupula membranacea, $\frac{1}{2}$ lin. lata, extus rimuloso-substriata. l. c. An stipitata? an sessilis?

182. *Chailletii*, sparsa sessilis, cupulis suburceolatis carnosulis gilvis, margine denticulato nigrescente.

In Principatu Neoburgensi observata a felicissimo plantarum cryptogamicarum praesertim fungorum indagatore Chailletio ad caules *Angelicae sylvestris*; in Vogesia etiam provenire videtur in *Gentiana lutea*.

Juniores suburceolatae fuscescentes; adultiores magis depressae subventricosae, aqua irrigatae subturgidae et margine dentibus subtriangularibus erectis distinctius ornatae; in statu sicco, concavo-delapsae margine subinflexo. Color subcervinus aut pallide fuscescens. Species bene distincta, nec cum *Sphaeria* (?) *Patella* (*Peziza Ligustici* Decand.) confundenda, cui primo intuitu non valde dissimilis est.

Obs. Quinque quae praecedunt species, sub divisione particulari (*Odontoloma*) ob cupulae marginem dentatum enumerari merentur.

183. *Persoonii*, cupula urceolata aurantiaca, margine prominente membranaceo (scarioso) albido, stipite cylindrico roseo. *Tab. XII. f. 1.*

Elegantem plantulam una cum delineatione sub nomine imposito transmisit cl. Mougeot, aestate, in pratis uliginosis ad *Equisetum* arida detectam sed rarissime. Cupula lin. magna

na subventricosa, aurantio-rubra, basi dilu-
tior. Thecae 5-6-8-sporae. Stipes brevis.

134. *rhodoleuca*, gregaria stipitata albido-ro-
sea, cupula orbiculari marginata utrinque
plana. *Fries Obs. Mycol. 2. p. 306. †*

Vere, in stipitibus *Equiseti arvensis*, lo-
cis humidis dejectis crescit. Substantia valde
tenuis et forte *Helotii* species. Stipes tenuis,
interdum 3 lin. longus. Cupula 1-2 lin. lata,
orbicularis. *Fries.*

Diversitas ab antecedente satis luculenta
videtur, licet loco natali et colore ambae in-
ter se conveniant.

135. *Calopus*, sparsa, cupula urceolata flavi-
cante, stipite firmo elongato roseo. *Fries*
Obs. Myc. 2. p. 307.

In Suecia ad ramulos occurrit. Stipes te-
res, aequalis. Cupula flavicans, rarius rosea,
denuim explanata, lin. 1. lata, duplo longior.
Fries.

136. *Mougeotii*, gregaria conferta subsessilis,
cupulis tereti-ventricosis (suburceolatis) cum
stipite confluentibus, disco planiusculo au-
rantiaeo, marginè tenuissimo fimbriato albi-
cante. *Tab. XII. f. 5.*

Quoque in Vogesia habitat in montium
summitate prope nives liquescentes, ad terram
sed substrata crusta quadam pulverulenta. Spe-
cies pulchra, minuta, carnosae, hinc vix excava-

ta, margine scarioso lacero. Stipes non distinctus et cum cupula confluit.

187. *cupressina*, substipitata flavo-rufescens, cupulis carnosis, primo clavaeformibus dein hemisphaericis, stipite crasso basi villosa. — *Micheli Gen. pl. t. 86. f. 20.*

Peziza rufa, γ . *cupressina*. *Syn. fung. p. 667.*

In Italia secundum Micheliū in Cupressi foliis nascitur, his in locis ad folia Juniperi *Sabinae* inventa a D. Thuillier. Multam habet affinitatem cum *Peziza pithya*, sed dimidio minor, et fere glabra nec non carnososa est, etiam agmine denso folia investit, colore lacte flavo oculos alliciens.

188. *mellina*, dispersa sessilis, cupulis subcarnosis profunde concavis luteo-cerinis glaberrimis, basi villo albo subdivergente cinctis.

Hab. ad corticem abietinum prope Neuchatel. Cupulae majusculae suburceolatae, apertae, ceraceae, colore dilute flavo veloque basilari, qui vero non subiculum format, distinctae.

189. *Scleropyxis*, majuscula cervina vel gilva, cupulis subcartilagineis late apertis subobliquis intus albido-pallentibus.

Ex Helvetia eam habeo, ubi in caulibus plantarum, semper fere in consortio cum *Scleroglossa conico* aut ejus varietate adest, et

quoad naturam ad hujus familiam pertinere videtur. 2 lin. lata.

190. *aeruginosa*, substipitata glabra aeruginosa subinfundibuliformis, margine subrepando, disco concolore aut albidō. *Syn. fung. p. 662. Consp. fung. p. 334.*

Helvella aeruginosa. Sowerb. fung. t. 347. Flor. Dan. t. 534. f. 2. Syn. fung. p. 617.

Hab. passim in ligno quercino aut betulino aerugini tincto, autumnō. Majuscula, parca.

191. *Urceolus*, stipitata, extus brunnea nigro-striata, intus fuligineo-nigricans, ore contracto erecto. *Consp. fung. p. 332. t. 3. f. 4.*

Vere in ramulis alneis decorticatis. Substantia subcoriacea, ab initio ore clauso pyriiformis. Stipes crassus.

192. *elatina*, firma, recens virescens, sicca fusco-aenea, extus nigro-striata, margine lato inflexo, stipite crasso mediocri in cupulam concavam intumescente. *Consp. fung. p. 330. t. 2. f. 3.*

Vere nascitur ad ramos Pini *Piceae* emortuos. Cupula crassiuscula, obconica, 3 lin. lata, extus subfibrillosa, margine interdum subserrata. Primo fusco-virens, dein tota viridis, in disco demum explanato sordide lutescens, vel olivascens. *Consp.*

193. *fascicularis*, caespitosa sessilis opaca, cupulis tenuibus patentibus subhemisphaericis difformibus rugosis sordide olivaceis extus nigris. — *Consp.* p. 315. t. 12. f. 2.

Innascitur corticibus Populi *Tremulae* putrefactis. Cupulae 2-4 lin. latae, admodum irregulares, intusque valde plicatae. *Consp.*

194. *uveata*, caespitosa dilute fusco-umbrina subhemisphaerica concaviuscula, margine tumido, stipite brevissimo crassiusculo apice dilatato. *Schum. Saell.* p. 430.

Hab. in Dania et Suecia, in fibrillis aridis graminum, ad agrorum margines. *Julio.*

195. *ochracea*, gregaria subcaespitosa, primo elongata clavata turbinata, demum disco plano ochraceo, margine tumido subtusque pallido in stipitem sublongum tenuem flexuosum desinens. *Schum. l. c.* p. 431.

Octobri in ramo dejecto subputrido *Sambuci nigrae* inventa. Lin. 1. lata. Stipes $2\frac{1}{2}$ lin. longus. *S.*

196. *serotina*, subgregaria majuscula laete flavida, cupula subtenui planiuscula, stipite brevi. *Syn. fung.* p. 661. *Consp. fung.* p. 331.†

Ad ramulos dejectos, autumnis. "Stipes constanter praesens, 2-3 lin. plerumque longus, firmus, crassiusculus, concolor. Cupula in junioribus aequae ac in adultis convexo-planata, 2-4 lin. lata." *Consp.*

197. *citrina*, conferta subconfluens citrina, cupulis carnosis concavo-planis, stipite crasso brevi obconico dilutiore. *Syn. fung. p. 663. Batsch El. fung. t. 39. f. 218. Hedw. Musc. frond. p. 28. t. 8. f. 3.*

Peziza aurea. Sowerb. fung. t. 150.

Ubivis frequens, autumnno, supra truncos caesos, aut eorum assulas, in sylvaticis. Substantia firma, crassiuscula. Exsiccata subrufescit in disco depresso. Juniores excavatae; adultae convexiusculae; per marginem vix ullum, cum vicinis saepe coalescunt.

- β. *albicans*, subsparsa fere sessilis, primo concava hemisphaerica alba, demum applanata aut subconvexa, disco sublutescente.

Ad ramulos aridos quercinos passim.

- γ. *filicina*, substipitata, cupula excavata aurantia, stipite firmo pallidiore. *Fries Obs. Mycol. 2. p. 306. β.*

Hab. in Polypodiis putridis, locis umbrosis, roridis. An species distincta?

- * *lenticularis*, gregaria adpressa laete flava, stipite vix ullo papillaeformi nigrescente. *Syn. fung. l. c.*

Ad truncos quoque provenit, et *var.* antecedentis modo videtur. Juniores excavatae, adultiores lentiformes.

- * *nigripes*, gregaria parva concavo-plana aquo-

se pallida, stipite longiore nigrescente. *Syn.*
p. 661. *Consp. fung.* p. 332.

Non tantum in truncis semiputribus ab-
iētinis gregaria aut subsparsa occurrit, sed et
in foliis deciduis *Pseudoplatani*, autumnno et
vere. *Consp.* In eodem libro etiam enumera-
tur *var. cellaris*, cupula fere plana subdiffor-
mi sordide lutescente. Ad ligna in cellis in-
venta, mense Jan.

198. *pallescent*, conferta glabra minor dilute
flava aut albido - pallens, cupula concava,
stipite brevi crassiusculo. *Obs. Mycol.* 2. p.
85. *Sowerby t. 151.?*

Peziza lenticularis. *Hoffm. Flor. Germ.*
2. t. 13.

Hab. ad ligna exsiccata aut assulas, in syl-
vis, plerumque subiculo subcrustaeformi nigres-
cente increscens, sed etiam absque illo inveni-
tur. *P. citrina* multo minor, sparsim crescit,
substantia tenerior, sicca quasi cornea, disco
semper excavata; margine tubtunido, erectius-
culo.

199. *flava*, sparsa, cupula orbiculari subde-
pressa, intus flava, extus pallida, margine
repando acuto, stipite brevissimo concolore.
Schum. p. 413. *Fries Obs. Mycol.* 1. p. 166.

In Dania et Suecia vegetat in strobilis de-
jectis Pini Abietis. Autumnno. Concaviuscula.
Stipes apice dilatatus.

200. *chionea*, gregaria alba, cupula concava, stipite brevi crassiusculo. *Fries Obs. Mycol.* 2. p. 306.

In foliis pineis, locis humidis, primo vere. Maxime affinis *P. herbarum*, sed cupula minor et concava est, stipes vero longior. *Fr.*

201. *nivea*, subsparsa sessilis tota alba glabra, cupulis concavis subflexuosis, margine tumido. — *Batsch El. fung.* p. 117. t. 12. f. 56.

Peziza imberbis. *Bull. Champ.* t. 467. f. 11.

Passim ad ligna et assulas obvenit. Majuscula, subplana.

202. *herbarum*, conferta subsessilis carnosa, primo albida, deinde pallens aut subrufescens, demum convexa, margine evanescente. *Syn. fung.* p. 664. *Fries*, l. c. p. 305.

Hab. in caulibus plant., praesertim *Urticae dioicae*.

Individua sibi valde approximata sunt, hinc subundulata caulique adpressa, stipite brevissimo instructa.

203. *epiphylla*, sparsa sessilis vel substipitata glabra convexo-plana marginata ex lutescente demum rufa. *Syn. fung.* p. 699.

Autumno in foliis delapsis et occultis, praesertim *Fagi sylvaticae* et *castaneae*. Majuscula depresso-subtenuis, $1\frac{1}{2}$ - 2 lin. lata, carnosula,

204. *theiochroa*, gregaria minuta, primo orbiculata, demum explanata concaviuscula sulphurea glabra, extus margineque tumido pallido. *Schum. Saelland. p. 428.* *Peziza sulphurea.*

In foliis putridis Pini Abietis, Septembri. Cupula deorsum stipitiforme protracta. S.

205. *faginea*, minuta planiuscula albida, stipite breviusculo crasso. *Syn. fung. p. 664.*
Consp. fung. p. 334.

Ad pericarpia sicca faginea, locis umbris dejecta.

206. *laricina*, gregaria orbicularis applariata lutea, subtus nigra. *Balb. Misc. Botan. p. 58. t. II. f. 2.*

In Pini *Laricis* ramulis prope Taurinum. Interdum stipite brevissimo ptaedita; ex icone 2 lin. lata.

207. *xanthomela*, gregaria sessilis, subtus margineque nigrescens, disco subconvexo lutescente. *Syn. fung. p. 665.*

In sylvis abietinis montosis ad terram se propagat. Majuscula, carnosa aut subceracea, margine lacera.

208. *leucoloma*, sparsa immersa rubra applanata, exsiccata concava, albo-fimbriata. *Nees Syst. t. 37. f. 268.*

Octospora leucoloma. *Hedw. Musc. frond. t. 4. f. 1.*

Vere, in caespitibus muscorum, Phasci praesertim *cuspidati*. Majuscula carnosa.

209. *Polytrichina*, submagna aurantiaca eonca-
va, stipite crasso longo albente.

P. leucoloma. γ . muscigena. *Conspect.*
fung. p. 335.

Ex Polytrichorum majorum axillis pro-
crescit, locis paludosis, aestate. Semunciam
fere lata et alta dicitur.

210. *humosa*, sessilis subcarnosa concava lae-
vis glabra sanguineo-rubra. *Fries Obs. My-*
col. 2. p. 309.

Peziza cartilaginea. *Sowarb. fung. t.* 369.
f. 2. ex Fries.

Ad terram nudam occurrit, in ericetis et
turfosis. Carnosa, nec cartilaginea, 1 - 2 lin.
lata. *Fr.*

211. *muscorum*, sessilis planiuscula glabra ful-
va, margine acuto. *Viborg-Holmsk. Fung.*
Dan. 2. p. 48. t. 21. †

In Dania supra muscos oritur, autumnii
initio. Ob mutuam pressionem nonnunquam
polymorpha, 2 lin. lata.

212. *convexula*, sparsa immersa convexa sub-
immarginata carneo-ochracea diluteque ru-
bra. *Syn. fung.* p. 667.

Ad terram nudam passim, qua saepe in-
quinata, lin. 1. lata.

213. *aquatica*, coccinea sessilis plana vel subconvexa, margine nullo. *Decand. Flor. Gall.* 2. p. 76. *Hall. Helv. no. 2245.*

In trabibus inundatis, vel aquaeductu ligneo, et in ipso fundo sub aqua vegetat, quatione ab omnibus fungis sane differret.

214. *Capreoli*, conferta ex livido purpurascens (sicca umbrina) carnosae obconica, margine membranaceo prominente.

Aestate in sylvula Vincennes (prope Parisios) supra stercus Cervi *Capreoli*. Cupulae dense approximatae, subcrassae, lin. $1\frac{1}{2}$ latae. Deorsum pyriformis, externe subvenosa subvillosa quasi, margine acuto distincta.

215. *granulosa*, gregaria carnosae planiuscula aurantio-rubra, margine subserrata extus papillis subpellucidis rugoso-granulosa. — *Bull. Champ. I. p. 258. t. 438. f. 3.* — *Flor. Dan. t. 655. f. 2.*

Ad stercus vaccinum exsiccatum autumnovulgaris, locis praesertim sylvaticis.

β .? *caballina*, gregaria sessilis dense conferta pallida, cupulis carnosis teretiusculis aut subventricosis crassiusculis, disco parum excavato, externe verrucosis. In stercore equinovariis.

γ . *leporum*, opacius rubra sordide purpurascens substipitata, constanter concava, sub-

tus rugosa, vix granulosa. *Conspect. fung. p.*
 337. In stercore leporino hinc inde.

216. *membranacea*, subgregaria magnitudine
 media sordide lutea subvirescens, primo or-
 bicularis depressa, demum concava, margi-
 ne submembranaceo crenulato patulo. *Schu-*
mach. Saell. p. 427.

In fimeto vaccino rarius in sylvis umbro-
 sis. Julio.

* *P. bovina*, minor subcespitosa pallida, primo
 plana convexiuscula subsessilis, dein subcya-
 thiformis: margine acuto repando, stipite
 longiusculo cylindrico apice dilatato. *Schu-*
mach. p. 430. (Pez. vaccinea).

In stercore vaccino, aestate, locis umbro-
 sis rarius. Cupula 1-2 lin. lata. Stipes 2-3 lin.
 longus. Ob stipitem tali in loco notabilis.

217. *ollaris*, conferta flava, cupulis crassiuscu-
 lis subturbinatis externe rubrugosis, margi-
 ne prominente membranaceo.

In ollis terra stercoreata repletis, subhu-
 midis, autumnus. Tota sessilis et filis brevibus
 terrae affixa. Facie fere *P. citrinae*.

218. *argillacea*, sparsa, cupulis subgloboso-
 ventricosus ore contracto, basi pilis distanti-
 bus affixa. *Sowerb. fung. t. 148. †*

Inventa fuit ad argillam, ex qua caput fi-
 guratum erat. Majuscula, lin. 1-2 lata, sessi-
 lis, concava; unum individuum adultum inci-

so-lacerum est pictum, substantiam inde carnosam esse, autumo.

219. *Lonicerae*, minuta sessilis subbyssiseda ex albo fuscescens globulosa, ore connivente crenato. *Consp. fung. p. 328. t. 11. f. 8.*

In caudicibus ramisque *Lonicerae Periclymeni*, vere. Primo tota alba, dein, a margine incipiens, fuscescit, tandem (in siccis) fusco-nigra evadit. Cupula in aliis hirtiuscula basique tomentosa, ligno insidens; in aliis paene laevis et vix byssiseda. *Consp.*

220. *Pteridis*, sessilis minutissima subdifformis, sordide lutescens, extus subgranulosa opacior, margine laxius crenato. *Consp. L. c. p. 538. t. 12. f. 7.*

Vere ad stipites humidos semiputres *Pteridis Aquilinae*. Substantia molliuscula, exsiccata dura. Vix hujus generis, an potius *Tremella*?

221. *betulina*, sparsa sessilis minutissima epiphylla orbicularis convexa, dein cupularis, subtremellosa, e pallido nigricans (?). *Consp. p. 339. t. 12. f. 5.*

In fol. vetustis *Betulae albae*, vere, Sphaerulaeformis, pallens aut rufescens, in siccis umbrina; adulta e fol. prominens formam habet congenerum, hemisphaerico-concavam.

222. *sanguinolenta*, gregaria sessilis minuta glabella, dilute ochracea (pallens) ad tac-

tum cruentata. *Conspect. l. c. p. 336. t. 8. f. 7.*

Ligna varia humida ramosque alneos aut coryleos ad terram udam dejectos incolit. Intacta tota unicolor straminea, at laesa maculis rubellis pingitur. *Consp.*

223. *carnea*, sparsa majuscula sessilis glabra carneo-rufa, cupulis carnosio-tremellosis hemisphaericis, disco subdepresso, margine tumido.

Ad ligna, in Gallia, ubi nunc aggregata, nunc sparsa crescit, scutelliformis; juniores subrotundae.

224. *Alismatis*, minuta glabra sessilis carnosio-tremellosa ex cinereo-subpurpurascens, cupulis depressis, margine subcomplatis.

Nascitur, vere, in caulibus siccis *Alismatis Plantaginis*, praeteriti anni, in aqua adhuc persistentibus. Cupulae discus passim pulvere rubello tectus.

225. *lurida*, subgregaria sessilis glabra incana, margine subtusque ex ochraceo-fuscescens subfarinosa. — *Consp. fung. p. 337. Syn. fung. p. 666.*

Peziza pineti, sessilis canescente-alba reflexa, margine ochraceo-fuscescente sublurido. *Batsch El. fung. p. 201. t. 26. f. 140.*

In acubus *Pini sylvestris* et in strobilis emortuis, Aprili-Majo. Sicca e fusco nigricat.

226. *callosa*, sessilis subinaequalis ochraceo-canescens carnosae glabra, margine crasso pulvinato subinflexo. *Batsch El. fung. p. 95. t. 39. f. 219. †*

In ligno putrido juxta cisternam aquaticam inventa. Cupula crassiuscula, disco pulvinato subnitens, in junioribus umbonato, margine calloso - tumente, demum subirregularis, lin. 1-2 lata. *B.* Sequenti nimis affinis.

227. *cinerea*, gregaria applanata subtremellosa mollis cinerascens subrepanda, margine subobsoleto. *Syn. fung. p. 634.*

P. cinerea, sessilis cinerea reflexa, margine crispo sinuato - lobato. *Batsch El. fung. p. 196. t. 26. f. 137. Sowerb. t. 64. Nees Syst. 2. p. 67. t. 37. f. 269.*

Peziza callosa. *Bull. Champ. 1. p. 252. t. 416. f. 1. ?*

β. pallida, disco pallida, subtus nigrescens, margine subcrenato. *Syn. l. c.*

γ. ? albicans, ab initio semiglobularis demum applanata ceraceo - mollis. *P. subdiaphana.* *Sowerby t. 339. f. 7.*

δ. grisea, sessilis obscure grisea planata, vel margine prominente concava. *Batsch. l. c. p. 118. t. 12. f. 55.*

Frequens, vere in autumnum usque ad ligna, ramulos, in sylvis. Forma item et colore

inconstans. Aqua emollita thecas suas depellit. *Nees.*

* *uda*, subtremellosa applanata sessilis livido-fuscescens. *Syn. fung. p. 634. Consp. fung. p. 307.*

In ramis locis udis dejectis. *Syn. Ad ligna varia madentia passim. Consp. Orbicularis, planissima, crassiuscula, 1 - 3 lin. lata. Color nunc lividus, nunc in olivaceum, in disco praesertim, vergens, subius fuscescens, subfuliginosus. Consp. Conf. Pez. callosam Batsch.*

* *Alni*, exigua sparsa sessilis, primo patellata, demum plana cinereo-caesia, margine obtusiusculo suberecto. *Schum. p. 417.*

In squamis strobilorum Betulae *Alni*. Octobr. in Dania.

228. *conigena*, minutissima orbicularis, tumidula subcinerea, margine subnullo. *Syn. fung. l. c. Consp. l. c. †*

In pinetis hercynicis ad Strobilos abietinos. *Syn. In Lusatia reperta in fissuris truncorum aut in aliis lignis abietinis, locis sylvaticis udis, Nov. — Dec. Consp.*

Gregaria, tumens, subpulvinata, sessilis aut substipitata, prorsus immarginata, forma regularissima, in nonnullis subsemiglobosa, convexa. *Consp.*

229. *vinosa*, minuta subtremellosa sessilis ful-

vo - carnea aut subrosea planiuscula immarginata. *Syn. fung. p. 635.*

Ad caules plantarum (*Urticae dioicae*).

Orbicularis, tenuis, nitida, $\frac{1}{2}$ - $\frac{3}{4}$ lin. lata, passim. Vere et aestatis fine. Non confundenda cum *Tremella lacrymali*.

Obs. P. vinosa. Consp. paullo diversa videtur etiam epixyla in rantiis quercinis projectis. Auctores laudati libri enumerant quoque varietatem *subteres*, subcylindrica, disco concaviusculo.

230. *rubella*, sessilis carnosio - tremellosa subflexuosa incarnato-rubra, margine sublacinato. *Syn. l. c.*

In corticibus et lignis putridis. Initio, pro multarum more, subglobosa, dein orbicularis, applanata, $1\frac{1}{2}$ - 2 lin. lata. Substantia semipellucida, attamen firmiuscula, colore carneo-rubella. *Consp.* Margo primum integer, dein serrulato - flexuosus. *Fries Obs. Myc. 1. p. 163.*

231. *aurea*, congesta minuta sessilis planiuscula subtremellosa fulvo-nitida. *Obs. Mycol. 1. p. 41.*

Peziza chrysocoma. Sowerb. fung. t. 152. (Color nimis pallidus).

Non infrequens ad ligna et truncos. Aestate et autumnno. Cupulae subconfluunt, exsiccatae concavae, tenues.

232. *subplana*, minuta caespitosa glabra sessilis, primo globosa demum depressa subplana, margine subinvoluta, intus aurantia, externe fusco-pallida. *Schum. Snell. p. 416.*

Ad ligna putrida Pini *Abietis*, Octobri.

233. *xanthostigma*, subgregaria minuta orbicularis concava flavescenti-aurea coriacea glabra. *Fries Obs. Mycol. I. p. 166-167.*

In ligno Pini *Abietis* putrido. Cupulis tenuibus coriaceis, margine orbiculari a *Pez. aurea*, cui proxima, distincta. *Fries.*

234. *mycophila*, gregaria minuta planiuscula pallide virescens. *Fries l. c. p. 164. †*

Nascitur in *Agaricis* putrescentibus (*A. adusto*). Conf. *Myrothecium viride*.

Coriacea, sessilis, minuta, glaberrima, margine aequali integro, dein planiuscula, subconfluens, colore virescente-albida. *Fries.*

235. *bryophila*, tota glabra minutissima orbicularis, disco applanata, obscure viridis, exsiccata nigrescens.

In Vogesia supra muscos detritos, praesertim *Jungermanniam scalarem*. Substantia videtur subtremellosa, nam sicca crispata fit, ceterum minutissima, oculo inermi ut puncta nigra apparens.

236. *seriata*, sessilis glabra, cupulis subseriatis orbicularibus subconcavis: recentibus dilute viridibus, exsiccatis nigris, disco fusco.

Peziza viridis. Chaillet. In ramis quercinis, cortice denudatis, crescit. Forma regulari, satis grata, distincta. Ad minores pertinet. A sequente omnino distincta videtur.

237. *atro-virens*, sparsa subtremellosa, junior globosa viridis, adulta hemisphaerica, disco plano subcarneo. *Syn. fung. p. 635. Consp. fung. p. 308.*

In lignis putridis et siccis, carie exesis.

* *viridi-nigra*, majuscula carnosio-gelatinosa virescente-nigra subglobosa, disco cupulari concolore. *Fries Obs. Mycol. I. p. 163.*

In ligno pineo putrido, in Suecia. Differt a *P. atro-virente*, substantia firmiore, colore magis opaco, forma globosa, urceolato-cupulaeformi, et intus concolor est. *Fries.*

238. *atrata*, sessilis subglobosa glabra nigrescente-olivacea, ore connivente albicante. *Syn. fung. p. 669. Consp. fung. p. 340. Nees System. t. 37. f. 266.*

Vere ad truncos. Subglauescens, concava, extus rugulosa.

239. *olivacea*, depresaa majuscula nigrescens externe rugulosa olivacea orbicularis, margine prominente tumido. *Batsch El. fung. p. 127. t. 12. f. 51. †*

Ad truncos, in Germania. 2 - 3 lin. lata, sessilis.

240. *Patellaria*, sparsa nigra glabra appplanata

orbicularis subcoriacea, margine brevi subtumido cincta. *Syn. p. 670.*

Peziza nigra plano - patelliformis nuda. *Nees p. 66. f. 265.*

Ad ligna et ramos cortice denudatos communis. E longinquo *Lecideam* refert.

β. foveolans, orbicularis subnitida ligno immersa.

γ. picea, minor, primo disco pruinata, demum opaca.

δ. valvata, erumpens transversalis seu oblonga, (subhysterina) margine utrinque subapproximata valviformis.

Distincta forsitan species, paullo major, opaca, rimulosa quasi, e ligno erumpens quare compressa orta videtur. E *Pedemontio* misit b. *Michel*.

241. *compressa*, minutissima tenuis nigra, exsiccata compressa conchiformis. *Syn. fung. p. 670. Consp. fung. p. 340.*

Sat frequens in assulis et reliquis lignis, in sylvaticis.

Recens aut humida formam habet regularem et colorem subrufescentem externe praesertim.

242. *inclusa*, albedo-incrustans, cupulis nigris ellipticis, primo ligno immersis demum suberumpentibus, margine inaequali sublacero, disco profunde concavo pallide livido.

Nascitur supra lignum *Salicis capreae*, quod eo in tractu, ubi apparent cupulae, quae situ subparallelae sunt, dealbatum, crustamque Lichenoideam simulat. Cupulae juniores ligno sunt inclusae et sphaeriaeformes. Species quoad genus parum ambigua, forsitan ad *Stictides* pertinet, sed adulta pezizoidea.

243. *nervisequa*, seriata nigra, cupulis in subiculo indurato lineari (e foliorum nervis erumpente) insidentibus.

In fol. emortuis *Plantaginis lanceolatae* cl. Mougeot reperit et hanc speciem, proveniendi modo ab aliis distinctam.

In fol. nervis observantur lineae, pariter nigrae, super quas sparsim positae sunt cupulae parvae, subcompresso-flexuosae.

244. *salicaria*, conferta sparsaque sessilis glabra erumpens varie flexuosa e spadiceo nigricans.

Hanc reperi speciem circa Parisios, vere, in ligno carioso *Salicis albae* excavatae, cui primo immersa.

Habitu Lichenis cujusdam, sed subiculum adest nullum. Cupulae, quae minutissimae, subcoriaceae etiam sunt; nunc sparsae, plerumque vero dense aggregatae, erumpentes.

245. *obtrita*, sparsa tenuis orbicularis depressa nigra, exsiccata crispa.

Rarius ad caules plantarum majorum. Mi-

nuta sphaerulaeformis (Sph. herbarum similis), irrigata, distincte pezizoidea, cauli arcte appressa; sicca applanata, convexa aut subexcavata, difformis.

246. *Sphaerioides*, minutissima sparsa globoso-hemisphaerica nigra, ore subcinereo, maculae crustaeformi passim insidens. †

Semel observavi ad caulem exsiccatum Valerianae officinalis, in sylvaticis, oculo inermi vix visibilis. An var. *P. atratae*? conf. Nees Syst. t. 37. f. 166.

247. *strobilina*, nigra parva subconferta glabra sessilis, cupulis atris nitidis globoso-hemisphaericis profunde concavis, ore erecto.

E Germania misit cl. Funck, qui pulchellam hanc speciem *vasculiformem* in strobilo Pini *sylvestris* detexit.

248. *Lecideola*, minuta gregaria subconfluens concava nigra, crustae cinerascenti insidens. Fries Obs. Mycol. 1. p. 166. t. 4. f. 2.

"Ad terram supra folia pinea in Helvetia. Cupulae punctiformes, glabrae, margine obtuso, primo aequales, dein irregulares, crustae effusae tenui insident, *Lecideis* hinc affinis est haec species. Fr."

249. *viticola*, subseriata crassiuscula subcoriacea regularis sessilis, cupulis orbicularibus planiusculis margine tenui cinctis sordide nigris, disco superficiali rufo-carneo.

In rimis ramorum *Vitis viniferae* nascitur (*Cauvin*). Subiculum tenue, nigrescens, subpulverulentum simul adest, an ei proprium? Cupulae paullo majores, et iis *Pez. atro-virentis*, praesertim colore, haud absimiles.

250. *leucomela*, erumpens sessilis, cupulis subcoriaceis concavis, extus nigris, intus albidis. *Syn. fung. p. 670.*

Peziza fascicularis, β . *salicina*. *Conspect. fung. p. 315?*

E ramis *Coryleis* erumpentem inveni. Majuscula, substantia sicca, subtenuis, non eodem coriacea, subelevata, margine erecto.

251. *immersa*, sparsa atra minuta, terrae fere tota immersa, cupulis subglobosis margine erecto denudato. — *Sowerb. fung. t. 389. f. 2.*

In Anglia a D. *Harriman*, Lichenologo, eximio, reperta, hinc non *Verrucaria* esse videtur, cum qua habitu et crescendi more convenit.

252. *Gentianae*, subgregaria erumpens minuta, cupulis concavis umbrinis nigricantibus, disco submolli griseo-albido, margine sublacero.

Hab. in Vogesia in caulibus emortuis *Gentianae luteae*. *Mongeot.*

Caulis epidermide lacera, solitariae aut in seriem aggregatae sunt cupulae, quae substan-

tia sunt ceraceae, disco planiusculo lutescente etiam distinctae. Sub epidermide crusta quaedam nigrescens observatur, in qua residet haec species.

253. *bicolor*, plano - patelliformis nuda extus margineque cinnamomea, disco atra. *Nees ab Esenb. Syst. 2. p. 66. t. 37. f. 294.*

Ad ligna quercina, Vere. $\frac{3}{4}$ lin. lata. Thecae minutissimae, persistunt. N.

254. *leucostigma*, gregaria minuta coriacea alba, disco subcinerascente. *Fries Obs. Mycol. 1. p. 163.*

"In ligno nigricante fagineo, in quo ut granula alba apparet. Concava, junior punctiformis dein subapplanata, disco subcinerascente, exsiccato flavescente, margine integro." *Fr.*

255. *Abbotiana*, sparsa majuscula hemisphaerica sessilis glabra coriacea (?), extus alba, interne laete flava. *Sowerby fung. t. 389. f. 8. †*

Hab. in Anglia ad ligna. Primo intuitu Lichenem, ex. gr. *L. cerinum* refert, etiam auctor haesitavit, numne plantula Lichenoidea sit, sed cum nullam invenerit crustam, eam Pezizis adnumerandam melius arbitratus fuit. Cupulae lin. 1. latae, disco planae.

256. *virens*, gregaria parvula, cupulis subsparsis subrotundo-hemisphaericis excavatis unicoloribus carneo-luteis. †

P. virens, sparsa vel subgregaria crustae sordide virescenti insidens, minutissima glabella albida vel subcarnea, margine integerrimo. *Alb. et Schw. Consp. fung. p. 338. t. 10. f. 10.*

In sylvis opacis paludosis, minime rara incolit ligna abiegnae pineaeque, in maculis satis latis effusis A Lichenibus removetur substantia carnosa, in statu humido pellucida. Cupulae eumorphae, etc. *Consp.*

Obs. In specimine, jam ante plures annos, a cl. auctoribus mecum communicato, cupulae minime albae vel carneaе sunt, sed colore exacte fere conveniunt, imo et forma, cum *Lecidea luteola*, a qua tamen hanc speciem diversam puto.

Peziza diluta α . *Syn. fung. p. 668.* verus est Lichen, nempe *Lecidea Pineti* Achar.

257. *Carpinea*, caespitosa carneo-pallens, cupulis minutis subdifformibus planiusculis in subiculo carnoso concolore aggregatis. *Syn. p. 673.*

β . *fraxinea*, cupulis subglobis, exsiccatis conniventibus.

γ . *fasciculata*, erumpens, aurea, convexa, margine obsoleto compresso. — *Tode fung. Mechl. 1. p. 20. t. 4. f. 32.* sub *Tubercularia*.

Non equidem rara in cortice exsiccato *Carpini Betuli*, β . in *Fraxino* rarius oritur.

Obs. Si plures detegantur species, tali gaudentes subiculo, hae in subdivisione particulari, imo subgenere, etiam propter habitum peculiarem, a ceteris separari merentur. Var. γ . distincta species videtur, etiam ob substantiam *gelatinosam*.

258. *ocellata*, erumpens sparsa orbicularis depressa rufescens, subtus pallidior, margine prominente subinvoluta. *Syn. fung. p. 667.*

Rarissime in corticibus ramorum Populi *Tremulae*, cui immersa, margine tantum prominente. Cupulae majusculae, glabrae, applanatae, subnitidae.

259. *Lecanora*, erumpens sparsa carnosio-tremellosa patellaeformis fulva, margine sublacero furfuraceo-albido. *Kunze et Schmidt Fung. Germanici, Fasc. 7. no. 174.*

Hab. in ramis exsiccatis Salicinis, vere, in Lusatia. Priori valde affinis, et uti illa, a reliquis *Pezizis*, facie parum recedit.

260. *rufa*, sessilis gregaria minuta rufa, exsiccata subferruginea, cupula crassiuscula concava. *Obs. Mycol. 2. p. 79.*

Hab. ad strobilos deciduos Pini *sylvestris* in Hercynia.

Obs. *P. rufa*, gregaria sessilis, cupula concaviuscula rubra, extus dilutior. *Fries Obs. Mycol. 2. p. 307.* omnino alia species est (an forsitan *P. rufa*, var. β ?), nam secundum auct.

in pinetis densis ad terram crescit, ejusque cupula concava, 1 - 4 lin. lata, et sub lente ad marginem tenuissime pubescit.

261. *juncina*, sparsa sessilis glabra applanata obscure rufa, margine subprominente.

Hab. non infrequens in uliginosis ad culmos siccos, subputridos Junci *conglomerati* aut *effusi*, aestate. Minuta, adpressa, linearis.

262. *lepida*, sparsa suberumpens dura rufescens nitida, cupulis vasculosis subconniventibus.

Hab. prope Parisios, rarius ad caules plantarum, e quibus erumpere videtur, saltem iis arcte adnata stipite brevissimo. Ob substantiam compactam, ad tactum aspera, $\frac{1}{2}$ - 1 lin. magna. Primo subteres.

263. *axillaris*, minuta muscigena aurantia vasculiformis sessilis, ore lato aperta.

P. axillaris, hemisphaerica, excavata, margine obtuso, aurantia, basi pallidior, in muscis parasitica. *Nees Syst. 2. p. 67. t. 37. f. 267.*

Oritur, vere, in axillis foliorum muscorum, (*Phasci cuspidati*, *Barbulae ruralis*) ubi, primo obtutu, pro propagulo istarum plantularum haberi posset; ceterum $\frac{3}{4}$ lin. lata est et firmula. N.

264. *Amenti*, sparsa minuta substipitata glabra albido-cinascens, cupula demum convexa, margine reflexo, †

P. amenti, stipite brevi sursum diffuso in discum pulvinatum reflexum et cicatrice notatum, tota alba. *Batsch El. fung. Cont. I. p. 211. t. 27. f. 148.*

Capsulis foemineis *Amenti salicini* adnata et fine Aprilis inventa. Pileus (cupula) subfuscescens, petiolus (stipes) cinereus et pellucidior. *B.*

265. *umbonata*, gregaria glabra sessilis albido-caesia, disco subgelatinoso, ut plurimum umbonato. *Icon. et Descript. fung. Fasc. 2. p. 35. t. 9. f. 5.*

Ad ramos, locis umbrosis subudis; etiam in pericarpis alneis, fagineis, pineis provenit haec species minus communis, quae substantia mollis est, colore dilute caesia, forma dolii.

266. *subteres*, gregaria glabra subcylindrica, ex albido-glauca subfuliginosa, disco laete flavo. *Consp. fung. p. 341. †*

Sub finem Augusti, in Lusatia, observata in ligno pineo. Parva et forma prioris dicitur, ejusque discus plano-depressus est.

267. *dentata*, sparsa minutissima sessilis albido-pallens subfarinacea, margine dentata. *Id. et Descr. fung. I. p. 5. t. 1. f. 6 et 7.*

In ligno carioso, materie virescente obducta inventa, ab initio subglobosa, dein hemisphaerica, tenera, margine dentato-incisa.

268. *serrata*, minuta albida, intus carnea, externe tenuissime pilosa, margine serrata.

Hoffm. Veget. crypt. 2. p. 26. t. 7. f. 4.

Elvela pusilla. *Flor. Dan. t. 179. f. 3.*

Syn. fung. p. 668. (ex Fries).

In ligno putrescente nascitur, primo in globulum contracta.

269. *hyalina*, gregaria minutissima sessilis subglobosa aquoso-hyalina, externe subpilosa.

Syn. fung. p. 33. Consp. fung. p. 326.

β . *granulata*, mollis albo-hyalina, aetate explanata, exterius granulata. *Nees fung. 2. p. 67. t. 37. f. 270. P. hyalina.*

Ad truncos humidos ut puncta alba apparet.

γ . *Typharum*, gregaria alba plana margine pilosiuscula, stipite brevissimo.

In caulibus putrescentibus *Typhae latifoliae*, unica modo vice inventa, mense Julio. Minutissima, vix marginata. An distincta species?

270. *urceolata*, acaulis urceolata cinerea pilosiuscula. *Flor. Dan. t. 1017. f. 3.*

In assulis, in Dania occurrit. Albida, subventricosa, lin. 1. magna.

271. *diaphana*, acaulis subcampanulata aquoso-hyalina glabra.

Peziza hyalina. *Hoffm. Veget. in Hercyn. Subterr. p. 10. t. 6.*

Hab. in trabeculis semiputridis, in Hercyniae fodinis, ultra 100 *Lachter* a terrae superficie remotis. Lin. 1-2 lata, colore dilute cornea.

272. *cuticulosa*, acaulis cyathiformis membranacea alba, marginē acuta. *Dicks. Pl. Cryptog. Brit. Fasc. 3. p. 22. t. 9. f. 11.*

Ad culmos putrescentes, in Anglia reperta. Quoad iconem juniores turbinatae sunt, adultae subventricosae et pendulae?

273. *triquetra*, sessilis obconica glabra flavescent, ore triquetro.

Octospora cryptophila. *Humb. Flor. Friberg. Specim. p. III. t. 3. f. 13.*

In Saxoniae fodinis ligno saxoque adnascens copiose adesse dicitur. Gregaria, membranacea?, arhiza, catino liquefactorio similis. Substantia mollis, deliquescent. *H.*

274. *Acuum*, gregaria minutissima sessilis subfarinacea grisea, cupulis tenerrimis vasculosis, ore latiusculo.

Cresc. in foliis adhuc subviridibus Pini sylvestris, prope Neuchatel. Juveniles orbiculares, depressae, grandescendo obconicae fiunt, subventricosae.

275. *perpusilla*, gregaria albido-pallens, cupulis stipitatis tenerrimis membranaceo-subpelucidis, ore subfarinaceis.

Etiam et haec in fol. *Pini sylvestris*, inventa a D. *Chaillet*.

276. *Arenula*, glabra sessilis minutissima subglobosa, intus albida, extus subfuliginosa. *Consp. fung. p. 341. †*

In fronde putrescente *Pteridis Aquilinae*. Majo. *Arenae* granulo simillima, humens paene tota clausa. *l. c.*

277. *Scybalorum*, exigua sessilis suboblonga difformisque, margine involuto subtuberculato, intus fusca, extus nivea. *Schumacher Saelland. p. 418.*

Omnium mihi notarum minutissima, dicit autor, quae in scybalis cervinis punctatim adpersa obvenit.

278. *inconspicua*, albida glabra teretiuscula hyalina.

In fimo equino arido, aestate, post pluvias, eam vidi, sed ita minuta, ut sub lente solummodo in conspectum veniat.

††††† *BURCARDIA*. *Schmidel, Fries. Cupula* tremellosa, aut carnosio-tremellosa, glabra, disco demum pulvinato, margine in plerisque ab initio cohaerente-clausa.

Obs. Species majusculae, heteroclitae, quarum nonnullae sub peculiari genere separari possent, ex. gr. tres priores; forsam quoque *P. Turbo* et *Clavus*.

279. *inquinans*, erumpens magna atro-inqui-

nans, demum pulvinata obconica, extus rugosa subumbrina. *Syn. fung. p. 631. cum* *Synonymis, quibus adde:*

Peziza nigra. Bull. champ. 1. p. 238. t. 460. f. 1. Sowerb. fung. t. 428.

Peziza brunnea. Batsch El. fung. t. 11. f. 50. P. polymorpha Anglor.

Ascobolus inquinans. Nees fung. t. 39. f. 296. Link Berl. Mag. 1809. p. 41.

Hab. autumnno et hyemi leni ad truncos quercinos, rarius fagineos, e quorum cortice erumpit. Juniores globulosae aut teretes, cortice exteriori omnino clausae; apice demum lacero et aperto discus denudatur, e thecis octosporis compositus, quas vero prominentes, multo minus dissilientes nondum observavi.

280. *Burcardia*, solitaria terrestris umbrina globosa, margine subcontracto, disco excavato, externe reticulata. *Syn. fung. p. 632.*

Burcardia globosa. Schmidel Anal. pl. Man. 3. p. 261. t. 69. An Lycop. truncatum Linn.? vid. Act. Suec. 1812.

Hyeme crescit in pinetis umbrosis.

"Plantula juvenilis absque rugis notabilibus esse solet, etiam colore est fusca, sed paullo dilutior. Adulta 4 lin. ad. unc. 1. et ultra magna, externe mollis. Substantia interna tota viscida est, ac si gluten videas. Suprema pars (discus) opaca, nigra." *Schmid.*

281. *pura*, gregaria magna laevis gelatinosa obconica dilute carnea, margine subflexuoso.
Obs. Mycol. I. p. 40.

Nascitur ad truncos abietinos in Hercynia et Suecia (*Fries*). Formam et magnitudinem prioris habet, sed substantia valde mollis est, hinc exsiccata diminuitur, tenuis et difformis fit.

282. *sarcoides*, caespitosa polymorpha firmula incarnato - rubra, disco excavato, externe subvenosa. *Syn. fung. p. 633. Schaeff. fung. t. 323. f. 2 - 7. Bull. t. 410. f. 1. Batsch El. t. 12. f. 53. (P. porphyria).*

Lichen sarcoides. *Jacq. Misc. Austr. 2. p. 378. t. 22.*

Autumno in sylvis ad truncos caesos.

Individua interdum occurrunt ingentia, caespitibus intermixta, unc. 1. lata, regularia, modice stipitata, externe albido - subtomentosa.
Consp. fung. p. 305.

Obs. Cl. Schumacher sub hac specie, quam non male *metamorpham* denominat, sequentes enumeravit varietates: α . *sarcoides*. *Syn. l. c.*

β . *obconica*, minor, substipitata, pileo obconico, disco concaviusculo, margine acuto undulato crenato; extus subvenosa. *Flor. Dan. t. 1017. f. 1.*

γ . *lobata*, subsessilis, pileo lobato: lobis oblongo - ovatis rotundatis vel retusis, erectis
 vel

vel decumbentibus, subvenosis, magnitudine varia. *Schaeff. vol. 4. t. 313. f. 1, 3 et 6.*
Tremella dubia. Syn. fung. p. 630.

δ. *turbinata*, substipitata turbinata glabra (laevis?) margine denticulato. *Flor. Dan. t. 1017. f. 2.* *Pez. turbinata. Syn. fung. p. 634.*

ε. *cylindrica*, caespitosa minor subcylindrica glabra, disco concaviusculo, margine tumido elevato. *P. porphyria. Batsch. El. t. 12. f. 53.*

ζ. *labyrinthiformis*, lamellosa cellulosa: lacunis inaequalibus. *Schum. Saell. p. 414-415.*

η. *pellucens*, gregaria magnitudine media turbinata stipitata fuligineo-nigra, primo obconica convexa, demum disco concaviuscula subplana, extus scabriuscula, stipite brevi adscendente. *Schum. p. 413.*

In Dania, in *Betulae* ligno putrido. Septembr. Substantia aquoso-gelatinosa, dilute fuliginea. Pileus $\frac{1}{2}$ unc. latus et altus. Stip. 1-2 lin. longus. 18.

284. *Turbo*, gregaria clavato-turbinata externe tomentosa pallida, disco subexcavato convexoque rufo (aut flavo).

Helotium radicum, volva nivea primo involuocratum, stipite simplice ramosoque crasso subrugoso albo-tomentoso; capitulis subrotundis planiusculis aureis. *Consp. fung. p. 348-349. t. 8. f. 5.*

Tuberoularia flavescens. Rebentisch.

β. Pelta, minor, cupula hemisphaerica latiore, stipite brevissimo.

Tremella Peziza. Syn. fung. p. 628. no. 18.

"Fungus gregarius obvenit intra cavitates sepium putrium fabrefactorum e Pino sylvestri. Ambigit inter *Pezizae* et *Helotii* genus. Substantia stipitis firma, subcoriacea, capituli carnososa. Junior involucrio candido lanato obductus est, quod in pileo paullo post in frustula quasi farinacea delapsum fatiscit, ac totum dein evanescit. Stipes in ramos plerumque binos ternosve divisus, deorsum radicis ad instar lignum intrat. Capitulum orbiculare aut subdifforme convexo-planum, dein saepe medio subdepressum." *Consp.*

Var. *β.* cupulas offert tumidas, subemarginatas, fere sessiles, forma quadantenus *Pezicitrinae*, 1-2 lin. latas.

Obs. Color in omnibus meis speciminibus ambarum varietatum rufus est et non luteus, forsitan exsiccatione ita mutatus.

285. *Clavus*, gregaria stipitata glabra carnososubtremellosa firmula dilute purpurascens, disco convexo-plano pileiformi. *Consp. fung. p. 306. t. II. f. 5.*

β. violascens, tota amoene violacea, obconica disco plano interdum subconcavo. *Consp. l. c.*

"Foliis variis deciduis madidis, saepe intra ipsam aquam increscit, in sylvarum latebris imis umbrosissimis paludosis, praesertim Majo, Junio, et var. β . ramulos, stipites, folia incolit locis prorsus similibus, sed Sept. et Octobri.

Clavum aliquatenus refert, ad *Helotium* tamen ableganda vix esset. Admodum polymorpha est quoad stipitem, qui nunc bilinearis, nunc ad semunciam elongatus; mox teres, mox subincrassatus; sed semper firmus et crassiusculus. Discus a 3 ad 6 lin. et ultra latus, e pallido dilute purpurascens aut sublivens." *Consp.*

286. *flavo-virens*; sparsa tremelloso-coriacea parvula, (recens) flavo-virens, primo subumbilicata, demum convexo-plana, stipite crasso brevissimo.

Hab. in Vogesia in ligno mudido prope rivulos. Applanata, vix marginata, deinde subconvexa, lin. 1-lata. Color in siccis rufescit.

287. *umbilicata*, sparsa orbicularis minuta coriaceo-tremellosa nigrescens, disco planato umbilicato, vix marginata.

In caulibus exsiccatis *Gentianae luteae* in Vogesia etiam, uti prior inventa a D. *Mougeot*. Fungillus dubius, aquae immersus tremellosus parum fit, aliud ad genus forsan pertinet.

+++++ *Scleroderris* (*Dermea Fries*). *Cupulae* coriaceae, plerumque sessiles et ex cortice (fasciculatim) erumpentes, colore ut plurimum nigrae.

288. *Frangulae*, erumpens solitaria aut cespitosa coriacea opaca umbrina, cupulis applanato - truncatis vix marginatis, stipite crasso brevissimo.

Hab. in Rhamno *Frangula*, in Vogesia. *Mougeot*.

Forma a ceteris *Pezizis* aberrat, et *Tuberculariis* subsimilis, aqua tamen emollita reliquis speciebus magis conformis fit, etiam thecas profert, sporulis forma et numero variis foetas. Recens fusco - olivascens, magis convexa, et substantia minus coriacea. Magnitudine specierum minorum. An melius ad praecedentem divisionem referenda?

289. *Ledi*, subgregaria sessilis e globoso hemisphaerica regularis extus cum micore viridi fusco-nigra farinacea, ore crenato. *Consp. fung. p. 343. t. 10. f. 7.*

Crescit in ligno excorticato *Ledi palustris*, rarius in cortice ipso. Majo. Substantia dura, coriacea, sicca; forma regularis, primo margine connivente globosa, aut subventricosa, ore praesertim pulchre viridi-micans, intus dilute fuliginea. Ceterum minutissima. *Consp.*

290. *alnea*, erumpens solitaria aut subcaespitosa opaca umbrino-nigrescens, cupula subflexuosa obsolete marginata. *Syn. fung. p. 673.*

Rarius ad ramos *Alneos* (circa Gottingam hactenus inventa ibique non frequens). Vera est *Peziza*, saltem ceteris hujus sectionis speciebus non valde dissimilis. Parva, exsiccata disco excavata, vel potius margine magis prominente distincta, non vero clausa.

291. *truncatula*, subcaespitosa erumpens atra, disco plano dilatato, margine tenui cincta.
— *Syn. l. c. †*

Crescit in Germania et Suecia ad truncos *Fagi*? Cupulae 2-4, sibi adpressae, caespitulum efficiunt rotundum aut ellipticum, ad tactum durum, vix lin. latum.

292. *Ariae*, gregaria erumpens tota nigra sessilis, cupulis orbicularibus applanatis margine subelevato cinctis.

Hab. in cortice *Crataegi Ariae* Lin.

Ab initio subrotunda et indurata sparsaque crescit; cupulae demum dilatatae denseque congestae, non excavatae, satis magnae.

293. *Betuli*, erumpens subcaespitosa nigra, margine tumido.

In cortice *Carpini Betuli*, in *Vogesia*. Minuta, opaca, passim solitaria. Cupulae demum hemisphaericae.

294. *globularis*, sparsa minuta nigra, cupulis globoso-ventricosis ore connivente.

Autumno ad ramos *Betulae albae*, in Vo-
gesia. *Mougeot*. Cupulae minutissimae, for-
ma grata distinctae, atrae, non cespitosae, mar-
gine tumidulo.

295. *Pini Strobi*, solitaria erumpens minuta
opaca nigra, cupulis sessilibus profunde ex-
cavatis.

Hab. in Pino *Strobo*; Hanovia mihi mis-
sa a D. *Gaertner*. Cupulae hemisphaericae,
apertae, dimidio minores quam in sequente.

296. *Piniastri*, erumpens subcespitosa atro-ni-
tida, margine acuto, stipite brevi crassius-
sulo. *Syn. fung. p. 672*.

Satis frequens, hyeme et vere, ad ramos
abietinos pineosque. Cupulae nitidae, aequa-
les aut subflexuosae, majusculae.

β. *ramosa*, difformis, stipite longo crasso sub-
flexuoso, in ramos 2-5 cupuliferos diviso.
Consp. fung. p. 344 - 345. (Sub cortice
abiegn).

297. *farinacea*, erumpens sessilis depressa so-
litaria nigro-subrufescens disco farinacea,
margine prominulo dilatato. *Syn. fung.*
p. 672. Consp. p. 343.

In ramis exsiccatis pineis, rarius. Sub-
difformis, minuta, disco lato albido-pulve-
raceo distincta.

298. *Pyri*, caepitosa sphaeriaeformis sessilis? cinerea pulverulenta. *Syn. fung. p. 671.* †

In cortice *Pyri Mali*. Parva, colore nitide cinerea. — Specimen mihi nunc deest, ut ampliorem hac de specie proferre possim descriptionem, nec ab aliis enumeratam eam invenio.

299. *pulveracea*, parva gregaria dura basi nigra stipitata turbinata, pulvere cinereo dense oblecta, cupula subglobosa. *Consp. fung. p. 342. t. 8. f. 2.*

In cortice *betulino* sicco obvia, ubi in greges satis longos latosque effunditur, sed in caespites non colligitur. Stipes validis, cupula longior. *Consp.*

300. *Aucupariae*, caespitoso-pulvinata biformis: cupulis paucis stipitatis subturbinatis discoideis, corpusculis pluribus subuliformibus subdigitatis cinctis.

Singularis hicce fungillus a D. Mougeot in Sorbo *Aucuparia* detectus fuit. Cespitulum crassum, 3 linearem, subpulverulentum efformat, Sphaeriae *Aucupariae* non dissimilem, sed inter corpuscula teretia simplicia aut tridentata, cupulas forsitan abortas, aut aliam ob causam non evolutas, formaque regulari indutas seu hemisphaericas offert, structura cum aliis *Pezizis* consimiles. Imo adsunt quoque cespituli e cupulis solis compositi.

301. *furfuracea*, subsolitaria majuscula coriacea, margine depresso pulvinato primato fusco-grisea, intus fusco-nigricans. *Roth Catal. Botan.* 2. p. 357. t. 9. f. 3. *Consp. fung.* p. 343. †

In caudicibus *Coryli Avellanae*, aut *Alni*.

Subsemuncialis, arcte caespitosa: inde polymorpha, difformis. Color externe e griseo terreus, in fungo humectato fuscescens, fufure dilutiore satis dense conspersus. Substantia membranaceo-coriacea. Margo latus, pulvinatus, involutus. *Consp.* An cupulae stipitatae? aut sessiles?

302. *Sphaerioides*, sessilis glomerata nigricans concava, ore contracto, extus rugosa albo-pulverulenta. *Roth in Usteri Annal. Bot.* Vol. I. p. II. t. I. f. 6. †

P. sphaerioides, fasciculis erumpens, nigra glabra, primum globosa, disco demum explanato, pedicello brevi. *Nees System.* 2. p. 69. f. 281.

Sphaeria versiformis, simplex caespitosaque globifera nigricans, primo clausa sphaeroidea, dein disco aperta pezizoidea. *Consp. fung.* p. 52. t. 9. f. 3. (excluso synonymo) ex *Nees l. c.* p. 263.

Hab. in cortice ramorum juniorum praecipue *Populi Tremulae*, hyemali tempore. Fungilli subgelatinosi, tamen duriusculi, in glo-

merulos sessiles sparsos, cortice circumdatos aggregati, nigricantes; siccati: grisei, rugosi, ore contracto, punctis albis pulveraceis non raro adpersi, intus glabri, atri. *Roth.*

Obs. Subsimilem fungum, paullo tamen minorem, in cortice quercino et Pruni *spinosa* ortum ex Helvetia mecum communicavit cl. Chaillet.

303. *populnea*, cespitosa majuscula coriaceo-membranacea sessilis, cupulis subrugosis griseis; recentibus patulis. *Syn. fung. p. 671.*

Hyeme, etiam vere, in Germania, ad ramos Populi Tremulae potissimum vegetat. Substantia multo tenerior quam in reliquis, et humida subtremellosa est, cupulae 2-3 lin. latae, exsiccatae compressae, intus subnigricant.

β. fusco-aenea, erumpens majuscula aeneo-fusca rugulosa,

Rarius in Populi *nigrae* cortice provenit.

γ. bracteata, erumpens; cupulis coriaceo-membranaceis patulis, intus fuscis, extus rugulosis nigricantibus.

In Vogesia in cortice Pruni *Cerasi*? inventa. Habitu antecedentis, sed major, multoque tenerior et colore disci differt.

304. *Cerasi*, erumpens coriacea argillaceo-rufescens, primo clausa sphaerioidea, dein in cupulam superne nigricantem dilatata. *Syn. fung. p. 673. Moug. Stirp. Crypt. Fasc. V.*

no. 494. *Sphaeria dubia*. Pers. Ic. pict.
fung. Fasc. 4. p. 48. t. 20. f. 1.

Non infrequens in ramis aridis *Cerasi*. Initium hujus speciei valde dissentit ab ejus statu adulto; cespitulum aut tuberculum nimirum sistit rugosum pallide rufescens, sensim evolvendo sphaerulae subglobosae papillatae, interdum spinulosae simile. Fungi adulti forma est eadem ac congenerum, discus tunc dilatatur et margine cinctus. In eodem acervo simul non raro adsunt individua sphaeriae formia et pezizoidea.

305. *Prunastri*, substipitata opaca nigricans, cupulis subconcavis marginatis. *Syn. fung.* p. 673.

β *discolor*, subtus e luteo olivascens. *Consp. fung.* p. 345.

Nascitur in caudice *Pruni domesticae*, satis rare. Transverse et seriatim erumpit, cupulis dense pressis, hinc subflexuosis; antecedente minor. Sed in hac quoque specie, in acervulo tamen separato, aut sparsae observantur *subulae* steriles, nigrae, rugulosae, quae ostiola sphaeriae cujusdam sub cortice latentis simulant, quae verosimile ut cupulae non evolutae habendae sunt.

* *rigida*, stipitata subfasciculata utrinque nigra, junioribus tereti-conicis crassiusculis.

Sphaeria rigida. Decand. Flor. Frans. VI.
p. 152. (Supplem.)

Hab. in ramis *Pruni spinosae*. Sub cortice exoritur in formam *sphaeriae* ostiolo crasso stricto praeditae, subrufescentis aut nigrae; adulta vero in cupulam antecedentis similem, paullo longius stipitatum mutatur; perfecta talia individua in eodem specimine, mecum etiam a D. Chaillet benevole communicata, aderant.

306. *Ribesia*, substipitata laevis nigra, cupulis turbinatis conniventibus, primo clausis
Syn. fung. p. 672.

β. *Ribis nigri*, duplo major obconica, velo tomentoso hirsuto badio fatiscente obvoluta
Consp. fung. p. 344.

Sat. vulgaris in ranuis *Ribis nigri* et *rubri*.

Obs. Ab initio et haec pro *Sphaeria* summi posset, tota omnino clausa est, sed matura in discum scutelliformam aperitur. Alius et novi generis (*Hylostoma* denominandi) fungillus videtur, cujus unam alteramve adhuc possideo speciem.

LVIII. TRIBLIDIUM. *Cupula* carnosomembranacea, ut plurimum subcompressa, libera, ab initio clausa, demum rima longitudinali aut triquetra hians.

Obs. Species hujus generis, quod concedo caractere firmo non bene esse limitatum,

naturam participant coriaceam cum *Pezizis*, et forma quodammodo sunt *Hysteriorum*.

1. *pineum*, gregarium erumpens ex olivaceo nigrescens, recens orbicularis, demum compressum, oris subinflexis.

β. *strobilina*, substipitata, e badio nigricans. *Consp. fung. p. 342.*

Peziza Abietis. Syn. fung. p. 671.

Hab. in Germania, Helvetia et Vogesia ad ramos Pini *sylvestris*. Aetate juniore totum fere viride, formaque regulari, et cupulae tunc membrana *clausae* sunt. Statu siccio subcorrugantur et fere nigrae sunt, opacae, latere utrimque compressae. Var. β. ab auctoribus libri citati semel hucusque reperta fuit inter squamas coni abietini, quorum individua alia sessilia, alia brevius longiusve stipitata, et quae hanc ob causam forsan distincta aut potius varietas *P. pinastri* est. Adest quoque varietas minor magisque regularis in ramis Pini *maritimi* in Gallia meridionali inventa.

2. *crispum*, elongatum subventricosum nigrum, marginibus subcrispis. *Syn. fung. p. 101. Icon. pictae fung. Fasc. 2. p. 20. t. 7. f. 1.*

β. *elatinum*, sparsum prominens subflexuosum, disco subdenudato rufescente. *Syn. fung. add. p. XXVIII. Moug. et Nestl. Stirp. Cryptog. Vol. 5. no. 474.*

Hab. in ramis gracilioribus et (β) in ipso caudice Pini *Abietis*.

3. *caliciforme*, sparsum nudum orbiculare aut angulatum rugulosum, adultum rima oblonga aut triquetra hians. *Tab. II. fig. 3 et 4.*

Peziza Hysterium. *Syn. fung. p. 671.*
Consp. fung. l. c.

Triblidium caliciforme. *Rebentisch Index pl. circa Berol. sponte cresc. p. 40.*

Hab. in Germania, Gallia (circa Parisios: passim) supra caudices praesertim quercinos, ex aliis etiam ad tiliaceos et salicinos. Iuvenilis plantula globoso-depressa et omnino clausa est; adulta oblonga, flexuosa, et rima oblonga aut triquetra margine denticulato dehiscit, disco non denudato. Color niger, intus pallens.

4. *quercinum*, gregarium hypodermum flexuosum, primo subteres, dein subventricosum nigrescente-cinereum. *Syn. fung. p. 100.*
Tode Fung. 2. p. 5. t. 8. f. 64. (sub Hysterio). *Nees Syst. t. 39. f. 300.*

Variolaria corrugata *Bull. champ. t. 432. f. 4. (bona).* Hypoderma quercinum. Dec.

Haec species vulgaris est ad ramulos arescentes quercinos. Ab initio margines connati sunt, perithecium tunc bene clausum, quod in veris Hysteriis locum non habet, substantia insuper est mollis, subcarnosa; autumno,

dum matura est, medio valde dilatatur et aperta est fere in cupulae formam. Specimina, locis umbrosis subhumidis dejecta, etiam sporulas nonnunquam elastice protrudere videntur, hoc tamen curatiore observatione confirmari debet.

Obs. Nonne potius huc quoque referri debet *Hysterium valvatum* Nees Syst. 2. p. 75. t. 39. f. 299? seu *Xyloma Pini* Consp. fung. p. 60 — 61. t. 5. f. 8. Dehiscencia perithecii enim fere eadem est ac in caliciformi. Saltem *Phacidia libera* et *epixyla*, quibus haec species relata est in *Mycol. Hefsten* 1. p. 30., cum ceteris *Phacidii epiphyllis* non bene congruunt.

LIX. SOLENIA. *Tubuli* solitarii, elongati, membranacei. *Thecae* sporigeræ nullae?

1. *ochracea*, sparsa ferruginea subtomentosa, tubulis sursum latioribus. — *Syn. fung.* p. 675. *Hoffm. Flor. Germ.* 2. t. 8. †

Ad truncos passim. *Pezizam* omnino refert, sed ad basin fere cava, intus alba.

2. *candida*, sparsa solitaria glabra erecta candida.

P. Solen. candida. *Disp. method. fung.* p. 36. *Hoffm. t. 8. f. 1.*

In ligno fagineo in Germania mihi rarissime obvia fuit. *Tubuli* tenuissimi, cylindri-

ci, colore albi, nec fasciculati, sed solitarii sunt et inter se subdistant.

3. *fasciculata*, tubulis fasciculatis subdepressis, apicibus subincrassatis albidioribus. *Tab.*

XII. fig. 8 et 9.

Peziza S. incana. *Consp. fung. p. 346* —
347. var. forsan *dicolor*.

Peziza Solenia. *Decand. Flor. Gall. 2.*
p. 80. (excl. Synonymis).

Hab. ad ligna putrescentia aut cariosa pinea aut abietina in Helvetia et Vogesia. (*Mougeot*). Tubuli glabri, per fasciculos seu cespitulos versus apicem latiores dispositi, albidobuberruginei (qualia erant specimina vogesiaca), aut lati fere albi in specim. in Helvetia collectis, quae etiam paullo majora erant.

Obs. Haec et antecedens, in serie fungorum (praesertim generum majorum) a completis ad simplices aut vice versa, *Boleti* (*Poriae*) esse videntur, qui se tantum ut tubulos exhibent.

LX. STICTIS. Coriaceo - membranacea, cupulaeformis, sicca, ligno immersa, limbo prominente aut obsoleto.

* *Forma magis completa limboque instructa.*

1. *chrysophaea*, disco concavo rubro, limbo crassiusculo aureo. *Syn. fung. p. 674. Ic. pict. spec. fung. rariorum. Fasc. 2. p. 17. t. 8. f. 13 et 21.*

In ceteris thetini pulchre haec species
sua provenit, in quibus alia similitudo
immersa. Natura et forma magis penetrans
est, quam reliquae.

2. *radiata*, immersa tota sive. Imbo radi-
um loco aut integerrima. *Syn. fung. L.*
Consp. fung. p. 348. (Observatio.)

Peziza marginata. Sowerb. fung. citat.

3. *Peziza acididoides*, disco immerso au-
rantiaeo, margine accessorio candido subleoni
pulverulento. *Nees ab Esch. Synonym. n. 1011.*
t. 57. f. 294.

Hab. in diversarum arborum ramis et
ligulis, etiam in culmis exsiccatis. An un-
de distinguenda var. illa, Imbo integerrimo
subcalloso instructa? Acididem simulat.

Obi. *Peziza acididoides* Neesii ob mole-
rem disci aurantiacum omnino diversam spe-
ciem diceret, nisi auctor ipse ad suum fun-
gillum, *Pez. radiatam* ut synonymum addu-
xisset, et specimen sub hoc nomine benevolit
communicatum etiam totum album fuisset.

4. *arundinacea*, gregaria alba nitida, margine
prominente integerrimo, interne caerulea.

In tectis vetustis arundinaceis observata a
D. Chaillot. Primo obtutu antecedenti quidem
similis, sed aberrat margine non inciso, supra
culmum elevato, coloreque interne caeruleo.

4. mol.

4. *mollis*, albida canescens, ore parvo cinereo-nigro subprominente. +

Hab. rarius in *Pruni spinosae* ramis. Sphaeriam *ovinam* mentiens, etiam externe subtomentosa est, substantia tamen est mollis, et disco non ostiolo, sed apertura manifesta, uti in ceteris speciebus distincta.

** *ALOMEA*. *Forma subvaria*, limbo obsoleto, aut nullo.

5. *Pelvicula*, orbicularis majuscula profunde excavata laevis rufescente-pallida.

Rarius in ramis siccis *Salicis* obvenit prope *Neocomium*. Lin. 1-2 lata, exacte orbicularis, subnitida, margine corticis subprominente cincta in cupulae formam.

6. *Hippocastani*, erumpens subrotunda intus fusca, extus pallescens, limbo subobsoleto inciso.

Nascitur ad ramos *Aesculi Hippocastani*. Dimidiam lin. lata.

7. *saligna*, erumpens disciformis tenuis cereacea alba cinereo-pruinosa, nigrescens. *Consp. fung. p. 303. t. 9. f. 7. sub Tremella.*

In *Salice alba* semiviva, autumno. Nunc planissima aut subdepressa, nunc orbicularis aut subangulata, rami cortice stellatim rupto ac reflexo cincta. Substantia cereacea, non tremellosa. Vicens subglauescit. *Consp.*

8. *rufa*, gregaria erumpens subimmersa compacta subrotunda aut elliptica planiuscula immarginata.

Rarius in Germania ad ligna oritur. Vix hujus generis est, nec bene ad Tremellas amandanda esset, quamvis substantia *subtremellosa* gaudere videatur. Cum *Schizoxylis* affinitatem quoque habere, palam est, sed crusta adest omnino nulla. Tubercula ceterum sunt minuta, ultra dimidium supra lignum eminent, convexo-plana, obscure rufa.

9. *pallida*, punctiformis latens pallida, ligno ore elliptico vel rotundo dehiscente. *Obs. Mycol.* 2. p. 74. t. 6. f. 7.

Peziza punctiformis. *Syn. fung.* p. 674

Hab. ad ligna exsiccata, cariosa. Particulae ligni, sub quibus degit, parum prominent et magis laevigatae videntur. Forma plerumque ovata aut elliptica est, margine obtuso, substantia cariosa, sed valde tenui.

10. *viridis*, subrhomboidea aut elongata applanata aeruginosa. — *Dufour* (*Peziza viridis*, ined.)

β. hysterioides, oblonga subangusta utrinque attenuata virens, ligno prominente marginata.

Cresc. in Gallia meridionali ad ligna *Castaneae vescae*; *β.* ad palos. Lin. 1-2 longa, $\frac{1}{2}$ lin. lata subfarinacea.

11. *cinerascens*, majuscula coriacea elliptica immersa, disco denudato cinerascente pulverulenta. †

Ad lignum *Salicis albae*? olim circa Göttingam obvia, eamque tunc ut *Thelephoram* quandam juniorem habebam, cum qua naturam participat, verum non crescendi morem, nec formam; haec plus minusve elliptica, passim orbicularis; disco subdepresso, aetate rimosa, 2-3 lineari.

12. *farinosa*, orbicularis convexo-plana, disco farinaceo subrimoso, margine subprominente nigro.

In *Populo nigra* a D. Dufour collecta. Specimen communicatum in ipso ligno vegetat, corticis arbore forsitan postea delapso, hinc non immersum. Disciformis, etiam elliptica est, lin. 1. lata, extus nigrescens.

13. *bullata*, orbicularis convexa laevis albida.

Etiam in Gallia australi a D. Dufour detecta, sed in ramis exsiccatis *Castaneae vescae*, ubi bullatim prominet, nulloque margine cincta, nisi epidermide corticis talem referente; punctis vel maculis albis similis.

14. *nivea*, elliptica tenerrima pustulata alba.

In foliis *Pini maritimi* Lam. sat profunde observantur foveolae, a fungillo nostro evolvendo excavatae, ubi membranulae albae cer-

nuntur; ab initio pustulas efficiunt suborbiculares, quae deinde plerumque delabuntur, et foveas relinquunt.

LXI. ASCOBOLUS. Carnosus, hemisphaericus, (pezizoideus). *Thecae*, in disco, punctatim prominentes, humore sporulisque, plerumque octo, repletae, dissilientes.

1. *furfuraceus*, gregarius subconcauus fuscus aut virescens, externe furfuraceus. *Obs. Mycol.* I. p. 33. t. 4. f. 3-6. *Bull.* p. 256. t. 376. *Sowerb.* t. 18. (*P. stercoraria*).

Peziza fusca. *Bolton. fung.* t. 109. f. 4 (mala, e specimine vetusto deprompta.)

Aestate et autumnno, in pascuis, frequens supra stercus vetustum vaccinum. *Pezizam* omnino refert. In disco demum subrufescente, thecae ut puncta nigra apparent, quae vero mox dissiliunt. Cupulae marginatae.

2. *ciliatus*, subhemisphaericus glaber aurantiacus, disco plano marginato; margine tumido ciliato albo. *Schmidt in Mycolog. Heften*, I. p. 90.

Autumno in fimo vaccino, locis humidis, sylvaticis; agmine denso plerumque provenit. Cupulae minutae, sed thecae pro ratione magnae sunt. S.

3. *glaber*, minutus congestus glaber nitens

subconvexus lutescente - fuscus vel dilute spadiceus. *Obs. Mycol.* I. p. 34. t. 4. f. 7.

β . *varius*. *Ascobolus glaber*, aurantiacus vel albus congestus nitens, forma (coloreque) varius. *Bivona-Bernardi Stirp. rarior. in Sicilia prov. Manip.* 4. p. 27. t. 6. f. 4.

γ . *ruber*. *Asc. glaber*, gregarius minutus hemisphaericus convexusculus, margine acuto cinnabarinus - ruber. *Schum. Saell.* p. 437.

In fimo vaccino autumnno sat communis. Forma parum variabilis, sed (in meo fungillo) non item colore, hinc β et γ forsitan ut species distingui merentur.

4. *carneus*, glaber applanatus gregarius colore carneo. *Syn. fung.* p. 676. *Rebent. Prodrom.* p. 388.

Occurrit quoque in fimetis in hortis, sylvis, pratis, sed uti videtur, satis rarus. Attamen bene distincta species est, "semipellucens, disciformis, convexo-plana, dilute vinosa seu carnea; margine vix ullo, thecis copiosis prominentibus e more generis nigris." *Consp. fung.* p. 348.

5. *immersus*, sparsus immersus irregularis subconicus, externe subfurfuraceus. *Obs. Mycol.* I. p. 35. t. 4. f. 7. fig. d. e.

Fimo vaccino fere totus immersus est, thecis solummodo prominentibus. Species vi-

detur subdubia, nec ab aliis Mycologis inventa aut descripta.

6. *lignatilis*, major stipitatus e sordide luteo subvirens externe albido-furfuraceus; stipite breviusculo subcrasso in cupulam planam marginatam dilatato. *Consp. fung.* p. 347. t. 6. f. 6.

A cl. auctoribus visa fuerunt ingentia huius speciei *epixylae* agmina in ligneis pineis fabre-factis subterraneis; aqua *cloacina* inquinatis, ac restaurationis causa effossis aquae-ductuum.

Ab A. *furfuraceo* differt stipite, forma, magnitudine et loco. Stipes 1-2 lin. longus et crassus. Cupulae 4-6 lin. latae, theculis creberrimis prominentibus nigro-punctatae. Sed thecae istae maturae disruptae sporulas *fumi* ad instar projaculantes laticem s. humorem simul fundunt. *l. c.*

7. *Trifolii*, epiphyllus minutus gregarius dilute flavus planiusculus. *Bivona-Bernardi* *l. c.* p. 27. t. 6. f. 3. †

Hab. in Sicilia, in fol. languescentibus *Trifolii hybridi*, stercore vaccino semiobrut.

"Tuberculi epiphylli, rarius hypophylli, urediniformes, subhemisphaerico-convexi, sessiles, carnosius, glabri, epidermide rupta, aperti, ore circulari, plerumque denticulato, interdum nigrescenti, disco subdepresso punctis

nigris notato. Thecae clavatae, albae. Sporae primum albae, demum nigrae." *Bernardi*.

LXII. HELOTIUM. *Pileolus* carnosus, demum convexo-hemisphaericus, immarginatus, basi a stipite sublongo distans.

Obs. Fungilli parvi, inter *Helvellas* et *Pezizas* intermediis, his tamen affiniores, etiam superne fructificantur; sed in disco tumido seu convexo; ab iis interea habitu peculiari distincti sunt.

1. *aciculare*, gregarium minutum candidum, capitulo primo excavato. *Syn. fung. p. 677.*
Obs. Mycol. 2. p. 20 - 21. t. 5. f. 1. sub Leotia.

Helvella acicularis. *Bull. Champ. 1. p. 296. t. 473. f. 1.*

Helvella agariciformis. *Sowerb. fung. t. 57. Bolt. fung. t. 98. f. 6.*

β. *elegans*, pileo hemisphaerico roseo, stipite elongato albo, basi villosa. *Fries. Obs. Mycol. 2. p. 310.*

Ad truncos caesos quercinos. Stip. subramosus, 3 lin. longus.

2. *subtile*, gregarium minimum candidum, pileo plano-convexo, stipite brevissimo. *Fries l. c. p. 311.*

In fol. abietinis. Stip. $1\frac{1}{4}$ lin. longus,

detur subdubia, nec ab aliis Mycologis inventa aut descripta.

6. *lignatilis*, major stipitatus e sordide luteo subvirens externe albido-furfuraceus, stipite breviusculo subcrasso in cupulam planam marginatam dilatato. *Consp. fung. p. 3. t. 6. f. 6.*

A cl. auctoribus visa fuerunt ingentia jus speciei *epixylae* agmina in ligneis pilae fabre-factis subterraneis; aqua cloacina irrigata, ac restorationis causa effossis adductum.

Ab A. *furfuraceo* differt stipite, fo magnitudine et loco. Stipes 1-2 lin. latus et crassus. Cupulae 4-6 lin. latae, the creberrimis prominentibus nigro-punctatis. Sed thecae istae maturae disruptae sporulanti ad instar projaculantes laticem s. humorem simul fundunt. *l. c.*

7. *Trifolii*, epiphyllus minutus gregarius, stipite flavus planiusculus. *Bivona - Berni. c. p. 27. t. 6. f. 3. †*

Hab. in Sicilia, in fol. languescens *Trifolii hybridi*, stercore vaccino semio-

"Tuberculi epiphylli, rarius hypogaei, urediniformes, subhemisphaerico-conversiles, carnosiusculi, glabri, epidermide rupta, ore circulari, plerumque denticulati, interdum nigrescent disco subdepresso.

subaequalis, saepe curvatus. Pil. lin. 1 latus, junior planus. *Fr.*

β. album, gregarium album minutum, pileo primum plano, demum convexo-hemisphaerico, margine acuto subrepando, stipite brevi crassiusculo, basi angustato. *Schum. Saell. p. 412.*

In fol. subputridis graminum. Lin. 1 longus. *S.*

5. nigripes, sparsum minutum, pileo convexusculo, margine subrepando, hymenioque (subtus?) flavo, stipite brevi fuligineo-nigricante. *Schum. l. c. Fries l. c.*

In Dania et Suecia ad asseres putridos. Novembr.

4. elongatum, gregarium, primo pallescens, demum dilute umbrinum, stipite elongato tenui flexuoso, basi byssoideo hymenioque (pileolo inferne) concavo albo. *Schum. l. c. H. aciculare β. abietinum. Consp. fung. p. 349?*

In ligno putrido Pini *Abietis*. Stipes diaphanus, 2 lin. longus. Pileus dimidiam lin. latus. *Sch.* — Pileolus a primordio convexo-hemisphaericus, ex albo dein pallens, ac subrufescens. In trunco abiegnio putredine corrupto repertum fuit exeunte Martio. *Consp.*

5. aureum, gregarium fulvum, pileolo hemis-

phaerico convexo, stipite subtenui, basi tomentoso. *Syn. fung. p. 678. Consp. l. c.*

Hab. in Germania, Helvetia et Vogesia, in truncis lignisque siccis Pini *Abietis* et *Piceae*. Vere *H. acicularis* habitu, sed minor. Pileolus lenticularis; ab initio etiam convexus. Tomentum stipitis baseos niveum, delicatum.

6. *finetarium*, parvum amoene rubrum, pileolo planiusculo subangulato. *Syn. fung. l. c.*

Leotia finetaria. Obs. Mycol. 2. t. 5. f. 4. et 5.

β. leporinum, minus nitidum, pileolo subgloboso.

Autumno ad stercus vaccinum oritur; et *β. finum piluliformem cuniculorum* et *leporum* in sylvis, fere totum involvit, sub finem Novembris, praevis pluviis autumnalibus. Stilbis affine, saltem *β.*

7. *sessile*, cespitosum minutum pallidum, disco dilute umbrino, stipite brevissimo deorsum subbadio. *Schum. Saell. p. 412.*

In Dania, supra ramulum dejectum repertum.

+ *Species ulterius examinandae, et quoad genus dubiae.*

8. *fibuliforme*, cartilagineum, pileo superne ochroleuco, stipite brevi nigro-fusco.

Ordo secundus.

347

Anglia vulgaris dicitur. Var. α .
acta, 2-3 lin. longa videtur, sti-
gentibus, longiusculis et cras-
culi instructa; β . habitu a
ita distincta, ut fortasse
stipes simplex duplo-

illa purpurea Lin.,
ura intermedia ta-
non proposita,
nus refert.

ideo albo.

uscoides albus, se-
oliis vel lignis semipu-
Mich. Nov. Gen. pl. p. 201.

carnatum, gregarium subacaule incar-
im.

ungoidaster minimus incarnati coloris,
s ferme, Anagyridi non foetidae et semi-
ae innascens. *Micheli p. 201. t. 82. f. 4.*
tuberculariis accedere videtur.

LXIII. STILBUM. *Capitulum* cum stipite
iens, primo molle subdiaphanum, matu-
do opacum, turbidum, ex sporulis
ue thecis?) farctum. (*Fungilli muco-*
nes).

ssisedum, congestum majusculum, capi-

Helvella fibuliformis, Bolt. *Fung. append.* p. 176. t. 176.

Hab. ad ramulos aut radices alneas, prope rivulos, in Anglia. Pileolus 2-3 lin. latus crassus, Stipes brevis, crassus, vix lin. longus.

9. *umbelliforme*, pileo hemisphaerico expanso subtenui flavo, stipite subbrevis.

Helvella aurea, stipite brevi, pileo umbellifero aureo. Bolt. *fung. vol.* 3. t. 98. f. 3.

In Anglia, ad ramulos, 3-4 lin. latum.

10. *flavescens*, stipitatum glabrum flavescens, stipite basi attenuato, pilei disco elevato hemisphaerico. *Holmsk. Fung. Dan.* 2. p. 34. t. 16. *Peziza convexa*.

Hab. in Dania, aestate. Fungillus ab initio est concavus, extus albicans, intus flavus, adultus vero habitum immutat: discus tunc elevatus et margo reflectitur, hinc pagina inferior concava evadit. l. c.

11. *coccineum*, subfasciculatum, capitulo subrotundo coccineo, stipite crassiusculo flavescente.

Clavaria coccinea. Sowerb. *fung.* t. 294. fig. sinistra.

β. *brachypus*, gregarium, capitulo subhemisphaerico convexo (semigloboso) coccineo, stipite brevi crasso. — Sowerby l. c. fig. dextra.

Species in Anglia vulgaris dicitur. Var. α . in ramulo depicta, 2-3 lin. longa videtur, stipitibus basi cohaerentibus, longiusculis et crassitudine fere capituli instructa; β . habitu a priore, quod iconem, ita distincta, ut fortasse speciem sisteret, ejusque stipes simplex duploque brevior et crassior est.

Obs. Non est Tremella *purpurea* Lin., uti auctor putat, interea figura intermedia tabulae citatae, lente vero aucta non proposita, Tuberculariam vulgarem aliquatenus refert.

12. *album*, parvum, pileolo discoideo albo.

Fungoidaster minimus, discoides albus, semine ovato donatus, foliis vel lignis semiputridis innascens. *Mich. Nov. Gen. pl. p. 201. t. 82. f. 3.*

13. *incarnatum*, gregarium subacaule incarnatum.

Fungoidaster minimus incarnati coloris, acaulis ferme, Anagyridi non foetidae et semimortuae innascens. *Micheli p. 201. t. 82. f. 4.* — Tubercularius accedere videtur.

LXIII. STILBUM. *Capitulum* cum stipite confluentis, primo molle subdiaphanum, maturescendo opacum, turbidum, ex sporulis (absque thecis?) farctum. (*Fungilli mucoriformes*).

1. *byssisedum*, congestum majusculum, capi-

tulo subrotundo pallescente, stipitibus basi byssoidea connexis.

Hab. in Vogesia, ad ligna pinea mucida. Ab initio totum albidum et subuliforme, adultum capitulo globoso subovato terminatum. Stipes crassiusculus, albidus.

2. *strigosum*, capitulo rotundato, stipite ochraceo strigoso hirtio. *Syn. fung. p. 680. Hoffm. Flor. 2. t. 10. f. 2.*

Rarissime ad truncos putrescentes obvenit. Verum est stilbum; capitulum saltem est farctum, non pulveraceum.

3. *erythrocephalum*, capitulo globoso roseo, stipite crassiusculo piloso albido. *Ditmar fung. germ. p. 91. t. 45.*

Ad excrementa avium? inventum. Capitulum ovatum?

4. *xanthocephalum*, gregarium capitulo globoso fulvo, stipite tenui nigrescente rigido. *Ditmar loc. cit. t. 60.*

Stilbum minimum, α . *nigripes. Tod. Fung. Meckl. 1. p. 11. t. 2. p. 19.*

Aestate-autumno super vetustos truncos nascitur.

5. *rigidum*, stipite subulato rigido nigro persistente, capitulo primum lacteo, demum compacto griseo deciduo. *Usteri Annal. Bot. Fasc. 1. p. 31. t. 2. f. 2.*

Hab. vere ad truncos caesos, ubi, capitulo deciduo, stipites ut setulae, nonnullis Byssoideis similes remanent.

β. atrum, gregarium, capitulo globoso, primo albedo demum nigro subturbinato, stipite passim prolifero crassiusculo rigido. *Ditmar l. c. p. 49. t. 59. St. rigidum.*

Nonne species distincta? habitu valde differt. Stipes multo crassior, interdum ramosus, et capitulum in valde adultis pyriforme fit totumque nigrum est, quod in *α.* nondum observavi.

* *nigrum*, totum nigrum durum, capitulo etiam maturo persistente. *Decand. Synpl. p. 15.*

In cortice *Juniperi* crescere dicitur.

6. *smaragdinum*, gregarium subcespitosum viride, stipite rigido persistente deorsum nigro, capitulo obovato pellucente demum deciduo. *Consp. fung. p. 355. t. I. f. I.*

"Ligna varia, praeprimis abietina, locis umbrosis humidis occupat, vere-aestate, magnitudine lineare. Stipes primo unicolor, dein nigrescit, apice vero nitorem smaragdinum conservat." *Consp.*

7. *turbinatum*, capitulo turbinato ochraceo, stipite cylindrico gelatinoso superne albo pulverulento, inferne glabro subpellucido di-

bus, sero autumnio. Parvum, $\frac{1}{2}$ lin. aut plus longum. Ab initio totum cylindricum, demum capitulum oritur stipite multo latius et pellucidum.

14. *bicolor*, minutum, capitulo subgloboso albido, stipite longo attenuato olivaceo-albido. *Syn. fung.* p. 632. *Consp. fung.* p. 353.

Crescit ad truncos, etiam in Agaricis corruptis. Capit. molle, primo globosum, albidum, demum dilute olivascens et subovatum fit. *Consp.*

15. *ventricosum*, gregarium, capitulo ovato subgloboso albido, primo diaphano, stipite subulato albo (in sterilibus bitrifido), deorsum ventricoso subfusco diaphano. *Schum. Saell.* p. 240.

In Dania, autumnio, in stercore bovino oritur.

16. *fimetarium*, confertum, capitulo subrotundo albido evanescente, stipite persistente rigido lutescente.

St. rubicundum β . *fimetarium*. *Consp. fung.* p. 354?

Inveni hanc speciem, initio Octobris, supra fimum Cervi *capreoli*, quod magna in copia obtegebat. Magnitudine vulgari; capitulum parvulum; guttulam nebulosam sistens, verum mox evanescit, ita ut stipites, qui crassitu-

situdine varii, nec non apice quasi tomentosi sunt, soli perdurent.

17. *murinum*, sparsum simplex, stipite stricto albo, capitulo globoso luteo. *Syn. fung. p. 201.* sub *Mucore.* †

Aestatis fine, in umbrosis, ad stercus murinum occurrit. Minutissimum, stipite crassiusculo suffultum. Capitula primo pellucida, demum turbida; sed an farcta, an cava? hinc vera peridia sint, nondum constat.

18. *equinum*, minutissimum piliforme pellucidum, capitulis subovalibus, ex albido sublateritiis, demum turbidis.

Ascophora Stilbum. Tode Fung. Mechl. p. 14. t. 3. f. 24?

Nascitur in fimo equino humido. Oculi inermi vix perceptibile, sub lente ut pili apparet. Stipes pellucidus.

19. *byssinum*, minutissimum confertum nitidum, capitulo primo rotundo, demum obovato stipiteque candidis. *Syn. fung. p. 680.*

In ramis, locis humidis dejectis, ut byssus albida apparet. Quoad *Consp. fung.* etiam in Agaricis putrefactis nascitur.

20. *piliforme*, confertum subfasciculare minutum, stipite tenuissimo nigro, capitulo subrotundo aquoso. *Annal. bot. l. c. Decand. Flor. Gall. 6. p. 16.*

ADMONITIO.

- Pag. 147. lin. ultima leg. fuscosecente-cinerea.
 — 149. — 18. leg. membranacea sebacea.
 — 151. — 1. loc. *leucocoma* leg. *leucoloma*.
 — 156. — 5. loc. Muscos leg. Muscae.
 — — — 5. ab inf. loc. subramosa leg. subcarnosa.
 — 163. — 7. ab inf. loc. fungus leg. fungum.
 — 172. — 4. ab inf. loc. *Bolt.* leg. *Roth*.
 — 173. — 7. loc. Mero leg. Mire, et videsis.
 — 190. — 15. loc. fuit leg. fit.
 — 192. — 11. leg. *Phacorhiza*.
 — 193. — 10. leg. *Sclerotio Semine*.
 — 198. — 7. post compressa *add.* A cl. Esenbeck ut
 species distincta habita est, saltem etc.
-

Species quorundam generum, hac in Sectione
 enumeratorum, jam cognitae sed fortuito
 omissae, aut sub impressionem noviter detectae,
 et praefatio, nec non terminologiae et iconum
 explicatio, cum sectione secunda, propediem edi-
 tura, sequentur.

25. *pubidum*, candidum, capitulo ovato, stipite inferne incrassato pubido. *Tod. l. c. p. 32. t. 3. f. 21.*

Repertum in fol. *Pini sylvestris*, Augusto, post pluvias.

26. *bulbosum*, capitulo albo sphaerico, stipite attenuato subflexuoso bulboso. *Tod. l. c. p. 10. t. II. f. 7. †*

In culmis, stipulis foliisque aridis superne tectis. Stipes desinit in bulbum hemisphaericum griseum. Capitulum pulviusculo seminali ochroleuco obducitur. *T.*

27. *leiopus*, clavatum sporangio (capitulo) subgloboso depresso roseo, in stipitem crassum laevem album transeunte. *Ehrenberg Sylvae Mycolog. p. 24.*

Prope Berolinum vegetat in stercore murino, et lectum, vere, in trunco muscoso. *E.*

28. *micans*, clavatum subcarnosum, micante-roseum, capitulo obovato, stipite brevi albicante. *Clavaria micans. Syn. fung. p. 604. Comment. de fung. Clavaef. p. 85.*

Clavaria Acrospermum. Hoffm. Flor. German. 2. t. 7. f. 2.

Fungillus pulchellus provenit, Aprili-Majo, in caulibus plantarum siccis. Submollis est (non coriaceus), lin. 1-2 longus. Stipes fere pellucidus.

Obs. In ejus capitulo seu clavula nullas observare potui thecas, hinc eum a *Clavariis* removi, nec vera *Stilbi* species videtur, interim tamen melius ob substantiam et habitum huc referri debere mihi videtur, quam ad *Sclerotia*, quae consistentia sunt dura, subcompacta.

29. *laetum*, gregarium, capitulo subrotundo saepe geminato laete rubro, stipite brevi dilutius albidio. †

Sclerotium laetum. *Ehrenb. l. c. p. 27. t. 3. f. 2. a. b.*

In caule plantae annuae lectum. Antecedenti simillimum. Structura est carnosocellulosa. *Ehr.* An varietas?

30? *aquigenum*, stipitatum persistens, capitulo globoso atro-purpureo. *Rebent. Flor. Neomarch. p. 382.*

In *Charae* vulgaris caulibus hinc inde dispersum. Aestate. Stipes lin. longus, dilute purpureus. Capitulum nitidum, massa subgelatinosa alba repletum! Exsiccatum collabescit. *Reb.* — An insectorum ovula?

Obs. De hujus generis sede systematica, auctores dissentiunt, ob conformitatem aliquam cum *Helotiis*, inter *Helvellas* illud adhuc servandum duxi.

ADMONITIO.

Cum auctor aliquas tantummodo priores et posteriores plagulas a vitiis typographicis (nonnullisque forsitan et scriptoriis) purgare potuisset, eorum plura sese irrepsisse dolendum est. Sed cum taedeo esset, omnia sphalmata indicare, notandum sufficit, quod litterae praecipue *vocales* commutatae sint, et quod *commata* nonnunquam aut plane omissa, aut aliter posita sint. Hinc sequentia, quae majoris momenti sunt, ab lectore ut benevole corrigantur, praecor.

- Pag. 26. lin. 5. ab infer. loc. 3. leg. *M.*
— 32. — 4. ab infer. loc. *verne* leg. *verae*.
— 56. — 11. ab infer. leg. *examinanda*.
— 55. *Obs. etc.* ad finem generis post descriptionem numeri 6. collocari debebat.
— 56. *lin. infima*, loc. sequente leg. antecedente.
— 82. lin. 10-11. leg. *nigri*. — *agglutinati*.
— 87. — 6. ab inf. post *strictis*. add. *Tab. VIII. f. 1*
et 2. et p. 88. post *radiantibus* add. *Tab. VIII. f. 3.*
— 110. — 14. *Descr.* lineam separatam incipere debebat.
— 112. — 8. loc. prima leg. *pruina*.
— 121. — 1. leg. hac modo nota.
— — 8. *varicosa* praeposito asterisco lineam separatam inchoare debebat.
— 126. — 6. ab inf. loc. *Pontuy* leg. *Pontivy*.
— 130. — 5. post *pallida*. add. *Fries. l. c.*
— 142. — 1. loc. *Hersi* leg. *Sterei*

ADMONITIO.

- Pag. 147. lin. ultima leg. *fucescente-cinerea*.
 — 149. — 18. leg. *membranacea sebacea*.
 — 151. — 1. loc. *leucocoma* leg. *leucoloma*.
 — 156. — 5. loc. *Muscos* leg. *Muscas*.
 — — — 5. ab inf. loc. *subramosa* leg. *subcarnosa*.
 — 163. — 7. ab inf. loc. *fungus* leg. *fungum*.
 — 172. — 4. ab inf. loc. *Bolt.* leg. *Roth*.
 — 173. — 7. loc. *Mere* leg. *Mire*, et *videsis*.
 — 190. — 15. loc. *fuit* leg. *fit*.
 — 192. — 11. leg. *Phacorhiza*.
 — 193. — 10. leg. *Sclerotio Semine*.
 — 198. — 7. post *compressa add.* A cl. *Esenbeck* ut
species distincta habita est, saltem etc.

Species quorundam generum, hac in Sectione enumeratorum, jam cognitae sed fortuito omissae, aut sub impressionem noviter detectae, et praeformatio, nec non terminologiae et iconum explicatio, cum sectione secunda, propediem editura, sequentur.


















Tab. V.






Tab. VI.









Tab. VIII.



I. Sturm sc.






1

1





Tab. XI.



Jac. Sturm sc.



Tab. XII.



Jac. Sturm sc.

